


Development of biotechnological tools for the *in vitro* propagation of elite trees for the Basque Country forest

Alejandra Rojas Vargas

2023



eman ta zabal zazu



Universidad
del País Vasco

Euskal Herriko
Unibertsitatea



**Development of biotechnological tools for
the *in vitro* propagation of elite trees
for the Basque Country forest**

Directoras

Paloma Moncaleán

Itziar A. Montalbán

Alejandra Rojas Vargas (2023)

ACTA DE GRADO DE DOCTOR O DOCTORA

ACTA DE DEFENSA DE TESIS DOCTORAL

DOCTORANDO/A DON/DÑA. Alejandra Rojas Vargas

TITULO DE LA TESIS: Development of biotechnological tools for the *in vitro* propagation of elite trees for the Basque Country forest

El Tribunal designado por la Comisión de Postgrado de la UPV/EHU para calificar la Tesis Doctoral arriba indicada y reunido en el día de la fecha, una vez efectuada la defensa por el/la doctorando/a y contestadas las objeciones y/o sugerencias que se le han formulado, ha otorgado por _____ la calificación de:

unanimidad ó mayoría

SOBRESALIENTE / NOTABLE / APROBADO / NO APTO

Idioma/s de defensa (en caso de más de un idioma, especificar porcentaje defendido en cada idioma):

Castellano _____

Euskera _____

Otros Idiomas (especificar cuál/cuales y porcentaje) _____

En _____ a _____ de _____ de _____

EL/LA PRESIDENTE/A,

EL/LA SECRETARIO/A,

Fdo.:

Fdo.:

Dr/a: _____

Dr/a: _____

VOCAL 1º,

VOCAL 2º,

VOCAL 3º,

Fdo.:

Fdo.:

Fdo.:

Dr/a: _____ Dr/a: _____ Dr/a: _____

EL/LA DOCTORANDO/A,

Fdo.: _____

A mis padres: Victor e Isabel
A mis hermanos: Cindy, Melissa y Felipe
A Julio e Ignacio

AGRADECIMIENTOS

Hace cuatro años iniciamos como familia un reto y aventura llamada doctorado. Dejamos nuestro país, familias, trabajos y zona de confort e iniciamos un viaje transatlántico. Hoy finaliza mi tesis y con ella una etapa en mi vida académica de un constante aprendizaje, e inicia una vuelta a casa.... y es el momento justo para agradecer a Dios e instituciones y personas que hicieron posible el desarrollo de mi tesis.

Doy gracias al Instituto de Investigación y Servicios Forestales (INISEFOR), Departamento de Junta de Becas y a la Universidad Nacional de Costa Rica por otorgarme una beca predoctoral y a Neiker-BRTA por el uso de instalaciones y recursos.

Gracias al financiamiento de los proyectos: MINECO (AGL2016-76143-C4-3R), MICINN (PID2020-112627RB-C32), CYTED (P117RT0522), MULTIFOREVER project, under the umbrella of ERA-NET Cofund ForestValue by ANR (FR), FNR (DE), MINCYT (AR), MINECO-AEI (ES), MMM (FI), VINNOVA (SE). ForestValue has received funding from the European Union's Horizon 2020 Research and Innovation Programme under grant agreement no. 773324 y al DECO (Basque government), que han permitido la realización de la tesis.

Quiero dar un agradecimiento muy especial a mis directoras, las doctoras Paloma Moncaleán e Itziar Montalbán. Mil gracias por aceptarme y permitirme ser parte de "Pino Team". Agradezco mucho toda la formación académica, confianza, tiempo, esfuerzo, dedicación, paciencia y consejos. Gracias por las oportunidades e incentivos brindados de crecer, conocer nuevos lugares y laboratorios para ampliar los conocimientos científicos. Las recordaré siempre.

Muchas gracias al personal administrativo y técnico de Neiker-BRTA por ser un apoyo en el día a día de mi tesis. Agradecimientos a mis inolvidables compañeros de laboratorio: Maiara, Ander y Cátia. También, doy gracias a mis compañeros becarios e Isi Guitiérrez por todos los momentos buenos y no tan buenos....por su paciencia en mis momentos de estrés.... Les compensaré a cada uno en Costa Rica.....

A los doctores Miroslav Strnad, Ondřej Novák y su equipo por la inolvidable estancia realizada en el Laboratory of Plant Growth Regulators, Palacký University, República Checa. Gracias al doctor Jorge Canhoto y especialmente a parte de su equipo João, Cátia, Mariana y Mónica Zuzarte por la estancia realizada en el Centre for Functional Ecology, Department of Life Sciences, University of Coimbra, Portugal. Gran parte del capítulo 3 es producto de esa estancia.

A mis padres Víctor e Isabel por su amor, apoyo incondicional, educación y motivación, mis palabras se quedan cortas para agradecerles. Mil gracias. A mis hermanos Cindy, Meli y Fe; agradecimientos por su amor y apoyo incondicional.

A mi esposo Julio, gracias infinitas por aceptar esta aventura juntos, por tu amor, paciencia, apoyo y noches de desvelo. Gracias totales !!!!!. A mi hijo Ignacio, mi motor....mi motivación. Quiero dar gracias a nuestras familias en Costa Rica y Guatemala por su apoyo. Gracias a Maiara, Diego y todos los amigos que hicieron de este doctorado un disfrute.

INDEX

ABSTRACT	3
RESUMEN	7
CHAPTER 1 INTRODUCTION	11
1. Cultivated forests in the Basque Country: current scenery.....	13
2. <i>Pinus radiata</i> D. Don characteristics.....	13
3. Current health situation of cultivated forest in the Basque Country.....	14
4. Possible alternative species to radiata pine for the Basque Country.....	15
4.1. <i>Cryptomeria japonica</i> (Thunb. ex L.f.) D. Don.....	15
4.2. <i>Pinus ponderosa</i> P. Lawson and C. Lawson.....	16
4.3. <i>Sequoia sempervirens</i> (D. Don) Endl.....	17
5. Biotechnology: an alternative for the forestry sector.....	18
6. Factors influencing the success of the micropropagation process.....	19
7. REFERENCES.....	21
CHAPTER 2 General Hypothesis and Objectives	31
CHAPTER 3 Influence of Physico-Chemical Factors on the Efficiency and Metabolite Profile of Adult <i>Pinus radiata</i> D. Don Bud Organogenesis	35
ABSTRACT	37
RESUMEN	37
1. INTRODUCTION.....	38
2. MATERIALS AND METHODS.....	39
2.1. Plant Material.....	39
2.2. Organogenic Process.....	41
2.2.1. Shoot Growth and Elongation.....	41
2.2.2. Root Induction and Acclimatization of Rooted Plants.....	41
2.3. Metabolite Extraction and Soluble Sugar and Amino acid Quantification.....	42
2.4. Data Collection and Statistical Analysis.....	43
3. RESULTS.....	43
3.1. Organogenic Process.....	43
3.2. Rooting Induction and Acclimatization of Rooted Plants.....	45
3.3. Metabolite Analysis.....	47
4. DISCUSSION.....	54
5. CONCLUSIONS.....	57
6. REFERENCES.....	57
CHAPTER 4 Adult trees <i>Cryptomeria japonica</i> (Thunb. ex L.f.) D. Don Micropropagation: Factors Involved in the Success of the Process	65
ABSTRACT	67
RESUMEN	67
1. INTRODUCTION.....	68
2. MATERIALS AND METHODS.....	69
2.1. Plant Material.....	69
2.2. Sterilization.....	70
2.3. Micropropagation Process.....	70

2.3.1.	Experiment 1	70
2.3.2.	Root Induction and Acclimatization of Rooted Plants	70
2.3.3.	Experiment 2	71
2.3.4.	Root Induction and Acclimatization of Rooted Plants	71
2.4.	Data collection and Statistical Analysis	71
2.4.1.	Experiment 1	71
2.4.2.	Experiment 2	72
3.	RESULTS	73
3.1.	Micropropagation Process	73
3.1.1.	Experiment 1	73
3.1.2.	Root Induction and Acclimatization of Rooted Plants	75
3.2.	Experiment 2	77
4.	DISCUSSION	78
5.	CONCLUSIONS	81
6.	REFERENCES	81
CHAPTER 5 Testing Explant Sources, Culture Media, and Light Conditions for the Improvement of Organogenesis in Pinus ponderosa P. Lawson and C. Lawson..... 87		
	ABSTRACT	89
	RESUMEN	89
1.	INTRODUCTION	90
2.	MATERIALS AND METHODS	91
2.1.	Plant Material	91
2.2.	Sterilization	91
2.3.	Organogenic Process	92
2.3.1.	Experiment 1	92
2.3.2.	Experiment 2	93
2.3.3.	Rooting and Adaptation	93
3.	Morphological Characterization of Needles	94
3.1.	Light Microscopy Analysis	94
3.2.	Scanning Electron Microscopy	94
4.	Data Collection and Statistical Analysis	94
4.1.	Experiment 1	94
4.2.	Experiment 2	95
5.	RESULTS	95
5.1.	Induction of Organogenesis	95
5.1.1.	Experiment 1	95
5.1.2.	Experiment 2	97
5.2.	Morphological Characterization of Needles	98
6.	DISCUSSION	102
7.	CONCLUSIONS	104
8.	REFERENCES	105
CHAPTER 6 Optimization of the micropropagation of elite adult trees of Sequoia sempervirens: forest species of interest in the Basque Country, Spain..... 111		
	ABSTRACT	113
	RESUMEN	113
1.	INTRODUCTION	115

2.	MATERIALS AND METHODS	115
2.1.	Plant material	115
2.2.	Shoot induction	115
2.3.	Shoot elongation	116
2.4.	Root induction and acclimatization of plants	116
2.5.	Data collection and statistical analysis	116
3.	RESULTS	117
3.1.	Shoot induction	117
3.2.	Root induction	120
3.3.	Acclimatization of rooted microplants	122
4.	DISCUSSION	123
5.	REFERENCES	125
	CHAPTER 7 GENERAL DISCUSSION	129
	CONCLUSIONS	139
	SUPPLEMENTARY MATERIALS	143
	CHAPTER 3 Influence of Physico-Chemical Factors on the Efficiency and Metabolite Profile of Adult <i>Pinus radiata</i> D. Don Bud Organogenesis	145
	CHAPTER 4 Adult trees <i>Cryptomeria japonica</i> (Thunb. ex L.f.) D. Don Micropropagation: Factors Involved in the Success of the Process	157
	CHAPTER 5 Testing Explant Sources, Culture Media, and Light Conditions for the Improvement of Organogenesis in <i>Pinus ponderosa</i> P. Lawson and C. Lawson	163

ABBREVIATIONS

2,4-D	2,4-dichlorophenoxyacetic acid
A	auxin type
AC	activated charcoal
AG	apical explants of 1.5 cm in length
ANOVA	analysis of variance
AP	apical explants of 1.0 cm length
ARN	ARN medium (Arnaud et al. 1993)
AS	apical explants <1.0 cm
AT	apical explants >1.5 cm
BA	6-benzyladenine
BG	basal explants of 1.5 cm length
BP	basal explants of 1.0 cm in length
BS	basal explants <1.0 cm
BT	basal explants >1.5 cm
C	cytokinin type
CK	cytokinin
CKs	cytokinins
DCR	DCR medium (Gupta and Durzan, 1985)
df	degrees of freedom
E	explant type
EFS	explants forming shoots
EM	elongation medium
FL	fluorescent light
FMOC	9-fluorenyl-methylchloroformate protecting group
HLP	half LP macronutrients medium
HPLC	high-performance liquid chromatography
IAA	indole-3-acetic acid
IBA	indole-3-butyric acid
IM	induction medium
KIN	kinetin
LED	light emitting diode
LEDs	light emitting diodes
LLR	length of the longest root

LP	LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)]
MS	MS medium (Murashige and Skoog, 1962)
m-T	meta-topolin
NAA	1-naphthaleneacetic acid
NR/E	number of roots per explant
ns	non significant
NS/E	number of shoots per explant
OPA	o-phthaldialdehyde
PGRs	plant growth regulators
PSR	percentage of shoots elongated sufficiently to be rooted
PVP	polyvinylpyrrolidone
QL	QL medium (Quoirin and Lepoivre, 1977)
REM	root expression medium
RI	root induction
RIM	root induction medium
S.E.	standard error
SE	somatic embryogenesis
SEM	scanning electron microscopy
SI	shoot induction
TDZ	thidiazuron

ABSTRACT

ABSTRACT

Located in the North of Spain, Basque Country has edaphoclimatic conditions that have allowed the establishment and development of an important forestry sector at European level. In this sense, 54.8% of the territory is dedicated to natural and cultivated forests. The most cultivated species in the Basque territory is *Pinus radiata* D. Don, whose first report of forest plantation establishment dates back to 1897. From that year to the present, its cultivation has increased in the Basque Country, reporting for the year 2022 the largest total area (25.9 %) and the highest forest productivity (80-85% of the annual cuts).

P. radiata has a natural distribution from the Central Coast of California to Mexico. The versatility and quality of the wood, together with the rapid growth of the species, has allowed its plantations to extend to countries such as Australia, Chile, New Zealand, South Africa and Spain. However, in recent years, *P. radiata* in the Basque Country has been affected by several diseases caused mainly by fungi: Pine canker caused by *Fusarium circinatum*, Dothistroma needle blight caused by *Dothistroma septosporum* and *Dothistroma pini*, and brown spot needle blight caused by *Lecanosticta acicula*. This last disease has affected 3,300 ha in Gipuzkoa (Basque Country). In addition, *P. radiata* plantations in the Basque Country have decreased from 123,921 ha (2016) to 102,488 ha (2022), a decrease that coincides with a historical outbreak of the needle blight during the years 2018-2019. For all of the abovementioned reasons, and together with the increase in the world population that causes a strong demand for timber and non-timber products, HAZI (www.hazi.eus) proposed a list of candidate species to be used, compatible with the demand of the forestry sector, as well as socially and culturally accepted.

Using the list of possible alternative species, the following were selected for this study: *Cryptomeria japonica* (Thunb. ex L.f.) D. Don, *Pinus ponderosa* P. Lawson and C. Lawson, *Sequoia sempervirens* (D. Don) Endl. and apparently tolerant genotypes of *P. radiata*. However, these species display desirable characteristics for clonal propagation at maturity, when traditional vegetative propagation and rooting are unfeasible. Therefore, the development of alternative propagation methods, such as *in vitro* propagation for these species is necessary. *In vitro* propagation through tissue culture can be performed mainly by organogenesis or somatic embryogenesis (SE); these techniques combined with traditional ones could be efficient for the vegetative propagation of the abovementioned alternative species. In addition, these methods allow to improve the quality, yield and health of commercially valuable forest species. In this context, organogenesis in species of the Pinaceae family is generally restricted to zygotic embryos, parts of them or young seedlings as initial source of explants and attempts to clone selected adult individuals have been feasible but have faced several difficulties. In this regard, in our laboratory at Neiker-BRTA, several *in vitro* propagation protocols have been successfully developed for *Pinus pinaster* Ait., *P. sylvestris* L., *P. pinea* L., *P. radiata* and *P. halepensis* Mill. among others.

Taking into account all this information, the general hypothesis of this work is that the modification of the physico-chemical environment in the micropropagation process can influence the regeneration efficiency of elite trees of forest species such as *C. japonica*, *P. ponderosa* and *S. sempervirens* to be considered as an alternative to *P. radiata* in the Basque Country. The second hypothesis to be corroborated is that adult *P. radiata* trees with apparent tolerance to pine needle blight can be used as a source of material for *in vitro* propagation.

In this context, the objective of the first chapter of this thesis was to analyze the influence of collection dates and different concentrations of 6-benzyladenine (BA) in the shoot induction phase, as well as the effect of different types of light on rooting. Also, the effect of these physico-chemical modifications in the content of amino acids and carbohydrates in the shoots developed *in vitro* was studied. Although reinvigorated shoots were obtained in both BA concentrations (22 or 44 μ M), the highest BA concentration promoted the best results in terms of shoot induction (SI) and number of shoots per explant (NS/E) when the

explants were collected in the first week of February. In addition, in shoots developed *in vitro*, significant differences were found in the levels of fructose, glucose and sucrose and in the threonine and tyrosine contents in the different phases of the rooting process.

The objective in the second chapter was to optimize the micropropagation method of *C. japonica* using adult trees as a source of plant material. In a first experiment, the effect of different types of initial explants and three basal culture media on SI was studied. In addition, two sucrose concentrations and two light treatments: light emitting diode (LED) versus fluorescent light (FL) were tested at rooting stage. In a second experiment, the effect of different plant growth regulators (PGRs) [BA, meta-topolin (m-T) and thidiazuron (TDZ)] on SI and subsequent phases of the organogenesis process was analysed. As main results, the highest number of shoots was obtained when using basal explants longer than 1.5 cm of length grown in the QL culture medium (Quoirin and Lepoivre, 1977). The shoots produced a higher number of roots under red LED lights. Finally, root induction (RI) was significantly higher in shoots previously induced with m-T.

The third chapter was focused on the analysis of the influence of different types of explants, basal culture media, cytokinins (CKs), auxins and light treatments on the *in vitro* propagation success in *P. ponderosa*. In this sense, whole zygotic embryos cultured in 44 μM BA performed better in terms of explant survival. For shoot organogenesis, whole zygotic embryos and LP with macronutrients halved LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] were selected. Regarding the light treatments applied, a significantly higher percentage of rootable shoots was obtained under blue LED light. However, the percentage of acclimatization was higher in shoots previously cultivated under FL. Furthermore, anatomical studies using optical and scanning electron microscopy (SEM) showed that light treatments promoted differences in anatomical aspects in *in vitro* shoots; needles of plantlets exposed to red and blue light emitted diodes (LEDs) showed less stomata compared with needles from plantlets exposed to FL.

The objective in the fourth chapter was to optimize the micropropagation of adult elite trees of *S. sempervirens*. To accomplish this objective, the effect of different CKs [BA, m-T and kinetin (KIN)], and four types of explants in SI were studied. At rooting stage, the effect of two types of auxins was evaluated: 1-naphthalene acetic acid (NAA), indole-3-butyric acid (IBA) and a mixture of both. The best induction rate was obtained when apical explants longer than 1.5 cm were induced with 4.4 μM BA. Regarding the rooting, shoots previously induced in 4.4 μM BA and rooted in a culture medium with 50 μM NAA showed the best result in terms of rooting percentage. However, acclimatization was better when explants had been cultivated with KIN and rooted in a culture medium with IBA.

This work has demonstrated that *in vitro* culture can be used as an alternative for the propagation of elite individuals of *C. japonica*, *P. ponderosa*, *P. radiata* and *S. sempervirens*. Furthermore, it has been evidenced that the *in vitro* propagation success in these species is influenced by genetic factors, collection dates and physico-chemical conditions of the culture environment.

RESUMEN

RESUMEN

Situado en el norte de España, el País Vasco cuenta con unas condiciones edafoclimáticas que han permitido el establecimiento y desarrollo de un importante sector forestal. En este sentido, el 54,8% del territorio está dedicado a bosques naturales y cultivados. La especie más cultivada en el territorio vasco es *Pinus radiata* D. Don, siendo el primer informe de su establecimiento del año 1897. Desde ese año hasta la actualidad su cultivo se ha incrementado en Euskadi, reportando para el año 2022 la mayor superficie total (25.9%) y la mayor productividad forestal (80-85% de las cortas anuales).

P. radiata tiene una distribución natural desde la costa central de California hasta México. La versatilidad y calidad de la madera unido al rápido crecimiento de la especie han permitido que sus plantaciones se extiendan en países como Australia, Chile, Nueva Zelanda, Sudáfrica y España. Sin embargo, en los últimos años, el *P. radiata* en el País Vasco se ha visto afectado por varias enfermedades causadas principalmente por hongos: Chancro resinoso del pino, causado por *Fusarium circinatum*; banda roja causada por *Dothistroma septosporum* and *Dothistroma pini* y banda marrón causado por *Lecanosticta acicula*. Esta última enfermedad ha ocasionado la pérdida de 3.300 ha en Gipuzkoa (País Vasco). Además, las plantaciones de *P. radiata* en Euskadi han disminuido de 123.921 ha (2016) a 102.488 ha (2022); disminución que coincide con un brote histórico de la enfermedad (bandas) durante los años 2018-2019. Todo esto unido al aumento de la población a nivel mundial provoca una fuerte demanda de productos maderables y no maderables. Por ello, HAZI (www.hazi.eus) creó una lista de especies alternativas para ser utilizadas en sustitución del *P. radiata*, compatibles con la demanda del sector forestal, así como social y culturalmente aceptadas.

En base a la lista de especies candidatas antes mencionada, se seleccionaron las siguientes para llevar a cabo nuestros estudios: *Cryptomeria japonica* (Thunb. ex L.f.) D. Don, *Pinus ponderosa* P. Lawson and C. Lawson, *Sequoia sempervirens* (D. Don) Endl. y genotipos aparentemente tolerantes de *P. radiata*. Sin embargo, las especies mencionadas muestran las características deseables cuando se ha producido el cambio de fase y no es posible su propagación mediante técnicas tradicionales.

Por lo tanto, es necesario el desarrollo de métodos alternativos como la propagación *in vitro*. La propagación *in vitro* a través del cultivo de tejidos se realiza principalmente mediante organogénesis o embriogénesis somática. Estos métodos permiten mejorar la calidad, el rendimiento y la sanidad de especies forestales comercialmente valiosas. En este contexto, la organogénesis en especies de la familia Pinaceae generalmente se restringe a embriones cigóticos, partes de ellos o plántulas jóvenes como fuente inicial de explantos; la clonación de individuos adultos seleccionados ha sido posible pero presenta ciertas dificultades. En este sentido, en nuestro laboratorio en Neiker-BRTA se han desarrollado de manera exitosa protocolos de propagación *in vitro* en *Pinus pinaster* Ait., *P. sylvestris* L., *P. pinea* L., *P. radiata* y *P. halepensis* Mill. entre otros.

En base a la información antes mencionada, la hipótesis general de este trabajo es que la modificación del entorno físico-químico en el proceso de micropropagación puede influir en la eficiencia de regeneración de plantas élite de especies forestales como *C. japonica*, *P. ponderosa* y *S. sempervirens* para ser consideradas como alternativa a *P. radiata* en el País Vasco. La segunda hipótesis a ser corroborada es que los árboles adultos de *P. radiata* con aparente tolerancia a la enfermedad de las bandas pueden usarse como fuente de material para la propagación *in vitro*.

En este contexto, el objetivo del primer capítulo de la tesis fue analizar la influencia de las fechas de recolección y diferentes concentraciones de BA en la fase de inducción de brotes, así como el efecto de diferentes tipos de luz sobre el éxito de la inducción de raíces. A su vez, se estudió el efecto de los factores físico-químicos antes mencionados sobre el contenido de aminoácidos y carbohidratos en los brotes desarrollados *in vitro*. Aunque se obtuvieron brotes revigorizados en ambas concentraciones de BA (22 o 44 μ M), la BA a mayor concentración mostró los mejores resultados en términos de inducción y número de brotes por explanto cuando estos se recolectaron en la primera semana de febrero. Además, se encontraron diferencias significativas en el

perfil de fructosa, glucosa y sacarosa y en el contenido de treonina y tirosina de los brotes desarrollados *in vitro* en las diferentes fases del proceso de enraizamiento.

El segundo capítulo tuvo como objetivo optimizar el método de micropropagación de *C. japonica* utilizando árboles adultos como fuente de material vegetal. En un primer experimento, se estudió el efecto de diferentes tipos de explantos iniciales y tres medios de cultivo basales en la inducción de brotes. Además, se probaron dos concentraciones de sacarosa y dos tratamientos de luz (LEDs versus luces fluorescentes) en la etapa de enraizamiento. En un segundo experimento, se analizaron los efectos de diferentes reguladores del crecimiento (BA, m-T y TDZ) en la inducción de brotes y las fases posteriores del proceso de organogénesis. Como principales resultados, el mayor número de brotes se obtuvo cuando se utilizaron explantos basales con una longitud superior a 1,5 cm cultivados en medio QL (Quoirin y Lepoivre, 1977). Los brotes obtenidos produjeron un mayor número de raíces cuando se cultivaron bajo luz LED roja. Finalmente, la inducción de raíces fue significativamente mayor en los brotes previamente inducidos con m-T

El objetivo del tercer capítulo fue analizar la influencia de diferentes tipos de explantos, medios de cultivo basales, citoquininas, auxinas y tratamientos de luz en el éxito de la propagación *in vitro* en *P. ponderosa*. En este sentido, los embriones cigóticos completos y cultivados en 44 μM BA mostraron los mejores resultados en términos de supervivencia de los explantos. Para la organogénesis de brotes, se seleccionaron los embriones cigóticos completos y el medio de cultivo LP [Quoirin y Lepoivre (1977) modificado por Aitken-Christie et al. (1988)] con sus macronutrientes reducidos a la mitad. En cuanto a los tratamientos de luz aplicados, se observó un porcentaje significativamente mayor de brotes susceptibles de ser enraizados cuando estos crecieron bajo luz LED azul. Sin embargo, el porcentaje de aclimatación fue mayor en los brotes previamente cultivados bajo luces fluorescentes.

Por otra parte, los estudios anatómicos mediante microscopía óptica y microscopía electrónica de barrido mostraron que los tratamientos con luz promovieron diferencias en aspectos anatómicos en brotes *in vitro*; así, las acículas de plántulas expuestas a LEDs rojos y azules mostraron menos estomas en comparación con las acículas de plántulas expuestas a luz fluorescente.

El objetivo en el cuarto capítulo fue optimizar la micropropagación de árboles élite adultos de *S. sempervirens*. Para lograr este objetivo se estudió el efecto de diferentes tipos de citoquininas (BA, m-T y KIN), y cuatro tipos de explantos en la etapa de inducción de brotes. Además, se evaluó el efecto de dos tipos de auxinas: NAA, IBA y una mezcla de ambas. La mejor tasa de inducción de brotes se obtuvo cuando se utilizaron explantos apicales con una longitud superior a 1,5 cm inducidos con 4,4 μM BA. El mejor resultado en términos de inducción de raíces se obtuvo cuando los brotes previamente inducidos en 4,4 μM BA se cultivaron en un medio de cultivo suplementado con 50 μM de NAA. Sin embargo, la aclimatación fue mejor cuando los explantos cultivados con KIN y enraizados en un medio de cultivo con IBA.

Este trabajo ha demostrado que el cultivo de *in vitro* puede ser utilizado como un método alternativo de propagación de individuos élites de *C. japonica*, *P. ponderosa*, *P. radiata* and *S. sempervirens*. Además, se ha evidenciado que el éxito de la propagación *in vitro* en estas especies está influido por factores genéticos, fechas de recolección y condiciones físico-químicas del ambiente de cultivo.

CHAPTER 1
INTRODUCTION

1. Cultivated forests in the Basque Country: current scenery

The Basque Country have an extension of 7,226 km² and it is located in the North of Spain (42°28'-43°27' N; 1°44'W-3°27'W). Historically, the edaphoclimatic conditions of the Basque Country have led to an important silvicultural development (Ibarrondo, 2008). In this context, the 54.8% (396, 191 ha) of the territory is dedicated to natural and cultivated forests, being an important surface in the European region (HAZI, 2022). Therefore, in the Basque Country, the forestry industry plays an important role at the economic, social and cultural level.

According with the last forest map of the Basque Country (2022), the wooded forest area is composed mainly of the following species: *Pinus radiata* (102,488 ha), *Fagus sylvatica* (55,291 ha), Atlantic mixed forest (38,138 ha), *Quercus faginea* (26,892 ha), *Q. ilex* (26,757 ha), *Eucalyptus* spp. (25,316 ha), *Q. robur* together *Q. petraea* (17,488 ha), *P. sylvestris* (17,493 ha), *P. nigra* (13,857 ha) and *Q. pyrenaica* (13,258 ha) (HAZI, 2022).

In Spain, the largest plantations of *P. radiata* are in the Basque Country (HAZI, 2022). The first plantation of *P. radiata* was established in 1897 (Puerta et al., 2004). From then until nowadays, its cultivation has increased in the Basque Country, showing in 2022 the largest total wooded area (25.9%) and highest forest productivity (80-85% of the annual cuts) (HAZI, 2022).

2. *Pinus radiata* D. Don characteristics

Pinus radiata D. Don, commonly known as radiata pine or Monterey Pine, is naturally distributed from the Central Coast of California to Mexico. Due to the versatility, quality of the timber and rapid growth of the species, its plantations have been extended along Australia, Chile, New Zealand and South Africa (Martínez-Álvarez et al., 2014; Escandón et al., 2017; Farjon, 2018; Castander-Olarieta et al., 2019).

Radiata pine is an evergreen tree (Figure 1A), 30 m tall, trunk to 0.9 m diameter and leaves (needles) in fascicles of 2 or 3 needles (8- 20 cm long) with dark green color (Figure1B). The cones (7-15 cm long) have ellipsoid-cylindric shape and need two years for maturing, the seeds are characterized by two parts, the body (0.6 cm long) of dark brown color and the wing (2-3 cm long) (Bussmann et al., 2020).



Figure 1. (A) *Pinus radiata* D. Don tree in a seed orchard established by Neiker-BRTA in Arkaute (Álava, Spain) and (B) apical buds and leaves (needles).

Radiata pine natural populations grow in climatic conditions characterized by monthly temperatures means ranging from 9 °C to 11 °C in winter, 16 °C to 18 °C in summer and annual precipitations ranging from 380 to 890 mm, with dry summers and snowless winters (Mead, 2013).

3. Current health situation of cultivated forest in the Basque Country

During the last years, *P. radiata* in the Basque Country has been affected by several diseases caused mainly by fungi. This situation has provoked a deep reduction of the survival, vigor and yield of cultivated forests (Elvira-Recuenco et al., 2020). Among the fungi that have caused the most damage to Basque Forest systems are the following:

a.-Pine pitch canker (PPC), caused by *Fusarium circinatum* Nirenberg and O'Donnell, provoked an affectation of 353,65 ha of radiata pine plantations during the years 2006-2011 in Gipuzkoa, Basque Country (Cantero, 2019).

b.-Dothistroma needle blight (DNB), caused by *Dothistroma septosporum* (Dorog.) Morelet and *D. pini* Hulbary.

c.-Brown spot needle blight caused by *Lecanosticta acicula* (Thümen) Sydow. According to Cantero (2019), during 2011-2012 an affectation of 3,300 ha of radiata pine plantations caused by this disease was reported in Gipuzkoa.

d.-*Mycosphaerella dearnessii* teleomorph of *L. acicula* also has been reported in *P. radiata* (Elvira-Recuenco et al., 2020; Mesanza et al., 2021a, 2021b).

In the last years (2018-2019), plantations of radiata pine in the North of Spain were affected by a historical outbreak of needle blight disease. In Basque Forest systems this disease was mainly caused by *D. septosporum* and *L. acicula* (Mesanza et al., 2021c), provoked an affectation of 53,227 ha in the abovementioned period (Cantero, 2019).

In the Basque country, the presence of *L. acicula* dates back to 1942, but for unknown scientific reasons, the disease has expanded quickly in the last years. In this context, the plantations of radiata pine in the Basque Country have decreased from 123,921 ha (2016) to 102,488 ha (2022), a decrease that coincides with the historical outbreak of this type of needle blight disease (HAZI, 2022). Also, several reports have mentioned problems in the commercial viability of these plantations. In this sense, it has been

described that climate change is the responsible of the emergence of needle blight disease (Jankovský et al., 2009; Allen et al., 2010; Woods et al., 2016; Castander-Olarieta et al., 2019).

The increase of population worldwide provokes a strong demand of wood and non-wood products (Farjon, 2018; Bueno et al., 2021). For this reason, it is urgent to find alternative species that could be used in the Basque Forest plantations and that are compatible with the demands of the industrial sector as well as being socially and culturally accepted.

At this point, a great effort has been made to find out which species could be planted in the areas where the radiata pine is being eliminated, which are resistant to the diseases mentioned and grow rapidly in our edaphoclimatic conditions. In recent years, forestry sector and HAZI (www.hazi.eus) proposed a list of alternative species to be used (Cantero, 2019): *Larix leptolepis*, *L. decidua*, *Pseudotsuga menziesii*, *Cryptomeria japonica*, *Sequoiadendron giganteum*, *Sequoia sempervirens*, *Picea abies*, *P. omorika*, *P. pungens*, *P. sitchensis*, *P. shrenkiana*.

4. Possible alternative species to radiata pine for the Basque Country

To carry out our experiments, three alternative species were chosen to work with: *Cryptomeria japonica* (Thunb. ex L.f.) D. Don, *Pinus ponderosa* P. Lawson and C. Lawson and *Sequoia sempervirens* (D. Don). Endl. Furthermore, *P. radiata* D. Don apparently tolerant genotypes were used for *in vitro* propagation experiments.

4.1. *Cryptomeria japonica* (Thunb. ex L.f.) D. Don

Cryptomeria japonica (Thunb. ex L.f.) D. Don (Figure 2), is monoecious conifer, belonging to the Cupressaceae family (20 genera with 125 species) and Taxodioidae subfamily (Onuma et al., 2023). This species also known as Japanese cedar or Sugi occurs naturally throughout East Asia (Taniguchi et al., 2020). According with Tsumura (2023), *C. japonica* forest area is naturally distributed and adapted to different ecosystems where the climatic conditions are characterized by an annual precipitation over 1500 mm and local temperatures ranging from 8.4 °C to 13.9 °C with cold winters. Sugi is an evergreen tree that can up to 50 m tall, it shows slender leaves, awl-shaped, bluntly pointed; male cones with axillary and lateral female globular cones with wedgeshaped scales (Farjon, 1999).



Figure 2. *Cryptomeria japonica* (Thunb. ex L.f.) D. Don adult tree in Arkaute, Álava, Spain.

In Japan, Sugi is an important forestry species, it constitutes around 44% of forest plantations, its timber is used for wood construction or for obtaining biomass (Taniguchi et al., 2020; Onuma et al., 2023; Tsumura, 2023). In this sense, Japanese Cedar tree has been introduced in different countries as an alternative species due to its valuable timber and fast-growing (Page, 1990; Shimizu and Maiochi, 2007; Koguta et al., 2017).

In the Paraná region of Brazil, it has been introduced for the last 40 years, and in other countries such Russia, England or Indonesia was introduced in the nineteenth century (Farjon, 1999; Shimizu and Maiochi, 2007; Koguta et al., 2017). In the Basque Country, *C. japonica* plantations occupied an area of 3,372 ha in 2022 (HAZI, 2022).

4.2. *Pinus ponderosa* P. Lawson and C. Lawson

Pinus ponderosa P. Lawson and C. Lawson belongs to the Pinaceae family, where the genus *Pinus* occupies the first place among all conifers in economic importance (Farjon, 2018; Charpin et al., 2022; Neale and Wheeler, 2019).

P. ponderosa (Figure 3) also known as ponderosa pine or western yellow pine is native to the United States, Southern Canada and Northern Mexico (Ellis and Bilderback, 1991; Raish et al., 1997). This species is the largest of the western pine species and is a major lumber tree in Western North America (Fitzgerald, 2005).



Figure 3. *Pinus ponderosa* P. Lawson and C. Lawson in a seed orchard established by Instituto Nacional de Tecnología Agropecuaria in Trevelin, Argentina.

Ponderosa pine is characterized for its protective outer corky bark and long needles (in fascicles of two or three needles) with high moisture content surrounding the terminal buds (Hall, 1980; Miller, 2000; Fitzgerald, 2005). Due to these adaptations that promote its survival, is considered one the most fires-resistant conifers in the West of the United States (Hall, 1980; Miller, 2000; Fitzgerald, 2005).

P. ponderosa forests are important in ecology terms, since they are wildlife habitats for animals and plants, growing in sites characterized by elevations ranging from sea level to 3,281 m, predominantly dry and moist. The seedling survival is affected by water availability and soil conditions; therefore, in the current climate change scenery, the development of new propagation technologies in the species is justified. They are also part of recreational, an esthetic and cultural values; the species itself

constitute a great economic asset due to the value of its timber (Zhang et al., 2022). In this respect, ponderosa pine cultivated forests have been established in California with lumber purposes on private lands, while in countries as Chile and Argentina has been introduced as a promising species to optimize and diversify the wood industry (Zhang et al., 2019; Espinoza et al., 2021; Solans et al., 2021).

4.3. *Sequoia sempervirens* (D. Don) Endl.

Sequoia sempervirens (D. Don) Endl. trees belong to the family Taxodiaceae (nine genera with 16 species), it shows an extreme size (400 Mg/ha biomass) and can live up 2,500 years (Figure 4) (Burns et al., 2018; Sillett et al., 2020). *S. sempervirens* is commonly known as coastal redwood and naturally occurs in the West Coast of North America; it is also cultivated in countries such as Romania as ornamental in parks and botanical gardens (Stoiculescu et al., 2009). Burns et al. (2018) mentioned that *Sequoia* has approximately 5% of primary forests remaining located in the native distribution of the genera, 740 km from Southwestern Oregon to central California, in sites below 1000 m elevation and north-south latitudinal gradient (Sillett et al., 2022).



Figure 4. *Sequoia sempervirens* (D. Don.) Endl. adult tree in Vitoria-Gasteiz, Álava, Spain.

Redwood is a shade-tolerant evergreen tree with fast growth rates that can reach up to 116 m in height. It has linear leaves (0.3-1.0 cm long), with 6-12 leaves arranged per centimeter along the axes and elliptical female cones (1.1-1.7 cm long) (Ma et al., 2005; Sillett et al., 2020; Watt et al., 2021).

Coastal redwood is an important species, due to the economic value of its wood (Laslo et al., 2015; Sillett et al., 2020). Several countries such as France, Chile, New Zealand and Germany have established *S. sempervirens* plantations with timber production purpose (Arnaud et al., 1993; Ibañez et al., 2009; Rapley, 2018; Breidenbach et al., 2020). In addition, in the Basque Country, it occupies an area of 1,586 ha (HAZI, 2022).

5. Biotechnology: an alternative for the forestry sector

In the last years the climate change, the increase of population and their demand over forestry products leads to find elite trees with higher productivity to be propagated for afforestation purposes.

Because of the often-poor correlation between juvenile and mature traits in conifers, proper testing in breeding programs generally requires at least about one-third to one-half the rotation age (Weng et al., 2008). This, coupled with the fact that conifer stem cuttings suitable for rooting are generally available only from juvenile plants, has led to limitations in traditional breeding techniques. Conifer trees usually develop the characteristics to determine their suitability for clonal planting after phase-change (during maturity), a developmental moment when rooting of cuttings becomes unfeasible (Bonga, 2015). Moreover, the intrinsic heterozygosity found in conifers makes difficult to fix desirable alleles by conventional techniques (Williams and Savolainen, 1996). Consequently, traditional breeding techniques should be combined with new biotechnological tools (forest biotechnology) such as *in vitro* tissue culture to effectively improve the performance, quality and health of commercially valuable forest species (Giri et al., 2004).

The generation of new plants *in vitro* can be achieved by three major pathways of micropropagation: 1) enhance of axillary bud break, 2) organogenesis and 3) SE (Loberant and Altman, 2010). In this way, the first uses pre-existing meristems in shoot tips and leaves and requires elongation and grow stage of the shoot. The axillary bud break is considered the simplest technique of micropropagation and the regenerated plants present the highest rates of genetic stability (Loberant and Altman, 2010; De Oliveira et al., 2012). In regard to organogenesis, it involves the production of adventitious buds (shoots or roots) directly (without callus formation) or indirectly via callus, where the regenerated plant are finally rooted (Phillips, 2004; Loberant and Altman, 2010; Loyola- Vargas and Ochoa-Alejo, 2012). Finally, SE defined as embryo regeneration from cells or tissues directly (without callus formation) or indirectly via callus (George, 2008).

Organogenesis has successfully employed to propagate a wide variety of angiosperm species with high economic importance, such as *Eucalyptus* and *Populus*, and has also shown great potential with certain conifer species, including *P. radiata* (Aitken-Christie et al., 1988; Montalbán et al., 2011). However, in the Pinaceae family this technique is restricted to zygotic embryos, parts of them, or young seedlings as explant sources, and attempts to clone selected adult trees, albeit feasible, have not shown the desired results. In our laboratory, organogenesis from vegetative buds of adult trees from various *Pinus* spp. has been carried out (De Diego et al., 2008; Cortizo et al., 2009; De Diego et al., 2010; Montalbán et al., 2013).

Micropropagation also known as *in vitro* culture is a procedure involving the manipulation of small plant parts (explants), tissues, organs or cells isolated from a mother plant that produces or regenerates plantlets under aseptic conditions on culture media and controlled environment (George, 2008; Loyola- Vargas and Ochoa-Alejo, 2012;). This procedure is based on the totipotency of plant cells which is defined as the ability of a single plant cell to regenerate a complete and functional plant (Haberlandt, 1902).

In general, organogenesis needs four stages for the success of the process: 1) establishment of the culture under *in vitro* conditions, 2) SI or multiplication, 3) rooting and 4) acclimatization to *ex vitro* conditions (George and Debergh, 2008; Dobránszki and Da Silva, 2010). The first stage starts with the selection of the explants from the mother plant, then surface sterilization of the explants and finally the culture on the culture media (George, 2008; Loberant and Altman, 2010; Loyola- Vargas and Ochoa-Alejo, 2012).

After several subcultures *in vitro* plants are rooted, this stage involves media and environmental changes as the addition or increase of auxins, reduced sugar levels, or modifications in light quality (Loberant and Altman, 2010). Finally, they are transferred from *in vitro* to *ex vitro* conditions to be acclimatized (George and Debergh, 2008).

The micropropagation technique compared to conventional propagation enables propagate independent on the season, high coefficient of multiplication in short time, aseptic cultures or disease-free cultures, fast propagation is a valuable technique for rare genotypes, sterile seeds or for the multiplication of species with economic or ecological importance (Tsubomura and Taniguchi, 2008; Valverde-Cerdas et al., 2008; Dobránszki and Da Silva, 2010; Sahu and Sahu, 2013). Although Bonga (2015) explained that conventional propagation will always be cheaper compared to biotechnological techniques, the cost is outweighed when, through micropropagation, plants with a genetic gain such as disease resistance or stem quality are obtained.

In this context, Neiker-BRTA tissue culture laboratory has developed several successful protocols for adult pine species: *Pinus pinaster* Ait. (De Diego et al., 2008), *P. pinea* L. (Cortizo et al., 2009), *P. sylvestris* L. (De Diego et al., 2010), *P. radiata* (Montalbán et al., 2013) and *P. halepensis* (Pereira et al., 2021).

6. Factors influencing the success of the micropropagation process

Plant tissue culture was developed more than half a century ago, but nowadays scientists use these biotechnological techniques not only to propagate plants, but to study physiological, morphological, biochemical and molecular aspects related to plant development, as these aspects are implicated in tissue culture success (Neelakandan and Wang, 2012).

It is generally known that the genotype, the source and explant type and the physico-chemical environment affect the success of *in vitro* morphogenic processes (Phillips and Garda, 2019). In this context, several works have demonstrated that the genotype has a significant role in the reliability and repeatability of a micropropagation protocol, since a different organogenic response can be obtained depending on the genotype (Coleman and Ernst, 1989; Meneguzzi et al., 2019). For example, in previous studies with *P. radiata*, *P. taeda* and *P. halepensis* a different organogenic response depending on the genotype was reported (Montalbán et al., 2013; C  zar et al., 2015; Pereira et al., 2021).

The size of the explants may play an important role in *in vitro* propagation (Da Silva et al., 2016; Bonga, 2017). Several studies have demonstrated that the explant size can affect axillary bud multiplication, due to larger explants having more endogenous hormones and nutrients to maintain the culture (Renau-Morata et al., 2005; George and Debergh, 2008; Desai et al., 2015). Furthermore, bigger explants from parts of shoot apex or stems with lateral buds show advantages over smaller explants for *in vitro* proliferation (George and Debergh, 2008). In this sense, an *in vitro* protocol was obtained in *S. sempevirens* using explants bigger than 1.5 cm in length (Clapa et al., 2010; Meneguzzi et al., 2020). On the contrary, explants with smaller size can perform better because a lower surface favours sterilization (George and Debergh, 2008; Sathyagowri and Seran, 2011). For example, the best survival response was observed in *Cedrus libani* (Renau-Morata et al., 2005) and *Taxus mairei* (Chang et al., 2001) explants shorter than 1 cm.

Physical factors such as light quality and intensity, temperature, humidity gaseous environment are manipulated in different ways to induce and optimize organogenesis or SE (Phillips, 2004; Kumar and Reddy, 2011). In this context, light quantity and spectral quality is one of the most important factors controlling plant growth and development and has effects on plant photosynthesis and photomorphogenesis (Fang and Jao, 2000; Alallaq et al., 2020). According to George and Davies (2008), the *in vitro* plant material can be affected during its growth and morphogenesis due to changes in the photo flux, the day length also called as photoperiod and the wavelength.

Generally, in most of plant tissue culture laboratories the use of tubular FL with light intensity between 25 and 150 $\mu\text{molm}^{-2}\text{s}^{-1}$ with 16 h photoperiod (8 h dark) per day is commonly used, depending on the species requirements (Larraburu et al., 2018; Phillips and Garda, 2019). However, FL increase costs due to high power consumption and limited efficiency, and culture rooms need to

remove the heat emitted by using air conditioners (Kulus and Woźny, 2020). In the last years, LEDs (red, blue, white and far-red) are recommended due their low power consumption, smaller size, durability, negligible heat production and the possibility to fix specific wavelengths, giving them advantages over FL (Fang and Jao, 2000).

Several studies reported that LEDs showed positive effects in *in vitro* propagation compared to conventional lighting (Gupta and Jatothu, 2013; Chen et al., 2020). For instance, *in vitro* shoots of *Gerbera jamesonii* growing under red LEDs displayed the greatest elongation rates (Pawłowska et al., 2018). A similar tendency was reported by Chen et al. (2020), where the highest stem length was observed in *in vitro* potato plantlets under red LEDs. Additionally, *in vitro* shoot regeneration was promoted in *Curculigo orchoides* and *Zingiber officinale* cultures exposed to blue LEDs (Gupta and Sahoo, 2015; Gnasekaran et al., 2021). Furthermore, *in vitro* explants of *Cucumis metuliferus* exposed to white LEDs and a mixed circuit of blue and red LEDs obtained higher RI rates (Lai et al., 2022). Also, *in vitro* shoots of *Cedrela odorata* growing under white LEDs showed a higher number of roots formed (Dos Santos et al., 2022).

Chemical factors including basal medium composition (for example, organic or inorganic nitrogen sources, pH, carbohydrate sources and concentrations) and PGRs (types and concentrations) are also factors involved in the induction and optimization of micropropagation success through organogenesis or SE (Phillips, 2004). The basal media are generally composed of macronutrient salts (N,P, K, Ca, Mg and S), micronutrient salts (Fe, B, Mn, Mo, Cu, Co and I), vitamins, PGRs (auxins, CKs, gibberelins, abscisic acid, etc.) and carbohydrates (Phillips and Garda, 2019). The basal medium supplies all the necessary mineral nutrients for the plant tissue growth and the basal medium composition and its modifications such as reducing macronutrients to half-strength, can affect the success of the micropropagation (Phillips and Garda, 2019). For example, in *P. pinea* differences in shoot proliferation were obtained when different culture media were tested (Sul and Korban, 2004). Likewise, several studies in *Pinus* species such as *P. halepensis*, *P. pinaster* and *P. ayacahuite* reported that salt concentrations in the basal medium influenced the *in vitro* morphogenesis response (Lambardi et al., 1993; Saborio et al., 1997; De Diego et al., 2011).

Carbohydrates in plants have an important role in the synthesis pathway of many compounds (Gibson, 2000; Calamar and De Klerk, 2002). Generally, sucrose is commonly used in micropropagation because it is cheap, relatively stable to autoclaving, readily assimilated by plants and it is the main sugar translocated in the phloem of several plants (Ragonezi et al., 2010; Kumar and Reddy, 2011). The carbohydrates are involved in the growth of adventitious roots in shoots and supply the energy necessary for root development (Corrêa et al., 2005). However, during the root expression stage, sucrose at high concentrations can have a negative effect (Ragonezi et al., 2010). In this context, in *Metroxylon sagu* a higher rooting response was obtained when sucrose 3% (w/v) was added to the medium instead of 4.5% and 6.0% (w/v) (Sumaryono et al., 2012).

As it was mentioned before, the basal medium can be supplemented with different PGRs depending on the morphogenic response to be induced (Phillips and Garda, 2019). According with van Staden et al. (2008), the balance and interaction between the growth regulators endogenously synthesized, and the PGRs added to the basal medium regulate the morphogenesis *in vitro*. Among all PGRs, auxins and CKs are the most used (Phillips and Garda, 2019).

The most common auxins used are NAA and IBA, followed by 2,4-dichlorophenoxyacetic acid (2,4-D) and indole-3-acetic acid (IAA). Auxins induce rooting and alone or in combination with CKs promote SE induction. For example, in *Pinus koraiensis* (Liang et al., 2022) the highest rooting rates were observed in root induction medium (RIM) supplemented with NAA and IBA. Moreover, different auxin/cytokinin (CK) ratios have been successfully applied for SE induction in *Pinus* (Montalbán et al., 2016).

Regarding CKs, they are derived from adenine (aminopurine) and BA is the most used, followed by KIN, TDZ and 2-isopentenyladenine (2-IP). The involvement of CKs in the promotion of *in vitro* shoot formation is due to their effect in plant cell

division stimulation; morphogenesis, organogenesis and senescence regulation (Bairu et al., 2007; De Diego et al., 2011; Aremu et al., 2012). For example, in *P. halepensis*, BA at 44 μ M provided a satisfactory SI response (Pereira et al., 2021). Furthermore, *P. pinaster* cotyledons cultured in the presence of 10 μ M BA showed a high organogenic response (Álvarez et al., 2009). A new class of aromatic CKs called topolins have been used in plant tissue culture as an alternative to conventional CKs (Aremu et al., 2012). The structure of topolins differs from that of BA in the presence of a hydroxyl group on the benzyl ring; and they have been used in *in vitro* cultures since topolins are involved in the improvement of parameters such as proliferation rates, RI and alleviation of physiological disorders (Werbrouck et al., 1996). For example, in *P. sylvestris* (De Diego et al., 2010) and *P. pinaster* (De Diego et al., 2011) a higher SI response was obtained when m-T [6-(3-hydroxybenzylamino) purine] was used.

The approaches related to environment modifications for the improvement of micropropagation procedures can be combined with micromorphological and metabolomic analysis to understand some of the mechanisms leading to their success. Together with the implementation of advanced micropropagation techniques, these strategies could contribute to achieve efficient propagation of elite trees of alternative species and meet the current necessities derived from current climate change.

7. REFERENCES

- Aitken-Christie, J.; Singh, A.P.; Davis, H. Multiplication of meristematic tissue: A new tissue culture system for radiata pine. In *Genetic Manipulation of Woody Plants*; Hanover, J.W., Keathley, D.E., Eds.; Plenum Publishing Corp: New York, NY, USA, 1988; pp. 413–432. https://doi.org/10.1007/978-1-4613-1661-9_26.
- Alallaq, S.; Ranjan, A.; Brunoni, F.; Novák, O.; Lakehal, A.; Bellini, C. Red light controls adventitious root regeneration by modulating hormone homeostasis in *Picea abies* seedlings. *Front. Plant Sci.* **2020**, *11*, 586140. <https://doi.org/10.3389/fpls.2020.586140>.
- Allen, C.D.; Macalady, A.K.; Chenchouni, H.; Bachelet, D.; McDowell, N.; Vennetier, M.; Kitzberger, T.; Rigling, A.; Breshears, D.D.; Hogg (Ted), E.H.; et al. A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *For. Ecol. Manag.* **2010**, *259*, 660–684. <https://doi.org/10.1016/j.foreco.2009.09.001>.
- Álvarez, J.M.; Majada, J.; Ordás, R.J. An improved micropropagation protocol for maritime pine (*Pinus pinaster* Ait.) isolated cotyledons. *Forestry* **2009**, *82*, 175–184. <https://doi.org/10.1093/forestry/cpn052>.
- Aremu, A.O.; Bairu, M.W.; Doležal, K.; Finnie, J.F.; Van Staden, J. Topolins: A panacea to plant tissue culture challenges? *Plant Cell Tissue Organ Cult.* **2012**, *108*, 1–16. <https://doi.org/10.1007/s11240-011-0007-7>.
- Arnaud, Y.; Franclet, A.; Tranvan, H.; Jacques, M. Micropropagation and rejuvenation of *Sequoia sempervirens* (Lamb) Endl: a review. *Ann. des Sci. For.* **1993**, *50*, 273–295. <https://hal.archives-ouvertes.fr/hal-00882846>.
- Bairu, M.W.; Stirk, W.A.; Dolezal, K.; Van Staden, J. Optimizing the micropropagation protocol for the endangered *Aloe polyphylla*: Can meta-topolin and its derivatives serve as replacement for benzyladenine and zeatin? *Plant Cell Tissue Organ Cult.* **2007**, *90*, 15–23. <https://doi.org/10.1007/s11240-007-9233-4>.
- Bonga, J. M. A comparative evaluation of the application of somatic embryogenesis, rooting of cuttings, and organogenesis of conifers. *Can. J. For. Res.* **2015**, *45*, 379–383. <https://doi.org/10.1139/cjfr-2014-0360>.
- Bonga, J.M. Can explant choice help resolve recalcitrance problems in *in vitro* propagation, a problem still acute especially for adult conifers? *Trees Struct. Funct.* **2017**, *31*, 781–789. <https://doi.org/10.1007/s00468-016-1509-z>.

- Breidenbach, N.; Gailing, O.; Krutovsky, K.V. Genetic structure of coast redwood (*Sequoia sempervirens* [D. Don] Endl.) populations in and outside of the natural distribution range based on nuclear and chloroplast microsatellite markers. *PLoS ONE* **2020**, *15*, e0243556. <https://doi.org/10.1371/journal.pone.0243556>.
- Bueno, N.; Cuesta, C.; Centeno, M.L.; Ordás, R.J.; Alvarez, J.M. In vitro plant regeneration in conifers: The role of WOX and KNOX gene families. *Genes* **2021**, *12*, 438. <https://doi.org/10.3390/genes12030438>.
- Burns, E.E., Campbell, R., Cowan, P.D. 2018. State the Redwoods Conservation. Report of Save the Redwoods. Available at: <https://www.savetheredwoods.org>.
- Bussmann, R.W.; Paniagua-Zambrana, N.Y.; Njoroge, G.N. *Pinus patula* Schlttdl. & Cham. *Pinus radiata* D. Don. PINACEAE. In *Ethnobotany of the Mountain Regions of Africa. Ethnobotany of Mountain Regions*; Bussmann, R.W., Ed.; Springer: Cham. 2020; pp.1-9.https://doi.org/10.1007/978-3-319-77086-4_124-1.
- Calamar, A.; De Klerk, G.J. Effect of sucrose on adventitious root regeneration in apple. *Plant Cell Tissue Organ Cult.* **2002**, *70*, 207–212. <https://doi.org/10.1023/A:1016356123367>.
- Cantero, A. 2019. Proyecto life healthy forest. *Informe final de HAZI Fundazioa*. Available at: <https://www.lifehealthyforest.com>.
- Castander-Olarieta, A.; Montalbán, I.A.; De Medeiros Oliveira, E.; Dell'Aversana, E.; D'Amelia, L.; Carillo, P.; Steiner, N.; Fraga, H.P.F.; Guerra, M.P.; Goicoa, T.; et al. Effect of thermal stress on tissue ultrastructure and metabolite profiles during initiation of radiata pine somatic embryogenesis. *Front. Plant Sci.* **2019**, *9*, 1-16. <https://doi.org/10.3389/fpls.2018.02004>.
- Cézar, T.M.; Higa, A.R.; Koehler, H.S.; Ribas, L.L.F. Influence of culture medium, explant length and genotype on micropropagation of *Pinus taeda* L. *Ciênc. Florest.* **2015**, *25*, 13–22. <https://doi.org/10.1590/1980-509820152505013>.
- Chang, S.H.; Ho, C.K.; Chen, Z.Z.; Tsay, J.Y. Micropropagation of *Taxus mairei* from mature trees. *Plant Cell Rep.* **2001**, *20*, 496–502. <https://doi.org/10.1007/s002990100362>.
- Charpin, D.; Sénéchal, H.; Poncet, P. Respiratory Allergy to Conifers. In *Conifers: Recent Advances*; Gonçalves, A. C., Fonseca, T. F., Eds.; Intechopen Limited: London, United Kingdom, 2022, pp. 203– 218. <http://dx.doi.org/10.5772/intechopen.95677>.
- Chen, L.; Zhang, K.; Gong, X.; Wang, H.; Gao, Y.; Wang, X.; Zeng, Z., Hu, Y. Effects of different LEDs light spectrum on the growth, leaf anatomy, and chloroplast ultrastructure of potato plantlets in vitro and minituber production after transplanting in the greenhouse. *J. Integr. Agric.* **2020**, *19*, 108–119. [https://doi.org/10.1016/S2095-3119\(19\)62633-X](https://doi.org/10.1016/S2095-3119(19)62633-X).
- Chen, Y.C.; Chang, C.; Lin, H.L. Topolins and red light improve the micropropagation efficiency of passion fruit (*Passiflora edulis* Sims) "Tainung No. 1". *HortScience* **2020**, *55*, 1337–1344. <https://doi.org/10.21273/HORTSCI15078-20>.
- Clapa, D.; Fira, A.; Hărșan, E.; Chiș, L.; Roman, G.; Cantor, M. *Sequoia sempervirens* in Transylvania. *Acta Hort.* **2010**, *885*, 85–90. <https://doi.org/10.17660/ActaHortic.2010.885.11>.
- Coleman, G.D.; Ernst, S.G. In vitro shoot regeneration of *Populus deltoides*: Effect of cytokinin and genotype. *Plant Cell Rep.* **1989**, *8*, 459–462. <https://doi.org/10.1007/BF00269048>.

- Corrêa, L.D.R.; Paim, D.C.; Schwambach, J.; Fett-Neto, A.G. Carbohydrates as regulatory factors on the rooting of *Eucalyptus saligna* Smith and *Eucalyptus globulus* Labill. *Plant Growth Regul.* **2005**, *45*, 63–73. <https://doi.org/10.1007/s10725-004-6125-z>.
- Cortizo, M.; De Diego, N.; Moncaleán, P.; Ordás, R.J. Micropropagation of adult stone pine (*Pinus pinea* L.). *Trees Struct. Funct.* **2009**, *23*, 835–842. <https://doi.org/10.1007/s00468-009-0325-0>.
- Da Silva, J. A. T.; Winarto, B.; Dobránszki, J.; Cardoso, J. C.; Zeng, S. Tissue disinfection for preparation of *Dendrobium* in vitro culture. *Folia Horticolt.* **2016**, *28*, 57–75. <https://doi.org/10.1515/fhort-2016-0008>.
- De Diego, N.; Montalbán, I.A.; Fernandez de Larrinoa, E.; Moncaleán, P. In vitro regeneration of *Pinus pinaster* adult trees. *Can. J. For. Res.* **2008**, *38*, 2607–2615. <https://doi.org/10.1139/X08-102>.
- De Diego, N.; Montalbán, I.A.; Moncaleán, P. In vitro regeneration of adult *Pinus sylvestris* L. trees. *S. Afr. J. Bot.* **2010**, *76*, 158–162. <https://doi.org/10.1016/j.sajb.2009.09.007>.
- De Diego, N.; Montalbán, I.A.; Moncaleán, P. Improved micropropagation protocol for maritime pine using zygotic embryos. *Scand. J. For. Res.* **2011**, *26*, 202–211. <https://doi.org/10.1080/02827581.2011.559174>.
- De Oliveira, L.F.; Ribas, L.L.F.; Quoirin, M.; Koehler, H.S.; Amano, E.; Higa, A.R. Micropropagation of *Pinus taeda* L. from juvenile material. *Tree For. Sci. Biotechnol.* **2012**, *6*, 96–101.
- Desai, C.; Inghalihalli, R.; Krishnamurthy, R. Micropropagation of *Anthurium andraeanum* an important tool in floriculture. *J Pharmacogn Phytochem.* **2015**, *4*, 112–117.
- Dobránszki, J.; Teixeira da Silva, J.A. Micropropagation of Apple—A Review. *Biotechnol. Adv.* **2010**, *28*, 462–488. <https://doi.org/10.1016/j.biotechadv.2010.02.008>.
- Dos Santos Junior, C.F.; Rech, T.D.; Costa, M.D.; Navroski, M.C.; Boff, P.; Boff, M.I.C. Efeito da qualidade de luz na multiplicação e enraizamento in vitro de *Cedrela odorata* L. *Ciênc. Florest.* **2022**, *32*, 2419–2432. <https://doi.org/10.5902/1980509866513>.
- Ellis, D.; Bilderback, D.E. Ponderosa pine (*Pinus ponderosa* P. Lawson & C. Lawson). In *Biotechnology in Agriculture and Forestry 16, Trees III*; Bajaj, Y., Ed.; Springer: Berlin, Germany, 1991; pp. 339–357. https://doi.org/10.1007/978-3-662-13231-9_21.
- Elvira-Recuenco, M.; Cacciola, S.O.; Sanz-Ros, A.V.; Garbelotto, M.; Aguayo, J.; Solla, A.; Mullett, M.; Drenkhan, T.; Oskay, F.; Aday Kaya, A.G.; Iturritxa, E.; Cleary, M.; Witzell, J.; Georgieva, M.; Papazova-Anakieva, I.; Chira, D.; Paraschiv, M.; Musolin, D.L.; Selikhovkin, A.V.; Varentsova, E.Y.; Adamčíková, K.; Markovskaja, S.; Mesanza, N.; Davydenko, K.; Capretti, P.; Scanu, B.; Gonthier, P.; Tsopeles, P.; Martín-García, J.; Morales-Rodríguez, C.; Lehtijärvi, A.; Doğmuş Lehtijärvi, H.T.; Oszako, T.; Nowakowska, J.A.; Bragança, H.; Fernández-Fernández, M.; Hantula, J.; Díez, J.J. Potential Interactions between Invasive *Fusarium circinatum* and Other Pine Pathogens in Europe. *Forests* **2020**, *11*, 7. <https://doi.org/10.3390/F11010007>.
- Escandón, M.; Valledor, L.; Pascual, J.; Pinto, G.; Cañal, M.J.; Meijón, M. System-wide analysis of short-term response to high temperature in *Pinus radiata*. *J. Exp. Bot.* **2017**, *68*, 3629–3641. <https://doi.org/10.1093/jxb/erx198>.
- Espinoza, S.E.; Quiroz, I.A.; Magni, C.R.; Yáñez, M.A. Variation in early growth and survival of *Pinus ponderosa* provenances established in Southern Chile. *New For.* **2021**, *52*, 959–970. <https://doi.org/10.1007/s11056-021-09833-6>.

Fang, W.; Jao, R.C. A review on artificial lighting of tissue cultures and transplants. In *Transplant Production in the 21st Century*; Kubota, C., Chun, C., Eds.; Springer: Dordrecht, Netherlands, 2000; pp. 108–113. https://doi.org/10.1007/978-94-015-9371-7_17.

Farjon, A. *Cryptomeria japonica*. *Curtis's Bot.* **1999**, *16*, 212-228.

Farjon, A. The Kew review: Conifers of the world. *Kew Bull.* **2018**, *73*, 8. <https://doi.org/10.1007/s12225-018-9738-5>.

Fitzgerald, S. Fire ecology of Ponderosa pine and the rebuilding of fire-resilient ponderosa pine ecosystems. In *Proceedings of the Symposium on Ponderosa Pine: Issues, Trends, and Management, Klamath Falls, Oregon, USA, 18–21 October 2005*; Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture: California, USA, 2005; pp. 197–225. <https://doi.org/10.2737/PSW-GTR-198>.

George, E.F. Plant Tissue Culture Procedure - Background. In *Plant Propagation by Tissue Culture*; George, E.F., Hall, M.A., De Klerk, G.J., Eds.; Springer: Dordrecht, The Netherlands, 2008; pp. 1–28. https://doi.org/doi.org/10.1007/978-1-4020-5005-3_1.

George, E.F.; Davies, W. Effects of the Physical Environment. In *Plant Propagation by Tissue Culture*; George, E.F., Hall, M.A., De Klerk, G.J., Eds.; Springer: Dordrecht, The Netherlands, 2008; pp. 423–464. https://doi.org/doi.org/10.1007/978-1-4020-5005-3_1.

George, E.F.; Debergh, P.C. Micropropagation: Uses and Methods. In *Plant Propagation by Tissue Culture*; George, E.F., Hall, M.A., De Klerk, G.J., Eds.; Springer: Dordrecht, The Netherlands, 2008; pp. 29–64. https://doi.org/doi.org/10.1007/978-1-4020-5005-3_1.

Gibson, S.I. Plant sugar-response pathways. part of a complex regulatory web. *Plant Physiol.* **2000**, *124*, 1532–1539. <https://doi.org/10.1104/pp.124.4.1532>.

Giri, C.C.; Shyamkumar, B.; Anjaneyulu, C. Progress in tissue culture, genetic transformation and applications of biotechnology to trees: An overview. *Trees* **2004**, *18*, 115–135. <https://doi.org/10.1007/s00468-003-0287-6>.

Gnasekaran, P.; Rahman, Z.A.; Chew, B.L.; Appalasamy, S.; Mariappan, V.; Subramaniam, S. Development of micropropagation system of *Zingiber officinale* var. *rubrum* Theilade using different spectrum light-emitting diode (LED) irradiation. *Ind. Crops Prod.* **2021**, *170*, 113748. <https://doi.org/10.1016/j.indcrop.2021.113748>.

Gupta, S.; Jatothu, B. Fundamentals and applications of light-emitting diodes (LEDs) in in vitro plant growth and morphogenesis. *Plant Biotechnol. Rep.* **2013**, *7*, 211–220. <https://doi.org/10.1007/s11816-013-0277-0>.

Gupta, S.; Sahoo, T.K. Light emitting diode (LED) induced alteration of oxidative events during in vitro shoot organogenesis of *Curculigo orchiooides* Gaertn. *Acta Physiol. Plant.* **2015**, *37*, 233. <https://doi.org/10.1007/s11738-015-1990-9>.

Haberlandt, G. Kulturversuche mit isolierten Pflanzenzellen. *Sitzungsber, Akad Wiss Wien Math-Naturwiss Kl. Abt J.* **1902**, *11*, 69–92.

Hall, F.C. Fire history-blue mountains, Oregon. In *Proceedings of the Fire History Workshop, Tucson, Arizona, USA, 20–24 October 1980*; Stokes, M.A., Dieterich, J.H., Eds.; Rocky Mountain Forest and Range Experiment Station, Forest Service, U.S. Department of Agriculture: Colorado, USA, 1980; pp. 75–81. <https://doi.org/10.2737/RM-GTR-81>.

HAZI. El bosque vasco en cifras 2022. *Informe de HAZI Fundazioa sobre el Inventario forestal del País Vasco-2022*. Available at: <http://www.nasdap.net/inventarioforestal>.

Ibañez, M.; Caru, M.; Herrera, M. A.; Gonzalez, L.; Martin, L. M.; Miranda, J.; Navarro-Cerrillo, R. M. Clones identification of *Sequoia sempervirens* (D. Don) Endl. in Chile by using PCR-RAPDs technique. *J. Zhejiang Univ. Sci. B.* **2009**, *10*, 112-119.

Ibarrondo, M. J. A. El monocultivo de pino radiata en el País Vasco: Origen y claves de permanencia de un sistema de explotación contrario al desarrollo sostenible. *Estud. Geogr.* **2008**, *69*, 335–356. <https://doi.org/10.3989/estgeogr.0426>.

Jankovský, L.; Palovčíková, D.; Dvořák, M.; Tomšovský, M. Records of Brown spot needle blight related to *Lecanosticta acicola* in the Czech Republic. *Plant Prot. Sci.* **2009**, *45*, 16–18. <https://doi.org/10.17221/39/2008-pps>.

Koguta, K.V.; Flores, P.C.; Alcantara, G.B.; Higa, A.R. Serial micropropagation and cuttings as rejuvenation methods for *Cryptomeria japonica* (L.F) D.Don. *Rev. Espac.* **2017**, *38*, 7.

Kulus, D.; Woźny, A. Influence of light conditions on the morphogenetic and biochemical response of selected ornamental plant species under in vitro conditions: A mini-review. *Bio Technol.* **2020**, *101*, 75–83. <https://doi.org/10.5114/bta.2020.92930>.

Kumar, N.; Reddy, M.P. In vitro plant propagation: A review. *J. For. Environ. Sci.* **2011**, *27*, 61–72. <https://doi.org/10.7747/JFS.2011.27.2.1>.

Lai, C.S.; Kho, Y.H.; Chew, B.L.; Raja, P.B.; Subramaniam, S. Organogenesis of *Cucumis metuliferus* plantlets under the effects of LEDs and silver nanoparticles. *S. Afr. J. Bot.* **2022**, *148*, 78–87. <https://doi.org/10.1016/j.sajb.2022.04.020>.

Lambardi, M.; Sharma, K.K.; Thorpe, T.A. Optimization of in vitro bud induction and plantlet formation from mature embryos of Aleppo Pine (*Pinus halepensis* Mill). *In Vitro Cell. Dev. Biol. Plant.* **1993**, *29*, 189–199. <https://doi.org/10.1007/BF02632034>.

Larraburu, E.E.; Correa, G.S.; Llorente, B.E. In vitro development of yellow lapacho (Bignoniaceae) using high-power light emitting diode. *Rev. Arvore.* **2018**, *42*, e420508. <https://doi.org/10.1590/1806-90882018000500008>.

Laslo, V.; Agud, E.; Vicaș, S. In Vitro Modeling of Morfogenesis Processes in *Sequoia Sempervirens* (D. Don) Endl, Via the Use of Phytohormones. *Natural Resources Sustainable Development*, **2015**, *5*, 73-78. <https://doi.org/10.31924/nrsd>.

Liang, Y.; Bai, X.; Xu, X.; Xu, H.G.; Wang, J.; Pan, P. Direct in vitro organogenesis from sprouted seeds of a highly economical and ecological valued tree, Korean pine. *Plant Cell Tissue Organ Cult.* **2022**, *148*, 197–207. <https://doi.org/10.1007/s11240-021-02164-6>.

Loberant, B.; Altman, A. Micropropagation of Plants. In *Encyclopedia of Industrial Biotechnology: Bioprocess, Bioseparation, and Cell Technology*; Wiley: New York, NY, USA, 2010; pp. 1–17.

Loyola-Vargas, V.M.; Ochoa-Alejo, N. An introduction to plant tissue culture: Advances and perspectives. In *Plant Cell Culture Protocols*; Loyola-Vargas, V., Ochoa-Alejo, N., Eds; Humana Press: New York, NY, USA, 2012; pp. 1–8. https://doi.org/10.1007/978-1-4939-8594-4_1

Ma, Q.W.; Li, F.L.; Li, C.S. The coast redwoods (*Sequoia*, Taxodiaceae) from the Eocene of Heilongjiang and the Miocene of Yunnan, China. *Rev. Palaeobot. Palynol.* **2005**, *135*, 117e129. 2005. <https://doi.org/10.1016/j.revpalbo.2005.03.002>.

Martínez-Álvarez, P.; Pando, V.; Diez, J.J. Alternative species to replace Monterey pine plantations affected by pitch canker caused by *Fusarium circinatum* in northern Spain. *Plant Pathol.* **2014**, *63*, 1086–1094. <https://doi.org/10.1111/ppa.12187>.

Mead, D. J. Sustainable management of *Pinus radiata* plantations. FAO Forestry. Paper No. 170. Rome, FAO, 2013. Available at: <https://www.fao.org>.

Meneguzzi, A.; Konzen, E. R.; Navroski, M. C.; Camargo, S. S.; Pereira, M. D. O.; Rufato, L. Shoot multiplication of two *Sequoia sempervirens* genotypes with addition of small concentrations of kinetin. *Pesqui. Florest. Bras.* **2019**, *39*, 1-8. <http://dx.doi.org/10.4336/2019.pfb.39e201701550>.

Meneguzzi, A.; Camargo, S. S.; Navroski, M. C.; Pereira, M. D.O.; Souza, P.; Rufato, L. Agar and activated charcoal influence the in vitro development of shoot and root *Sequoia*. *Rev. Ciências Agroveterinarias.* **2020** *18*, 20–24 <https://doi.org/10.5965/2238117118e2019020>.

Mesanza, N.; García-García, D.; Raposo, E.R.; Raposo, R.; Iturrutxa, M.; Pascual, M.T.; Barrena, I.; Urkola, A.; Berano, N.; Sáez de Zerain, A.; Iturrutxa, E. Weather Variables Associated with Spore Dispersal of *Lecanosticta acicola* Causing Pine Needle Blight in Northern Spain. *Plants* **2021a**, *10*, 2788. <https://doi.org/10.3390/plants10122788>.

Mesanza, N.; Hernandez, M.; Raposo, R.; Iturrutxa, E. First Report of *Mycosphaerella dearnessii*, Teleomorph of *Lecanosticta acicola*, in Europe. *Plant Health Prog.* **2021b**, *22*, 565–566. <https://doi:10.1094/PHP-03-21-0060-BR>.

Mesanza, N.; Raposo, R.; Elvira-Recuenco, M.; Barnes, I.; van der Nest, A.; Hernández, M.; Pascual, M.T.; Barrena, I.; San Martín, U.; Cantero, A.; Hernández-Escribano, L.; Iturrutxa, E. New hosts for *Lecanosticta acicola* and *Dothistroma septosporum* in newly established arboreta in Spain. *For. Pathol.* **2021c**, *51*. <https://doi:10.1111/efp.12650>.

Miller, M. Fire Autecology. In *Wildland Fire in Ecosystems: Effects of Fire on Flora*; Tech. Rep. RMRS-GTR-42; Brown, J., Smith, J., Eds.; Rocky Mountain Research Station, Forest Service, U.S. Department of Agriculture: Ogden, Utah, USA, 2000; volume 2, pp 257. <https://doi.org/10.2737/RMRS-GTR-42-V2>.

Montalbán, I.A.; De Diego, N.; Moncaleán, P. Testing novel cytokinins for improved in vitro adventitious shoots formation and subsequent ex vitro performance in *Pinus radiata*. *Forestry* **2011**, *84*, 363–373. <https://doi.org/10.1093/forestry/cpr022>.

Montalbán, I.A.; Novák, O.; Rolčík, J.; Strnad, M.; Moncaleán, P. Endogenous cytokinin and auxin profiles during in vitro organogenesis from vegetative buds of *Pinus radiata* adult trees. *Physiol. Plant.* **2013**, *148*, 214–231. <https://doi.org/10.1111/j.1399-3054.2012.01709.x>.

Montalbán, I.A.; García-Mendiguren, O.; Moncaleán, P. Somatic Embryogenesis in *Pinus* spp. In *In Vitro Embryogenesis in Higher Plants. Methods in Molecular Biology*; Germana, M., Lambardi, M.; Humana Press: New York, USA, 2016; vol 1359; pp.405-415. https://doi.org/10.1007/978-1-4939-3061-6_21.

Neale, D.B.; Wheeler, N.C. The Conifers. In *The Conifers: Genomes, Variation and Evolution*; Springer Nature: Cham, Switzerland, 2019; pp. 1–21. https://doi.org/10.1007/978-3-319-46807-5_1.

Neelakandan, A.K.; Wang, K. Recent progress in the understanding of tissue culture-induced genome level changes in plants and potential applications. *Plant Cell Rep.* **2012**, *31*, 597–620. <https://doi.org/10.1007/s00299-011-1202-z>.

Onuma, Y.; Uchiyama, K.; Kimura, M.; Tsumura, Y. Genetic analysis distinguished new natural population and old plantations of *Cryptomeria japonica*. *Trees For. People* **2023**, *11*, 100365. <https://doi.org/10.1016/j.tfp.2022.100365>.

Page, C.N. Taxodiaceae. In *The families and genera of vascular plants. V. 1: Pteridophytes and gymnosperms*; Kramer, K. U., Green, P. S., Kubitzki, K., Eds.; Springer-Verlag: Berlin, Germany, 1990; pp. 353–362.

Pawłowska, B.; Żupnik, M.; Szewczyk-Taranek, B.; Cioć, M. Impact of LED light sources on morphogenesis and levels of photosynthetic pigments in *Gerbera jamesonii* grown in vitro. *Hortic. Environ. Biotechnol.* **2018**, *59*, 115–123. <https://doi.org/10.1007/s13580-018-0012-4>.

Pereira, C.; Montalbán, I.A.; Pedrosa, A.; Tavares, J.; Pestryakov, A.; Bogdanchikova, N.; Canhoto, J.; Moncaleán, P. Regeneration of *Pinus halepensis* (Mill.) through organogenesis from apical shoot buds. *Forests* **2021**, *12*, 363. <https://doi.org/10.3390/f12030363>.

Phillips, G.C. In vitro morphogenesis in plants—Recent advances. *In Vitro Cell. Dev. Biol. Plant.* **2004**, *40*, 342–345. <https://doi.org/10.1079/IVP2004555>.

Phillips, G.C.; Garda, M. Plant tissue culture media and practices: An overview. *In Vitro Cell. Dev. Biol. Plant.* **2019**, *55*, 242–257. <https://doi.org/10.1007/s11627-019-09983-5>.

Puerta, F. R.; Cartagena, M. B.; Amiano, A. C. Aplicación del programa CUBICA a distintos regímenes de gestión del pino radiata en el País Vasco. *Cuad. Soc. Esp. Cien. For.* **2004**, *18*, 193–198.

Ragonezi, C.; Klimaszewska, K.; Castro, M.R.; Lima, M.; de Oliveira, P.; Zavattieri, M.A. Adventitious rooting of conifers: Influence of physical and chemical factors. *Trees Struct. Funct.* **2010**, *24*, 975–992. <https://doi.org/10.1007/s00468-010-0488-8>.

Raish, C.; Yong, W.; Marzluff, J. Contemporary human use of south western ponderosa pine forests. In *Songbird Ecology in South Western Ponderosa Pine Forests: A Literature Review*; Block, W.M., Finch, D.M., Eds.; Rocky Mountain Research Station, Forest Service, U.S. Department of Agriculture: Colorado, USA, 1997; pp. 28–42. <https://doi.org/10.2737/RM-GTR-292>.

Rapley, S. Redwood in New Zealand. *N. Z. J. For.* **2018**, *63*, 29–33.

Renau-Morata, B.; Ollero, J.; Arrillaga, I.; Segura, J. Factors influencing axillary shoot proliferation and adventitious budding in cedar. *Tree Physiol.* **2005**, *25*, 477–486. <https://doi.org/10.1093/treephys/25.4.477>.

Saborio, F.; Dvorak, W.S.; Donahue, J.K.; Thorpe, T.A. In vitro regeneration of plantlets from mature embryos of *Pinus ayacahuite*. *Tree Physiol.* **1997**, *17*, 787–796. <https://doi.org/10.1093/treephys/17.12.787>.

Sahu, J.; Sahu, R.K. A review on low cost methods for in vitro micropropagation of plant through tissue culture technique. *UK J. Pharm. Biosci.* **2013**, *1*, 38–41. <https://doi.org/10.20510/ukjpb/1/i1/91115>.

Sathyagowri, S.; Seran, T.H. In vitro plant regeneration of ginger (*Zingiber officinale* Rosc.) with emphasis on initial culture establishment. *J. Med. Aromat. Plants* **2011**, *1*, 195–202.

Shimizu, J. Y.; Maiochi, R. A. Criptoméria como espécie alternativa para produção de madeira no Paraná. 2007.

- Sillett, S.C.; Van Pelt, R.; Carroll, A.L.; Campbell-Spickler, J.; Antoine, M.E. Aboveground biomass dynamics and growth efficiency of *Sequoia sempervirens* forests. *For. Ecol. Manag.* **2020**, *458*, 117740. <https://doi.org/10.1016/j.foreco.2019.117740>.
- Sillett, S.C.; Antoine, M.E.; Carroll, A.L.; Graham, M.E.; Chin, A.R.O.; Van Pelt, R. Rangewide climatic sensitivities and non-timber values of tall *Sequoia sempervirens* forests. *For. Ecol. Manag.* **2022**, *526*, 120573. <https://doi.org/10.1016/j.foreco.2022.120573>.
- Solans, M.; Bernardi, G.C.; Raffaele, E.; Chaia, E. *Ochetophila*-infective *Frankia* colonization patterns of volcanic ash in Patagonia. *Symbiosis* **2021**, *85*, 359–369. <https://doi.org/10.1007/s13199-021-00820-1>.
- Stoiculescu, R.; Cogălniceanu, G.; Brezeanu, A.; Hristea, G. Influence of microstructured carbon materials on *Sequoia sempervirens* (D. Don.) Endl. in vitro culture. *Ann. For. Res.* **2009**, *52*, 137–142. <https://doi.org/10.15287/afr.2009.130>.
- Sul, I.W.; Korban, S.S. Effects of salt formulations, carbon sources, cytokinins, and auxin on shoot organogenesis from cotyledons of *Pinus pinea* L. *Plant Growth Regul.* **2004**, *43*, 197–205. <https://doi.org/10.1023/B:GROW.0000046013.47892.4f>.
- Sumaryono, W.M.; Ratnadewi, D. Effect of carbohydrate source on growth and performance of in vitro sago palm (*Metroxylon sago* Rottb.) plantlets. *HAYATI J. Biosci.* **2012**, *19*, 88–92. <https://doi.org/10.4308/hjb.19.2.88>.
- Taniguchi, T.; Konagaya, K.; Nanasato, Y. Somatic embryogenesis in artificially pollinated seed families of 2nd generation plus trees and cryopreservation of embryogenic tissue in *Cryptomeria japonica* D. Don (sugi). *Plant Biotechnol.* **2020**, *37*, 239–245. <https://doi.org/10.5511/plantbiotechnology.20.0220a>.
- Tsubomura, M.; Taniguchi, T. Micropropagation of the male sterile ‘Soushun’ Japanese cedar. *Comb. Proc. Int. Plant. Prop. Soc.* **2008**, *58*, 421–427.
- Tsumura, Y. Genetic structure and local adaptation in natural forests of *Cryptomeria japonica*. *Ecol. Res.* **2023**, *38*, 64–73. <https://doi.org/10.1111/1440-1703.12320>.
- Valverde-Cerdas, L.; Rojas-Vargas, A.; Hine-Gómez, A. In vitro propagation of *Albizia guachapele*, *Cedrela odorata*, *Platymiscium pinnatum* and *Guaicum sanctum*. *Plant Tissue Cult. Biotechnol.* **2008**, *18*, 151–156. <https://doi.org/10.3329/ptcb.v18i2.3397>.
- van Staden, J.; Zazimalova, E.; George, E.F. Plant Growth Regulators II: Cytokinins, their Analogues and Antagonists. In *Plant Propagation by Tissue Culture*; George, E.F., Hall, M.A., De Klerk, G.J., Eds.; Springer: Dordrecht, The Netherlands, 2008; pp. 205–226. https://doi.org/doi.org/10.1007/978-1-4020-5005-3_1.
- Watt, M.S.; Kimberley, M.O.; Rapley, S.; Webster, R. Comparing volume productivity of redwood and radiata pine plantations in New Zealand. *For. Ecol. Manag.* **2021**, *500*, 119628. <https://doi.org/10.1016/j.foreco.2021.119628>.
- Weng, Y.H.; Park, Y.S.; Krasowski, M.J.; Tosh, K.J.; Adams, G. Partitioning of genetic variance and selection efficiency for alternative vegetative deployment strategies for white spruce in eastern Canada. *Tree Genet. Genomes.* **2008**, *4*, 809–819. <https://doi.org/10.1007/s11295-008-0154-0>.
- Werbrouck, S.P.O.; Strnad, M.; Van Onckelen, H.A.; Debergh, P.C. Meta-topolin, an alternative to benzyladenine in tissue culture?. *Physiol. Plant.* **1996**, *98*, 291–297. <https://doi.org/10.1034/j.1399-3054.1996.980210.x>.

Chapter 1

Williams, C.G.; Savolainen, O. Inbreeding depression in conifers: implications for breeding strategy. *For. Sci.* **1996**, *42*, 102-117. <https://doi.org/10.1093/forestscience/42.1.102>.

Woods, A.J.; Martín-García, J.; Bulman, L.; Vasconcelos, M.W.; Boberg, J.; La Porta, N.; Peredo, H.; Vergara, G.; Ahumada, R.; Brown, A.; et al. *Dothistroma* needle blight, weather and possible climatic triggers for the disease's recent emergence. *For. Pathol.* **2016**, *46*, 443–452. <https://doi.org/10.1111/efp.12248>.

Zhang, J.W.; Finley, K.A.; Knapp, E.E. Resilience of a ponderosa pine plantation to a backfiring operation during a mid-summer wildfire. *Int. J. Wildland Fire* **2019**, *28*. <https://doi.org/10.1071/WF19033>.

Zhang, J.; Finley, K. A.; Young, D. H.; Fiddler, G. O.; Looney, C. Growth response of Ponderosa Pine to intensive cultural treatments varies with site quality and plantation age. *For. Sci.* **2022**, *68*, 212-225. <https://doi.org/10.1093/forsci/xfab065>.

CHAPTER 2

General Hypothesis and Objectives

General Hypothesis

This work wants to confirm the following hypothesis:

- The modification of the physical and chemical environment in the micropropagation process can influence the regeneration efficiency of elite plantlets of forestry species such as *Cryptomeria japonica*, *Pinus ponderosa* and *Sequoia sempervirens* to be considered as alternative to *Pinus radiata* in the Basque Country.
- Adult *Pinus radiata* trees with apparent tolerance to needle blight can be used as a source of material for *in vitro* propagation.

Objectives

To validate these hypotheses, this work has been divided in four different objectives:

1. To evaluate the effect of genotypes, collection dates and BA concentrations on the shoot induction stage; and investigate the influence of different light types on the success of root induction of *Pinus radiata* shoots coming from adult trees. To determine if the assayed environmental conditions modified the content of amino acids and carbohydrates during the rooting phase.
2. To study the effect of different types of explants, cytokinins and culture media in the micropropagation of *Cryptomeria japonica* adult trees. To evaluate the influence of different sucrose concentrations, ventilated culture containers and light treatments in the rooting stage.
3. To analyze the influence of different types of explants, cytokinins, culture media, and light treatments in the improvement of organogenesis in *Pinus ponderosa*. To determine through anatomical analyses if the resulting *in vitro* needles present anatomical differences after different light treatments.
4. To determine the influence of different explants and cytokinin types in the micropropagation process of *Sequoia sempervirens* elite adult trees. To study the effect of different auxins on the *in vitro* rooting and subsequent *ex vitro* acclimatization stage

CHAPTER 3

**Influence of Physico-Chemical Factors on the Efficiency and
Metabolite Profile of Adult *Pinus radiata* D. Don Bud
Organogenesis**

The content of this chapter corresponds to the published article "Rojas-Vargas, A.; Castander-Olarieta, A.; Montalbán, I.A.; Moncaleán, P. 2022. Influence of Physico-Chemical Factors on the Efficiency and Metabolite Profile of Adult *Pinus radiata* D. Don Bud Organogenesis. *Forests* 13, 1455. <https://doi.org/10.3390/f13091455>".

ABSTRACT

Genetic improvement programs for conifer species face the challenge of propagating elite individuals with superior characteristics in the present landscape of climate change; the problem is focused on the fact that when these individuals have shown the desirable traits, they have changed phase and therefore have lost the ability to be propagated by traditional methods. Based on our previous works on *Pinus* spp., regeneration of adult trees through organogenesis and trying to improve the protocol in *Pinus radiata*, our objective was to analyze the influence of collection dates and different BA concentrations in the first phase of SI, as well as the effect of different light types on the success of RI. Moreover, we were interested in studying the effect of the abovementioned physico-chemical factors on the amino acid and carbohydrate content in *in vitro* shoots. Reinvigorated shoots were obtained in both BA concentrations (22 or 44 μ M), although the highest BA concentration showed the best results in terms of SI (explants forming shoots (EFS) (46%) and NS/E (1.95 \pm 0.52)) when using initial explants collected in the first week of February. The percentage of EFS was genotype dependent. Highest EFS percentage (91%) was obtained when the explants from the better responding genotype were induced with the highest BA concentration. With respect to the light treatment applied, significant differences in RI (20%) and in the number of roots per explant (NR/E) (4.62 \pm 0.65) were observed in shoots cultured under white FL. Finally, significant differences in different phases of the rooting process were detected in the amounts of fructose, glucose and sucrose and in the content of threonine and tyrosine.

RESUMEN

Los programas de mejoramiento genético para especies forestales de coníferas enfrentan el desafío de propagar individuos élite con características superiores en el panorama actual de cambio climático; el problema se centra en que cuando estos individuos han mostrado los rasgos deseables, han cambiado de fase y por lo tanto han perdido la capacidad de ser propagados por métodos tradicionales. En base a trabajos previos sobre regeneración *in vitro* de árboles adultos de *Pinus* spp. mediante organogénesis, y tratando de mejorar el protocolo en *Pinus radiata*, nuestro objetivo fue analizar la influencia de las fechas de recolección de los explantos y diferentes concentraciones de BA en la fase de inducción de brotes, así como el efecto de diferentes tipos de luz en el éxito del enraizamiento. Además, nos interesó estudiar el efecto de los factores físico-químicos antes mencionados sobre el contenido de aminoácidos y carbohidratos en los brotes desarrollados *in vitro*. Se obtuvieron brotes revigorizados en ambas concentraciones de BA (22 o 44 μ M); sin embargo, la mayor concentración de BA mostró los mejores resultados en términos de inducción de brotes (46%) y número de brotes por explanto (1,95 \pm 0,52) al utilizar material recolectado en la primera semana de febrero. El porcentaje de explantos que forman brotes fue dependiente del genotipo. El mayor porcentaje de explantos que forman tallos se obtuvo cuando los explantos del genotipo de mayor respuesta se cultivaron en la concentración más alta de BA (91%). Con respecto al tratamiento de luz aplicado, se observaron diferencias significativas en la inducción de raíces (20%) y en el número de raíces por explanto (4,62 \pm 0,65) en los brotes cultivados bajo FL. Finalmente, se detectaron diferencias significativas en las distintas fases del proceso de enraizamiento en el perfil de fructosa, glucosa y sacarosa y en el contenido de treonina y tirosina.

1. INTRODUCTION

Conifers are an important gymnosperm group, distributed worldwide, and they form extensive, circumboreal forests across North America and Eurasia (Neale and Wheeler, 2019). Conifers are particularly important since they supply over 50% of the world's timber harvest and present technical and economic advantages for wood-based industries. In forestry plantations, conifers are preferred over angiosperms as they produce a much faster economic yield with more predictable shapes and sizes of timber (Farjon, 2018).

Among all conifers, the family Pinaceae far outweighs others in economic importance and the genera *Pinus* ranks first. Radiata pine (*Pinus radiata* D. Don), native to the Central Coast of California and Mexico, is the most planted pine species, and its plantations have been extended throughout Australia, Chile, New Zealand, South Africa and Northern Spain (Escandón et al., 2017). However, in recent years, many countries have reported problems regarding the commercial viability of plantations due to effects provoked by fungus and other stresses derived from climate change (Allen et al., 2010; Castander-Olarieta et al., 2019). Brown spot needle blight caused by *Lecanosticta acicola* and Dothistroma needle blight caused by the fungi *Dothistroma septosporum* and *D. pini* are forest diseases that affect this species (Jankovský et al., 2009; Woods et al., 2016). For this reason, great efforts are being carried out to develop efficient methods to propagate selected trees with tolerance to several biotic and abiotic stresses. In this sense, *in vitro* methods provide tools that can be used for the clonal propagation of *Pinus* spp., as well as for the creation of backup collections as an alternative to the *in situ* conservation of species (De Diego et al., 2008; Cortizo et al., 2009; De Diego et al., 2010; Reed et al., 2011).

In *in vitro* culture, the physico-chemical environment is very important and sometimes determines the success of the morphogenic process. The type and concentration of PGRs, vitamins, basal media, solidifying agents, pH, etc., are crucial to obtain a highly efficient *in vitro* regeneration procedure (Phillips, 2004; Kulus and Woźny, 2020). In this sense, CKs added to the culture medium directly affect organogenesis, having a direct response to the induction of axillary shoot buds due to their effect on the endogenous phytohormone balance (Phillips, 2004; de Almeida et al., 2012).

Light also affects plant morphogenesis, and it may not only induce plant development but also induce photo-inhibition when leaves are exposed to higher intensity of ore light than they can use (Thorpe et al., 2008). Usually, FL are used for plant micropropagation in growth chambers, with irradiances between 25 and 150 $\mu\text{molm}^{-2}\text{s}^{-1}$ for a 16 h photoperiod (Larraburu et al., 2018). However, this illumination method has some drawbacks, such as its short lifespan (10,000 h) and the fact that it produces heat, which leads to the need for an extensive cooling system and high maintenance costs (Jao and Fang, 2003; Thorpe et al., 2008). LEDs are being used currently as an alternative light source for controlled environment horticulture technology (Riikonen et al., 2016). The high efficiency in the energy conversion of LEDs reduces the heat emissions and thereby saves energy. LEDs are also environmentally safer than FL tubes and are made of recyclable material (Astolfi et al., 2012).

When optimizing a tissue culture protocol, the source of explant, the explant type and its developmental stage are also key factors (Kumar and Reddy, 2011). In the specific case of cells from adult explants, they are less prone to initiating dedifferentiation and reprogramming processes than juvenile explant cells (Narváez et al., 2019). Moreover, the effect of the initial explant collection date on the morphogenic and embryogenic success has been demonstrated (Montalbán et al., 2011a).

The development of a successful tissue culture protocol to achieve *in vitro* shoots from *Pinus* adult trees requires not only the optimization of adventitious shoot initiation and elongation but also the improvement of the rooting process. There are many chemical factors involved in the success of the process; the carbohydrate source has been described as one of them, involved in the growth of adventitious roots in shoots (Ragonezi et al., 2010). Traditionally, sucrose is commonly used in tissue culture

because it is readily assimilated by plants, but, in some cases, high concentrations of this compound could have negative effects in the root elongation phase (Kumar and Reddy, 2011).

In the case of *Pinus*, Neiker-BRTA team established successful protocols for adult pine organogenesis in *Pinus pinaster*, *P. sylvestris*, *P. pinea*, *P. radiata* and *P. halepensis* (De Diego et al., 2008; Cortizo et al., 2009; De Diego et al., 2010; Montalbán et al., 2011b; Pereira et al., 2021). These kinds of procedures showed low success and were sometimes extremely dependent on the genotype. For this reason and trying to increase the yield of this *in vitro* biotechnological tool, our main goal was to study the effects of different physico-chemical factors on the organogenesis process of *P. radiata* buds from adult trees. Moreover, we focused our study on the involvement of the assayed environmental conditions in the profile of amino acids and carbohydrates during the rooting phase.

2. MATERIALS AND METHODS

2.1. Plant Material

Vegetative shoot buds (1-7 cm length) (Figure 1A) were collected every 15 days from 13 January to the last week of January 2020, and every 8 days from the first to the fourth week of February 2020, comprising a total of 6 collection dates. Eight 13-year-old adult trees (A, B, C, D, E, F, G and J), located in Neiker-BRTA, Arkaute (Spain; 42°51'08.5" N, 2°37'37.1" W), and one 9-year-old juvenile somatic tree (N) (Spain; 42°51'5.51" N, 2°37'19.43" W) were chosen to carry out the experiment. After cutting and collecting buds from the trees, they were wrapped in moist paper to prevent dehydration and stored in polyethylene bags at 4 °C for a maximum of 3 days. The vegetative buds were disinfected as follows: all explants were washed with commercial detergent, rinsed under running water for 10 min, immersed in ethanol 96% for 2 min and then washed three times with sterile distilled water. After this, explants were disinfected in 1 mL⁻¹ silver nanoparticle solution (Argovit®, Vector Vita LLC, Novosibirsk, Russia) (1.2%, w/w) of metallic silver stabilized with 18.8% (w/w) of polyvinylpyrrolidone (PVP) suspended in water (80% w/w) for 15 min and then rinsed three times with sterile distilled water (Stephano-Hornedo et al., 2020). Finally, the bud scales were removed, and the explants were cut transversely with a surgical scalpel blade into 0.5-1.0 cm-thick slices (Figure 1B) under sterile conditions in a laminar-flow unit. The slices were directly used to study the effect of BA on the induction medium (IM).




Figure 1. Organogenic process in apical shoot buds from *Pinus radiata* D. Don adult trees: **(A)** apical shoot buds, bar=1 cm; **(B)** shoot bud slices in culture, bar=1 cm; **(C)** shoot bud slice after induction with 22 μ M 6-benzyladenine (BA), bar=1 cm; **(D)** shoots in elongation medium, bar=1 cm; **(E)** rooted shoots, bar=1 cm; **(F)** acclimatized shoots, bar=3 cm.

2.2. Organogenic Process

Four to five slices were cultivated vertically on Petri dishes (90 × 15 mm) containing 20 mL of bud IM medium (Figure 1B). IM consisted of LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)], supplemented with 3% (w/v) sucrose and solidified with 8 gL⁻¹ Difco Agar®. The effect of two concentrations of BA was evaluated (22 and 44 µM). The pH of the medium was adjusted to 5.8 before autoclaving (121 °C, 20 min). Four to five Petri dishes per BA concentration and genotype were laid on the growth chamber at a temperature of 21±1 °C, under a 16 h photoperiod, with 120 µmolm⁻²s⁻¹ of light intensity provided by cool FL (TLD 58 W/33; Philips, France).

2.2.1. Shoot Growth and Elongation

As soon as bud induction was observed (between 7 and 9 weeks in IM) (Figure 1C), explants were transferred to Petri dishes with elongation medium (EM). EM consisted of hormone-free LP medium supplemented with 2 gL⁻¹ activated charcoal and 3% (w/v) sucrose and solidified with 8.5 gL⁻¹ Difco Agar®. The pH of the medium was adjusted to 5.8. After 35-50 days in culture (when elongating needle fascicles were evident), explants were subcultured into baby food jars (150 mL) containing 25 mL of EM (Figure 1D). The shoots were subcultivated every six weeks to the same medium. The light intensity, photoperiod and temperature of the growth chamber were the same as described above.

When shoots were 1.0-1.5 cm in length in the original bud slice, they were separated and cultivated individually in fresh EM (Figure 1D). The explants that showed new axillary shoot formation were cultivated into fresh EM, comprising a total of 6 subcultures, and the remaining explants showing secondary needles were separated and cultivated again with IM treatment to promote axillary bud development (re-induction).

2.2.2. Root Induction and Acclimatization of Rooted Plants

After the elongation phase, shoots of at least 2.0-2.5 cm length were used for RI. The explants were cultivated in Ecoboxes (Eco2box/green filter: a polypropylene vessel with a “breathing” hermetic cover, Duchefa®) containing 100 mL of RIM, which consisted of LP medium supplemented with a mixture of 5 µM NAA and 10 µM IBA, 8 gL⁻¹ Difco Agar® and (A) 3% (w/v) sucrose or (B) 1.5% (w/v) sucrose. Eight shoots per genotype and BA concentration were cultivated for each sucrose concentration. The pH of the culture media was adjusted to 5.8 before autoclaving. The shoots were placed in dim light at 21±1 °C for 8 days, followed by four weeks under a 16 h photoperiod. Two different light treatments were tested: (A) white light, 120 µmolm⁻²s⁻¹ of light intensity provided by cool white fluorescent tubes (TLD 58 W/33; Philips, France); and (B) red light, 67 µmolm⁻²s⁻¹ of light intensity provided by adjustable LEDs (RB4K Grow LEDs, GureLED-Meetthings, Vitoria-Gasteiz, Spain).

After four weeks of culture in RIM, explants were cultured in Ecoboxes containing 100 mL of root expression medium (REM), which consisted of LP medium supplemented with 2 gL⁻¹ activated charcoal, 3% (w/v) sucrose and 8.5 gL⁻¹ Difco Agar®, for six weeks. Then, explants with visible roots were transferred to wet peat moss (Pindstrup, Aarhus, Denmark): vermiculite (8:2, v/v) and acclimatized in the greenhouse under controlled conditions at 21±1 °C, decreasing the humidity progressively over one month from 95 to 80%.

2.3. Metabolite Extraction and Soluble Sugar and Amino acid Quantification

Metabolite extractions from explants collected after four weeks of culture in RIM treatments and after six weeks cultured in REM medium were performed following the protocol described by Valledor et al. (2014). Five to eight explants were taken from each condition (previous BA treatment, light type and sucrose content in the medium); no particular selection of the material was made. Each explant was considered as a biological replicate. Explants were ground in liquid nitrogen and 100 mg of the resulting fine powder was transferred to 2 mL microcentrifuge tubes containing 800 μ L cold metabolite extraction buffer (methanol:chloroform:water, 2.5:1:0.5, v:v:v). Then, tubes were centrifuged at 20,000 g during 6 min at 4 °C and the supernatants were transferred to new tubes containing 800 μ L of phase separation mix (chloroform:water, 1:1, v:v). After centrifugation at 10,000 g for 5 min at room temperature, the upper aqueous phases containing polar metabolites were saved in new 1.5 mL tubes. Aliquots of 200 μ L were then obtained from these tubes and completely dried on a Speedvac to remove the remaining methanol.

Finally, the samples were resuspended in 100 μ L ultra-pure water and soluble sugars and amino acids were quantified by High-Performance Liquid Chromatography (HPLC) (Agilent 1260 Infinity II, Agilent Technologies, Santa Clara, CA, United States). Soluble sugars, including sugar alcohol (fructose, galactose, glucose, mannitol and sucrose), were separated using a Hi-Plex Ca column (7.7 mm \times 300 mm, 8 μ m) and detected using a refractive index detector at a flow rate of 0.15 mLmin⁻¹ pure water at 80 °C for 30 min, as described by Castander-Olarieta et al. (2021). In the case of amino acids, a Poroshell 120 column (4.6 mm \times 100 mm, 2.7 μ m, Agilent Technologies, Santa Clara, CA, United States) was used, coupled to a fluorescence detector. The samples were injected into the column at a flow rate of 1.5 mLmin⁻¹ at 40 °C for 18 min with a discontinuous gradient. Solvent A was a mixture of 10 mM Na₂HPO₄ (Scharlau Chemie, Barcelona, Spain) and 10 mM Na₂B₄O₇ (pH 8.2) (Scharlau Chemie, Barcelona, Spain) and solvent B was acetonitrile: methanol:water (45:45:10, v:v:v) (Scharlau Chemie, Barcelona, Spain, HPLC-grade). The gradient program was the following: min 0-13.40, solvent A 98% and solvent B 2%; min 13.40-13.50, solvent A 43% and solvent B 57%; min 13.50-15.80, solvent B 100%, and min 15.80-18, solvent A 98% and solvent B 2%. Amino acids were estimated after pre-column derivatization by mixing 1 μ L sample with 2.5 μ L borate buffer (Agilent Technologies, Santa Clara, CA, United States), 32 μ L diluent (100 mL solvent A and 0.4 mL concentrated H₃PO₄) (PanReac AppliChem, Barcelona, Spain, pure pharma-grade) and 0.5 μ L o-phthaldialdehyde (OPA) (Agilent Technologies, Santa Clara, CA, United States). In the specific case of amino acids hydroxyproline and proline, samples were mixed with 0.5 μ L of 9-fluorenyl-methylchloroformate protecting group (FMOC) (Agilent Technologies, Santa Clara, CA, United States) instead of OPA. Detection was performed by analyzing fluorescence with excitation at 260 nm and emission at 450 nm for OPA derivatives and emission at 325 nm for FMOC derivatives.

In the case of amino acids, concentrations were determined from internal calibration curves constructed with the corresponding commercial standards (Agilent Technologies, Santa Clara, CA, United States). Carbohydrates such as fructose, galactose, mannitol and sucrose (Merck, Darmstadt, Germany) and glucose (Duchefa Biochemie, Haarlem, Netherlands) were determined from internal calibration curves constructed with the corresponding commercial standards. Results were conveniently adjusted considering the concentration step after methanol removal (2 times), and the results were expressed as μ molgFW⁻¹. In the case of certain amino acids, samples were re-injected after 1/32 dilution to avoid signal saturation and the resulting concentrations were conveniently adjusted.

2.4. Data Collection and Statistical Analysis

Two to four Petri dishes per genotype (nine genotypes) and four to five bud slices per Petri dish were cultured for each BA concentration and collection date. Contamination, survival and EFS (%) at each collection date were measured after two months of culture. When the axillary shoots were isolated and cultured individually in elongation medium, the EFS (%) and NS/E were calculated with respect to the non-contaminated explants. A logistic regression model was used to analyze the collection date, genotype and BA concentration's effects on the EFS (%). Tukey's post hoc test ($\alpha=0.05$) was used for multiple comparisons. Data on the NS/E were analyzed by analysis of variance (ANOVA). When necessary, multiple comparisons were made using Duncan's post hoc test ($\alpha=0.05$).

The RI percentage (%), the NR/E and the length of the longest root (LLR) (cm) were recorded after six weeks of culture in REM medium. A completely randomized design was carried out using eight stems per BA and sucrose concentrations and light treatments.

To assess the effect of the CK and sucrose concentration and light treatments' effects on the RI (%), a logistic regression was performed. Data on NR/E were x^2 transformed to meet homoscedasticity and were analyzed by ANOVA. Data on LLR (cm) were subjected to ANOVA. When necessary, multiple comparisons were made using Duncan's post hoc test ($\alpha=0.05$). The acclimatization percentage was calculated after four weeks under *ex vitro* conditions.

Data on free amino acid and carbohydrate content after four weeks of culture in RIM medium and after six weeks cultured in REM medium were analyzed. The confirmation of the homogeneity of variances and normality of the data was performed, and, when necessary, they were $\log(x)$ and \sqrt{x} transformed. Data on free amino acids and carbohydrates were subjected to ANOVA, and, when necessary, multiple comparisons were made using Duncan's post hoc test ($\alpha=0.05$). The data were analyzed using R Core Team software® (version 4.2.1, Vienna, Austria).

3. RESULTS

3.1. Organogenic Process

Genotype A showed the highest survival of the explants, independently of the collection date (72%). On the contrary, the lowest survival was observed in genotype J (43%). The highest survival values of the explants were observed on the first collection date (second week of January). In the same way, the highest contamination rates were recorded in samples collected in the third and fourth weeks of February (44 and 43%, respectively), and the lowest were found from January to the second week of February (between 17 and 34%).

The collection date had a significant effect on the EFS (%) (Figure 2, Supplementary Table S1). A significantly higher EFS percentage was obtained during the first week of February than during the third and fourth weeks of February (4%). Shoots induced in the second and third weeks of January and the second week of February displayed intermediate values (29 to 34%).




Figure 2. Explants forming shoots (%) shown in buds of *Pinus radiata* D. Don adult trees collected on different dates and cultured in LP medium [Quoirin and Lepoivre (1977) modified according to Aitken-Christie et al. (1988)]. Data are presented as mean values \pm S.E. Different letters indicate significant differences by Tukey's post hoc test ($p < 0.05$).

Reinvigorated shoots were obtained after culture media were supplemented with both BA concentrations (22 and 44 μ M). Regarding EFS (%), BA concentrations alone did not provoke any statistically significant differences in the EFS (%) (Supplementary Table S2), but the genotype and the interaction between genotype and BA concentration showed significant differences (Figure 3). A significantly higher EFS (%) was obtained in buds of genotype A induced with 44 μ M BA than in the rest of the genotypes and BA concentrations tested. Buds of genotype N cultured with 44 μ M BA showed a significantly lower EFS (%) than buds from genotypes A, F, G and J and genotypes E and J cultured at 22 and 44 μ M BA, respectively (between 33 and 45%, Figure 3).




Figure 3. Explants forming shoots (%) shown in buds of different genotypes of *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified according to Aitken-Christie et al. (1988)], supplemented with 6-benzyladenine (BA at 22 and 44 µM). Data are presented as mean values±S.E. Different letters indicate significant differences by Tukey's post hoc test ($p<0.05$).

No statistically significant differences were found in NS/E when the effect of the BA concentration in the culture media during the induction phase was analyzed (Supplementary Table S3). The NS/E ranged from 1.66 ± 0.40 to 1.95 ± 0.52 in explants induced with 22 and 44 µM BA, respectively.

3.2. Rooting Induction and Acclimatization of Rooted Plants

The different BA concentrations (22 and 44 µM) used in the SI phase did not have a statistically significant effect on RI (18% and 14%), on NR/E (4.50 and 3.56) or on LLR (1.83 cm and 1.60 cm) (Supplementary Tables S4 and S5).

A significantly higher RI (%) was observed in shoots cultured under white FL when compared with shoots growing under red LEDs (Figure 4A). In the same way, NR/E was significantly higher in shoots cultured under white FL than in those under red LEDs (Figure 4B). Light treatments did not provoke any statistically significant differences in the LLR. Shoots growing under red LEDs showed the longest length (2.22 cm) (Supplementary Tables S4 and S6).




Figure 4. Effect of light treatments (white fluorescent (F) and red LEDs (R)) on root induction (%) **(A)** and number of roots per explant **(B)** of *Pinus radiata* D. Don shoots cultured in LP medium [Quoirin and Lepoivre (1977) modified according to Aitken-Christie et al. (1988)]. Data are presented as mean values \pm S.E. Different letters indicate significant differences by Tukey's post hoc test (a) and by Duncan's post hoc test (b) ($p < 0.05$).

For the LLR, a significant effect of the interaction between BA and light treatment (Figure 5A) and between BA and sucrose concentration (Figure 5B) was observed. Explants induced with 44 μM BA and exposed to red LEDs displayed higher values compared with explants induced at the lowest BA concentration and exposed to the same light conditions (Figure 5A). On the contrary, explants cultured in the presence of 22 μM BA and exposed to FL displayed significantly higher values on LLR compared with explants induced at the highest BA concentration and exposed to the same light conditions (Figure 5A). Additionally, explants induced in the presence of 44 μM BA and cultured in medium supplemented with 1.5% sucrose during RI showed significantly higher values than shoots induced at the same BA concentration followed by a medium supplemented with 3.0% sucrose (Figure 5B). When a culture medium containing 22 μM BA was used for SI, regardless of the sucrose concentration evaluated, intermediate values of LLR were obtained (Figure 5B).




Figure 5. Effect of 6-benzyladenine (BA) concentration (22 and 44 μM), light treatment (white fluorescent (F) and red LEDs (R)) (A) and BA concentration (22 and 44 μM) and sucrose concentration [1.5% or 3.0% (w/v)] (B) on length of the longest root (cm) of *Pinus radiata* D. Don shoots cultured in LP medium [Quoirin and Lepoivre (1977) modified according to Aitken-Christie et al. (1988)]. Data are presented as mean values \pm S.E. Different letters indicate significant differences by Duncan's post hoc test ($p < 0.05$).

Regarding the sucrose concentrations, shoots cultured in medium with 3% sucrose showed a significantly higher RI value (22%) than those cultured in 1.5% sucrose (9%). No statistically significant differences in NR/E (3.73 and 4.10) and LLR values (2.01 and 1.55 cm) were observed in shoots cultured in the presence of 1.5 and 3% sucrose, respectively (Supplementary Tables S5 and S6). Additionally, the interaction between BA concentration and light treatment and between BA and sucrose concentrations assayed throughout SI did not show significant differences in RI (%) and NR/E (Supplementary Tables S4 and S5). In the same way, no statistically significant differences were observed in the interaction between light treatment and sucrose concentration evaluated during the SI phase in RI (%), NR/E and LLR (Supplementary Table S6).

Shoots induced in the presence of 22 μM BA, rooted in medium supplemented with 3% sucrose and exposed to FL from genotypes A, E and K developed *in vitro* roots (Figure 1E). In all genotypes, in the bases of shoots, a large and profuse callus was observed (Figure 1E). Acclimatized shoots were successfully obtained only from genotype K (33.33%) (Figure 1F).

3.3. Metabolite Analysis

BA concentrations and light treatments, as well as the interaction between both variables, did not show significant differences for the concentration of carbohydrates in shoots analyzed after four weeks under RIM treatment (Supplementary Tables S7-S9). However, explants cultured in media with the highest BA concentration presented a lower sucrose concentration than explants cultured in media with 22 μM BA (Table 1). Additionally, the levels of all carbohydrates analyzed were lower in explants cultured under red LEDs, except for the galactose concentration (Table 1)

Table 1. Effect of 6-benzyladenine (BA) concentration (22 and 44 μM) and light treatment (white fluorescent or red LEDs) on carbohydrate content of *Pinus radiata* D. Don shoots cultured in rooting medium [LP medium (Quoirin and Lepoivre, 1977) modified according to Aitken-Christie et al. (1988)], supplemented with a mixture of 5 μM 1-Naphthaleneacetic acid (NAA) and 10 μM indole-3-butyric acid (IBA).

Carbohydrates (μmolgFW^{-1})	Retention time (min)	BA concentration (μM)		Light treatment	
		22	44	White Fluorescent	Red LEDs
Fructose	18.076	21.36 \pm 6.28	22.71 \pm 2.90	24.91 \pm 5.21	19.20 \pm 2.59
Galactose	16.096	4.82 \pm 2.18	2.75 \pm 0.28	2.96 \pm 0.37	4.25 \pm 1.82
Glucose	14.653	13.66 \pm 4.37	15.50 \pm 2.17	16.70 \pm 3.57	12.66 \pm 2.19
Mannitol	22.911	7.59 \pm 0.45	7.04 \pm 0.44	7.36 \pm 0.36	7.15 \pm 0.55
Sucrose	12.640	12.29 \pm 3.82	5.67 \pm 0.83	10.73 \pm 3.30	5.70 \pm 0.99

Data are presented as mean values \pm S.E.

Regarding the effect of the sucrose concentration on the carbohydrate levels of shoots analyzed after RI treatment, no significant differences were found in galactose and mannitol concentrations (Supplementary Tables S8 and S9). Galactose displayed values from 2.41 to 4.85 μmolgFW^{-1} and mannitol from 6.69 to 7.87 μmolgFW^{-1} in media supplemented with 1.5 and 3% sucrose, respectively. In the case of fructose, glucose and sucrose, significantly higher concentrations were obtained in shoots cultured in the presence of the highest sucrose concentration when compared with the results at the lowest sucrose concentration (Figure 6A-C). The interaction between BA and light treatment, the interaction between BA and sucrose concentration and the interaction between light treatment and sucrose concentration did not show statistically significant differences (Supplementary Tables S7-S9).




Figure 6. Effect of sucrose concentration [1.5 or 3.0 (% w/v)] on fructose (A), glucose (B) and sucrose (C) concentration of *Pinus radiata* D. Don shoots cultured in rooting medium [LP medium (Quoirin and Lepoivre, 1977) modified according to Aitken-Christie et al. (1988)] supplemented with a mixture of 5 µM 1-Naphthaleneacetic acid (NAA) and 10 µM indole-3-butyric acid (IBA). Data are presented as mean values±S.E. Different letters indicate significant differences by Duncan's post hoc test ($p < 0.05$).

At the end of the rooting phase, no statistically significant differences were found for any of the carbohydrates analyzed in shoots during the root expression phase in relation to BA, light treatments, sucrose concentrations or the interactions among them (Supplementary Tables S10-S12). The highest glucose content (26.44 µmolgFW⁻¹) was found when shoots were cultured in the presence of the lowest BA concentration during the induction phase (Table 2). Additionally, shoots cultured under white FL exhibited an increase in fructose, glucose and sucrose levels compared with those cultured under red LEDs (Table 2). The fructose, glucose and sucrose levels decreased in shoots cultured during the induction phase in medium supplemented with the highest sucrose concentration. The galactose concentration in shoots after the rooting phase was the lowest of all the sugars analyzed, independently of the chemical characteristics of the culture media.

Table 2. Effect of 6-Benzyladenine (BA) (22 and 44 μM) and sucrose concentration [1.5 or 3.0 (% w/v)] and light treatment (white fluorescent or red LEDs) on carbohydrate content of *Pinus radiata* D. Don shoots cultured in LP medium [Quoirin and Lepoivre (1977) modified according to Aitken-Christie et al. (1988)] supplemented with 2 gL^{-1} activated charcoal after the rooting phase.

Carbohydrates (μmolgFW^{-1})	Retention time (min)	BA concentration (μM)		Light treatment		Sucrose concentration (w/v)	
		22	44	White Fluorescent	Red LEDs	1.5%	3.0%
Fructose	18.076	25.74 \pm 7.18	22.87 \pm 1.97	25.98 \pm 7.79	21.57 \pm 2.36	27.17 \pm 4.44	19.58 \pm 1.95
Galactose	16.096	3.12 \pm 0.59	3.87 \pm 0.59	4.00 \pm 0.47	3.19 \pm 0.75	3.42 \pm 0.66	3.86 \pm 0.53
Glucose	14.653	26.44 \pm 15.01	14.41 \pm 1.55	24.22 \pm 9.88	12.46 \pm 1.37	22.50 9.11	13.51 \pm 2.28
Mannitol	22.911	5.38 \pm 0.75	5.85 \pm 0.36	5.66 \pm 0.56	5.71 \pm 0.41	5.83 \pm 0.56	5.49 \pm 0.33
Sucrose	12.640	10.10 \pm 4.24	5.29 \pm 0.79	8.63 \pm 2.88	5.15 \pm 0.94	8.38 \pm 2.71	5.12 \pm 0.79

Data are presented as mean values \pm S.E.

Amino acid concentrations in shoots after RI were not significantly different in relation to BA, light treatments, sucrose concentrations or the interactions between them (Table 3 and Supplementary Tables S13-S15). Alanine, arginine, asparagine and glutamine levels were the highest in shoots after the rooting phase, regardless of the *in vitro* treatment applied during the induction phase (Table 3).

Table 3. Effect of 6-Benzyladenine (BA) (22 and 44 μM) and sucrose concentration [1.5 or 3.0 (% w/v)] and light treatment (white fluorescent or red LEDs) on amino acid content (μmolgFW^{-1}) of *Pinus radiata* D. Don shoots cultured in rooting medium [LP medium (Quoirin and Lepoivre, 1977) modified according to Aitken-Christie et al. (1988)] supplemented with mixture of 5 μM 1-Naphthaleneacetic acid (NAA) and 10 μM indole-3-butyric acid (IBA).

Free amino acids (μmolgFW^{-1})	Retention time (min)	BA concentration (μM)		Light treatment		Sucrose concentration (w/v)	
		22	44	White fluorescent	Red LEDs	1.5%	3.0%
Alanine	5.323	19.14 \pm 9.72	16.08 \pm 4.82	15.40 \pm 6.71	19.43 \pm 6.64	11.98 \pm 4.54	22.54 \pm 8.08
Arginine	5.035	29.98 \pm 10.39	28.22 \pm 7.34	34.10 \pm 8.77	22.8 \pm 7.72	22.45 \pm 6.74	35.34 \pm 9.64
Asparagine	3.115	13.72 \pm 8.27	24.78 \pm 8.18	16.09 \pm 7.67	25.70 \pm 9.39	30.53 \pm 9.68	10.52 \pm 6.05
Aspartic acid	1.204	0.57 \pm 0.12	0.61 \pm 0.07	0.68 \pm 0.10	0.50 \pm 0.05	0.60 \pm 0.08	0.59 \pm 0.10
Cysteine	7.140	3.47 \pm 0.82	4.36 \pm 0.55	4.61 \pm 0.67	3.32 \pm 0.60	3.95 \pm 0.79	4.07 \pm 0.53
Glutamic acid	1.849	9.31 \pm 8.07	5.55 \pm 2.64	11.78 \pm 6.17	1.42 \pm 0.10	3.60 \pm 2.04	10.39 \pm 6.54
Glutamine	3.827	52.78 \pm 16.66	70.16 \pm 8.58	65.37 \pm 12.38	61.26 \pm 11.19	63.54 \pm 12.57	63.41 \pm 11.29
Glycine	4.231	0.11 \pm 0.02	0.12 \pm 0.01	0.13 \pm 0.02	0.10 \pm 0.01	0.10 \pm 0.02	0.13 \pm 0.01
Hydroxyproline	10.330	0.02 \pm 0.01	0.03 \pm 0.01	0.02 \pm 0.01	0.03 \pm 0.01	0.03 \pm 0.01	0.03 \pm 0.01
Histidine	4.016	0.15 \pm 0.04	0.18 \pm 0.02	0.18 \pm 0.03	0.16 \pm 0.02	0.18 \pm 0.03	0.17 \pm 0.02
Isoleucine	9.079	0.08 \pm 0.01	0.10 \pm 0.01	0.09 \pm 0.01	0.09 \pm 0.01	0.10 \pm 0.01	0.09 \pm 0.01
Leucine	9.592	0.09 \pm 0.02	0.11 \pm 0.01	0.11 \pm 0.01	0.10 \pm 0.01	0.10 \pm 0.01	0.11 \pm 0.01
Lysine	10.018	0.14 \pm 0.03	0.17 \pm 0.02	0.16 \pm 0.03	0.16 \pm 0.01	0.17 \pm 0.02	0.14 \pm 0.02
Methionine	7.905	0.02 \pm 0.01	0.03 \pm 0.01	0.02 \pm 0.01	0.03 \pm 0.01	0.02 \pm 0.01	0.03 \pm 0.01
Phenylalanine	8.927	0.05 \pm 0.01	0.06 \pm 4.9 ⁻³	0.06 \pm 0.01	0.06 \pm 0.01	0.05 \pm 0.01	0.06 \pm 4.3 ⁻³
Proline	12.543	0.27 \pm 0.07	0.30 \pm 0.08	0.33 \pm 0.09	0.23 \pm 0.06	0.25 \pm 0.07	0.31 \pm 0.08
Serine	3.335	3.82 \pm 3.16	1.01 \pm 0.11	3.18 \pm 2.24	0.83 \pm 0.12	0.92 \pm 0.14	3.27 \pm 2.42
Threonine	4.386	0.12 \pm 0.02	0.15 \pm 0.01	0.14 \pm 0.02	0.12 \pm 0.01	0.13 \pm 0.02	0.14 \pm 0.02
Tryptophan	8.653	0.05 \pm 0.01	0.06 \pm 0.01	0.05 \pm 0.01	0.06 \pm 5.0 ⁻³	0.05 \pm 0.01	0.06 \pm 4.6 ⁻³
Tyrosine	6.334	0.06 \pm 0.01	0.09 \pm 0.01	0.08 \pm 0.01	0.08 \pm 0.01	0.08 \pm 0.01	0.08 \pm 0.01
Valine	7.760	0.29 \pm 0.07	0.40 \pm 0.04	0.38 \pm 0.05	0.33 \pm 0.04	0.32 \pm 0.05	0.39 \pm 0.05

Data are presented as mean values \pm S.E.

Shoots cultured with the highest sucrose concentration showed higher levels of alanine, arginine and glutamic acid than those cultured at the lowest sucrose concentration (Table 3). In contrast, the asparagine content was higher in rooted shoots coming from explants cultured in induction media supplemented with 1.5% sucrose than in those cultured with 3.0% sucrose.

After six weeks of culture in REM medium, no significant differences were found for the amino acid concentrations in shoots coming from explants induced in the presence of different concentrations of BA, light treatment and the interaction between both variables (Supplementary Tables S16–S18). However, the concentrations of alanine, arginine, glutamine, glycine, and lysine exhibited a decrease when shoots had been cultured with the highest BA concentration (Table 4). *P. radiata* shoots contained concentrations below 1.00 μmolgFW^{-1} of aspartic acid, glycine, hydroxyproline, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan, tyrosine and valine, regardless of the BA concentration used in the induction media (Table 4).

Table 4. Effect of 6-Benzyladenine (BA) (22 and 44 μM) and sucrose concentration [1.5 or 3.0 (% w/v)] and light treatment (white fluorescent or red LEDs) on amino acid levels (μmolgFW^{-1}) of *Pinus radiata* D. Don shoots cultured in LP medium [Quoirin and Lepoivre (1977) modified according to Aitken-Christie et al. (1988)] supplemented with 2 gL^{-1} activated charcoal after the rooting phase.

Free amino acids (μmolgFW^{-1})	Retention time (min)	BA concentration (μM)		Light treatment		Sucrose concentration (w/v)	
		22	44	White fluorescent	Red LEDs	1.5%	3.0%
Alanine	5.323	27.12±9.12	15.29±5.17	18.89±7.55	19.96±5.62	24.53±6.87	12.74±5.63
Arginine	5.035	67.60±24.44	63.97±15.2	79.68±19.55	49.47±16.80	68.70±15.72	60.71±23.06
Asparagine	3.115	0.76±0.09	3.98±3.30	0.73±0.07	5.17±4.50	4.56±3.80	0.64±0.06
Aspartic acid	1.204	0.81±0.08	0.87±0.08	0.82±0.04	0.89±0.12	0.90±0.09	0.78±0.06
Cystine	7.140	7.08±0.74	7.39±0.53	7.17±0.60	7.41±0.62	7.74±0.66	6.69±0.42
Glutamic acid	1.849	19.35±10.99	27.51±6.85	20.38±8.18	29.34±8.35	31.59±9.19	15.67±5.12
Glutamine	3.827	52.86±12.07	39.56±6.66	49.22±10.51	38.70±5.38	46.58±5.06	41.08±12.57
Glycine	4.231	0.20±0.02	0.18±0.02	0.18±0.02	0.19±0.03	0.20±0.03	0.17±0.01
Hydroxyproline	10.330	0.05±0.01	0.09±0.01	0.07±0.01	0.08±0.01	0.09±0.02	0.06±0.0047
Histidine	4.016	0.24±0.03	0.23±0.02	0.23±0.03	0.23±0.03	0.26±0.03	0.20±0.02
Isoleucine	9.079	0.17±0.01	0.17±0.02	0.16±0.01	0.18±0.02	0.19±0.02	0.15±0.01
Leucine	9.592	0.22±0.02	0.21±0.02	0.20±0.01	0.22±0.03	0.22±0.02	0.19±0.01
Lysine	10.018	0.41±0.16	0.25±0.04	0.36±0.11	0.25±0.04	0.30±0.04	0.32±0.13
Methionine	7.905	0.02±0.01	0.04±0.01	0.03±0.01	0.04±0.01	0.03±0.01	0.03±0.01
Phenylalanine	8.927	0.11±0.01	0.10±0.01	0.09±0.01	0.11±0.01	0.10±0.01	0.10±0.01
Proline	12.543	1.00±0.18	1.02±0.26	0.91±0.17	1.12±0.32	1.28±0.28	0.67±0.13
Serine	3.335	1.44±0.14	1.41±0.15	1.31±0.11	1.53±0.19	1.60±0.16	1.19±0.09
Threonine	4.386	0.23±0.02	0.24±0.02	0.24±0.01	0.24±0.03	0.27±0.02 ^a	0.21±0.01 ^b
Tryptophan	8.653	0.15±0.02	0.12±0.01	0.14±0.01	0.12±0.02	0.13±0.01	0.13±0.01
Tyrosine	6.334	0.14±0.01	0.13±0.01	0.13±4.0 ⁻³	0.14±0.01	0.15±0.01 ^a	0.12±0.01 ^b
Valine	7.760	0.63±0.09	0.56±0.06	0.56±0.05	0.61±0.09	0.63±0.07	0.52±0.06

Data are presented as mean values±S.E. Different letters indicate significant differences by Duncan's post hoc test ($p < 0.05$).

Arginine, glutamine and lysine concentrations were the highest in shoots growing under white FL. In contrast, concentrations of alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glycine, hydroxyproline, isoleucine, leucine, methionine, phenylalanine, proline, serine, tryptophan, tyrosine and valine increased in shoots exposed to red LEDs (Table 4).

Threonine and tyrosine concentrations in rooted shoots cultured in induction media with the lowest sucrose concentration were significantly higher than in shoots cultured in media with the highest sucrose concentration (Table 4). Although not significant, it is worth noting that alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, hydroxyproline, histidine, isoleucine, leucine, proline, serine and valine levels at the end of the rooting phase were also lower in shoots cultured in medium with the highest sucrose concentration during RI (Table 4).

Significantly higher values of hydroxyproline were found in shoots developed in a culture medium with 44 μM BA and 1.5% sucrose than in those from 44 μM BA and 3.0% sucrose. Additionally, the hydroxyproline levels were significantly higher in shoots cultured in a medium with 44 μM BA and 1.5% sucrose than in those from 22 μM BA and 1.5% sucrose (Figure 7). The lowest hydroxyproline level was obtained in shoots induced under the lowest BA concentration and cultured in medium supplemented with 1.5% sucrose during the RI phase (Figure 7).




Figure 7. Effect of 6-Benzyladenine (BA) (22 and 44 μM) and sucrose concentration [1.5 or 3.0 (% w/v)] on hydroxyproline amino acid level of *Pinus radiata* D. Don shoots cultured in LP medium [Quoirin and Lepoivre (1977) modified according to Aitken-Christie et al. (1988)] supplemented with 2 $\text{g}\cdot\text{L}^{-1}$ activated charcoal. Data are presented as mean values \pm S.E. Different letters indicate significant differences by Duncan's post hoc test ($p < 0.05$).

After the rooting expression phase, the interaction between light treatment and sucrose concentration showed statically significant differences for the aspartic acid, isoleucine, phenylalanine, threonine and tyrosine levels (Figures 8A-E). Significantly higher concentrations of the abovementioned amino acids were detected in shoots from medium with the lowest sucrose concentration and exposed to red LEDs (Figures 8A-E).




Figure 8. Effect of light treatment (white fluorescent (F) or red LEDs (R)) and sucrose concentration [1.5 or 3.0 (% w/v)] on aspartic acid (A), isoleucine (B), phenylalanine (C), threonine (D) and tyrosine (E) amino acid levels of *Pinus radiata* D. Don shoots cultured in LP medium [Quoirin and Lepoivre (1977) modified according to Aitken-Christie et al. (1988)] supplemented with 2 gL^{-1} activated charcoal. Data are presented as mean values \pm S.E. Different letters indicate significant differences by Duncan's post hoc test ($p < 0.05$).

Additionally, shoots coming from RI at 3.0% sucrose and exposed to red LEDs presented the lowest aspartic acid, isoleucine, threonine and tyrosine levels (Figure 8A-E).

4. DISCUSSION

In this study, the genotype affected the survival of the explants independently of the collection date. This result could be partially explained since the genotype is an endogenous factor that has a significant role in the regenerative potential, overall repeatability and reliability of the tissue culture protocol (Coleman and Ernst, 1989). As it was reviewed (Fehér, 2005; Jiménez, 2005), the response of explants to tissue culture conditions is highly dependent on the genetic and physiological determination. In this sense, *P. radiata*, *P. taeda*. and *P. halepensis* showed a differential organogenic response *in vitro* depending on the genotype in previous studies (Montalbán et al., 2013; Cézar et al., 2015 Pereira et al., 2021).

The highest survival and the lowest contamination rates were found during the first collection date. Several protocols mention that factors such as the genotype, explant source and PGRs influence the *in vitro* regeneration of plants via organogenesis or SE (Parzymies, 2021; Rojas-Vargas et al., 2021; Tung et al., 2021). In our case, the developmental stage of the explant can be related to the contamination rate. On the first collection date, explants consisted of apical shoot buds with closed scales; this source of explant could be better for the effective elimination of contamination. Analogous results were obtained in *Pseudotsuga menziesii* organogenesis (Traore et al., 2005), where spring buds showed higher contamination rates than winter buds.

Silver nanoparticles have various applications in plant biotechnology, mainly to eradicate microbial contamination and promote *in vitro* development (Pastelín-Solano et al., 2020). In this sense, the use of a silver nanoparticle solution in our study favored the disinfection efficiency of the explants, in agreement with the results for *Fragaria × ananassa*, *Vanilla planifolia* and *Psidium friedrichsthalianum*, where the contamination rates of 23, 28 and 40%, respectively, (Andújar et al., 2020; Pastelín-Solano et al., 2020; Tung et al., 2021) were reported. In contrast, in *P. halepensis*, a negative effect of the silver nanoparticle solution for the sterilization of apical shoot buds was found with a contamination rate of 89% (Pereira et al., 2021).

The timing of explant collection and the explant developmental stage influence the response to organogenesis or SE (Andersone and Levinsh, 2002; Corredoira et al., 2012). When the EFS was evaluated, apical buds collected in the first week of February showed higher percentages than those collected during the third and fourth weeks of February. A similar pattern has been reported in other species, such as *Quercus alba* and *Allamanda cathartica*, showing that the explant developmental stage or the season of collection of explants played an important role in the *in vitro* response (Corredoira et al., 2012; Khanam and Anis, 2018). During our study, a morphogenic peak from dormancy to bud initiation was observed in samples collected during January and February, wherein tissues became very active in terms of morphogenesis by increasing the shoot formation. These results are in accordance with those obtained in shoot explants of *Larix decidua* (Bonga, 2004, 2017), where two short morphogenic peaks were observed, the first before bud break and the second in late summer.

When we focused on the study of the effect of the BA concentration on EFS, no statistically significant differences were obtained. In plant tissue culture, BA is the most used cytokinin (Phillips and Garda, 2019); in addition, BA, used alone or in combination with other CKs, has been shown to efficiently promote *in vitro* bud induction (De Diego et al., 2008; Valverde-Cerdas et al., 2008). In our experiments, BA at the concentrations tested (22 and 44 μ M) provided a satisfactory organogenic response, and, although not significant, a slightly higher response in explants cultured in media supplemented with 44 μ M BA was observed. An analogous result was obtained in *P. halepensis* using the same BA concentration (Pereira et al., 2021). Likewise, in *P. pinea*, a higher number of buds formed per explant was observed during the first 16 days of culture at the highest concentration of BA tested (44.4 μ M) (Moncaleán et al., 2005). This behavior may be attributed to the maturity of the explant used, as, most likely, the aged buds are not as physiologically active as the young ones. For the above, higher concentrations of exogenously applied growth regulators may be needed in mature explants to obtain similar morphogenic responses, as reported for *Tetraclinis articulata* (Juan-Vicedo

et al., 2022). In this regard, Wendling et al. (2014) mentioned that, through the continuous *in vitro* subculture of shoots in media supplemented with CKs, reinvigoration can be obtained. In our study, all genotypes evaluated were able to develop reinvigorated shoots in media with both BA concentrations tested. These results agreed with those observed in *P. pinea*, *P. pinaster*, *P. silvestris* and *P. halepensis*, where reinvigorated axillary shoots were obtained through organogenesis, using buds as initial explants (De Diego et al., 2008; Cortizo et al., 2009; De Diego et al., 2010; Pereira et al., 2021).

In order to obtain a RI response in the explants, the culture medium was supplemented with NAA combined with IBA, a mixture traditionally used to induce root differentiation in *Pinus* species (Álvarez et al., 2009; Cortizo et al., 2009; Montalbán et al., 2011b, 2013). However, in this work, the rooting percentage was low, as it was observed in *P. pinea* when NAA alone was applied (Cortizo et al., 2009). Montalbán et al. (2011b), studying *P. radiata*, reported that IBA alone was more efficient in the adventitious RI of shoots coming from explants of juvenile origin. In *Pinus elliotii*, Nunes et al. (2018) observed that different concentrations and combinations of IBA (4.9 and 9.8 μM) and NAA (1.1, 2.1 and 2.7 μM) led to low root frequency using shoots developed from juvenile material. Likewise, in *P. halepensis*, long exposure to IBA was not effective in inducing roots in shoots of mature origin (Pereira et al., 2021). It is widely recognized that the most common problem encountered in the micropropagation of mature conifers is the adventitious root formation in shoots (Cortizo et al., 2009).

As reviewed by Bairu et al. (2011), the development of basal callus is one of the main physiological disorders that affects the rooting competence of microplants; this basal callus could interfere with physiological processes by trapping essential growth constituents such as PGRs and creating a physical barrier. In our work, in the bases of shoots, a large and profuse callus was observed, and this fact could have affected the development of adventitious roots. In contrast, in slash pine, when inducing adventitious roots using IBA combined with NAA, no callus formation was observed in shoots developed from juvenile explants (Nunes et al., 2018). Similarly, in *Pinus roxburghii* using shoots developed from juvenile explants, the RI medium supplemented with NAA and solidified with 0.6% agar produced shoots with low or no callusing at the base of the shoots (Kalia et al., 2007). In *P. taeda* (De Oliveira et al., 2012; C  zar et al., 2015), a combination of auxin (2.68 μM NAA) with CK (0.44 μM BA) was evaluated and showed RI of 55.6 to 47.5%, respectively. More recently, a pulse with IBA 500 mgL^{-1} for 5 min was used as an *ex vitro* rooting preconditioning treatment in *Mitragyna parvifolia*, obtaining approximately 90% micropropagated shoots rooted *ex vitro* (Patel et al., 2020). Soumare et al. (2021) reported that the use of plant-growth-promoting microbes, specifically *Streptomyces griseorubens* and *Norcardiopsis alba*, increased the rooting and root hair in maize. In this sense, these recent strategies, or physical factors such as the photoperiod, temperature and substrates could be considered to improve *P. radiata* rooting.

Carbohydrates in plants have several essential functions: they are basic elements of macromolecules, constitute substrates for respiration, play an important role in the synthesis pathway of many compounds and are indispensable for many other processes related to plant development or gene expression (Gibson, 2000; Calamar and De Klerk, 2002). In our work, the sucrose treatment had a significant effect on the RI percentage, and the highest result was observed in 3% (w/v) sucrose. Similar results were observed when comparing sucrose concentrations (1-9%) for the RI of apple rootstock, concluding that sucrose had a direct effect on rooting (Calamar and De Klerk, 2002). Likewise, *Musa* spp. and *Metroxylon sagu*, the highest values for rooting were found in a medium supplemented with 3% (w/v) sucrose (Sumaryono et al., 2012; Ahmed et al., 2014). In *Eucalyptus globulus* and *P. pinea*, glucose had a positive effect during RI (Corr  a et al., 2005; Zavattieri et al., 2009). Additionally, in *E. globulus* (Corr  a et al., 2005), it was reported that the carbohydrates are an energetic requirement for root development and sucrose, which is commonly used in tissue culture because it is the main sugar translocated in the phloem of several plants (Ragonezi et al., 2010; Kumar and Reddy, 2011).

The light type and wavelength specificity have an influence on morphogenetic responses such as adventitious root formation (Ragonezi et al., 2010). In several studies, LED lights have been used as an alternative to conventional lighting sources for plant tissue culture (Gupta and Jatothu, 2013). Supporting this, the spectral properties of LEDs have been found to regulate the morphological, anatomical and physiological responses of *in vitro* plants (Gupta and Jatothu, 2013). In this work, it was stated that the effectiveness of red LEDs on rooting depends on the genotype and the concentration of growth regulators applied in the RIM (Gupta and Jatothu, 2013). LED treatments (red or blue) showed a higher response in the rooting of plantlets in *Tripterospermum japonicum*, *Gossypium hirsutum* and *Doritaenopsis* spp. (Moon et al., 2006; Shin et al., 2008; Li et al., 2010). In *Anthurium andreaeanum*, root formation was progressively induced under a higher portion of red LEDs in a mixed circuit of blue and red (Budiarto, 2010). However, in this work, shoots grown under FL showed a significantly higher RI and NR/E. A similar pattern has been reported in *Handroanthus ochraceus*, *Achillea millefolium* and *Alocasia amazonica*, where higher values of RI were obtained in plants growing under low FL irradiances (Jo et al., 2008; Alvarenga et al., 2015; Larraburu et al., 2018). In future experiments to improve the micropropagation procedure in radiata pine a combination of different types of light should be evaluated. In this way, in *P. radiata* the used of blue LEDs improved the shoot height of somatic plants (Castander-Olarieta et al., 2023).

When studying the effect of the BA concentration and light treatment on LLR, explants induced with 44 μM BA and exposed to red LEDs showed the highest values. A similar pattern has been reported in *Vitis ficifolia*, where longer roots were obtained in plants growing under red LEDs (Poudel et al., 2008). In contrast, higher values of root length were obtained in plants of *H. ochraceus* exposed to low FL irradiances (Larraburu et al., 2018).

Metabolite analysis showed significant differences for several carbohydrate and amino acid concentrations in the different treatments tested. Carbohydrates in *in vitro* culture media have functions such as the maintenance of osmotic potential and to serve as a carbon source for developmental processes including RI (Yaseen et al., 2013; Moradi et al., 2017). When studying the effect of the sucrose concentration supplemented in RIM, higher fructose, glucose and sucrose content was found in shoots growing in culture medium supplemented with the highest sucrose concentration. In *A. amazonica*, media supplemented with 3 or 6.0% sucrose showed higher glucose, fructose and sucrose content in *in vitro* leaves compared with *ex vitro* leaves (Jo et al., 2009). Moreover, Mingozzi et al. (2011), studying *Cydonia oblonga*, concluded that the sucrose content in leaves was associated with its content in the culture media, and shoots cultured with 30 g L^{-1} sucrose showed higher content of sucrose, glucose and fructose. Additionally, Thorpe et al. (2008) explained that *in vitro* plants release invertase enzymes to the culture medium that act in the hydrolysis of sucrose, giving rise to glucose and fructose. In the present work, the levels of the carbohydrates were unaffected by lighting conditions. Contrasting our results, the content of sucrose, starch and soluble sugars was higher in plantlets of the *Doritaenopsis* spp. and *G. hirsutum* exposed to red plus blue and red LEDs, respectively (Shin et al., 2008; Li et al., 2010). Amino acid metabolism plays an essential role in plant protein biosynthesis, represents a building block for other biosynthesis pathways and is essential during signaling processes (Hildebrandt et al., 2015). Apart from the abovementioned functions, amino acids may also be involved in the plant stress response through intracellular pH regulation and the detoxification of reactive oxygen species (Campalans et al., 1999; Nuccio et al., 1999). Tyrosine is the precursor of secondary metabolites such as betacyanin in *Alternanthera brasiliana* (and certain enzymes such as tyrosine decarboxylase, which is responsible for redirecting essential primary metabolites into secondary metabolic pathways (Silva et al., 2005; Zhang et al., 2011). In addition, in *P. radiata* embryonal masses initiated under stressful conditions, tyrosine seems to be involved in antioxidative processes and osmotic adjustment (Castander-Olarieta et al., 2019). Likewise, the tyrosine metabolism pathway serves as a starting point to produce

tyrosine-derived metabolites essential to plant survival, such as tocopherols, plastoquinone and ubiquinone (Xu et al., 2020). Regarding threonine, it is an essential amino acid, and it is a substrate for isoleucine synthesis (Joshi et al., 2010). Moreover, threonine metabolites have an important role in plant growth and development, cell division and responses to abiotic stresses (Joshi et al., 2010; Muthuramalingam et al., 2018). However, in our study, the shoots with higher content of these two amino acids showed worse *in vitro* rooting (those coming from RIM with 1.5% (w/v) sucrose, at the end of the rooting phase).

Krasenski and Jonak (2012), Lukan et al. (2010) and Joshi et al. (2010) observed amino acid accumulation in plants exposed to abiotic stress, such as drought, extreme temperature, salinity and osmotic stress. In this sense, tobacco plants showed high levels of aromatic amino acids, and this fact was related to the improvement of salt stress tolerance (Oliva et al., 2021). In this study, our results suggest that low sucrose concentrations could be related to abiotic stress similar to that observed in nutrient deficiencies or carbohydrate starvation, such as the results observed in *Camellia sinensis* leaves, where increased threonine levels were observed under phosphate starvation (Santosh et al., 2018). However, the amino acid pool differs and changes depending on the developmental and physiological stage of the plant (Hildebrandt et al., 2015).

5. CONCLUSIONS

The regeneration of *P. radiata* from the vegetative shoot buds of adult trees via organogenesis was achieved, and it depended on environmental physico-chemical factors. The collection date and genotype had a strong effect on the efficiency of the process. Reinvigorated shoots from adult trees were obtained at both BA concentrations tested. The optimal result in terms of shoot induction was obtained at the highest BA concentration. Our results suggest that the use of white FL and a 3% sucrose concentration was better for root induction. The low root induction percentage suggests that further research should be encouraged to improve this phase, which is still a bottleneck in the micropropagation of mature conifers. Moreover, shoots showing better *in vitro* rooting had higher fructose, glucose and sucrose content, whereas those from the worse root treatment showed higher threonine and tyrosine levels.

6. REFERENCES

- Ahmed, S.; Sharma, A.; Bhushan, B.; Singh, A.K.; Wali, V.K. Effect of carbohydrate source, pH and supporting media on *in vitro* rooting of banana (*Musa* Spp.) cv. Grand naine plantlets. *Afr. J. Agric. Res.* **2014**, *9*, 1135–1140. <https://doi.org/10.5897/ajar2013.7285>.
- Aitken-Christie, J.; Singh, A.P.; Davis, H. Multiplication of meristematic tissue: A new tissue culture system for radiata pine. In *Genetic Manipulation of Woody Plants*; Hanover, J.W., Keathley, D.E., Eds.; Plenum Publishing Corp: New York, NY, USA, 1988; pp. 413–432. https://doi.org/10.1007/978-1-4613-1661-9_26.
- Allen, C.D.; Macalady, A.K.; Chenchouni, H.; Bachelet, D.; McDowell, N.; Vennetier, M.; Kitzberger, T.; Rigling, A.; Breshears, D.D.; Hogg (Ted), E.H.; et al. A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *For. Ecol. Manag.* **2010**, *259*, 660–684. <https://doi.org/10.1016/j.foreco.2009.09.001>.
- Alvarenga, I.C.A.; Pacheco, F.V.; Silva, S.T.; Bertolucci, S.K.V.; Pinto, J.E.B.P. *In vitro* culture of *Achillea millefolium* L.: Quality and intensity of light on growth and production of volatiles. *Plant Cell Tissue Organ Cult.* **2015**, *122*, 299–308. <https://doi.org/10.1007/s11240-015-0766-7>.

- Álvarez, J.M.; Majada, J.; Ordás, R.J. An improved micropropagation protocol for maritime pine (*Pinus pinaster* Ait.) isolated cotyledons. *Forestry* **2009**, *82*, 175–184. <https://doi.org/10.1093/forestry/cpn052>.
- Anderson, U.; Levinsh, G. Changes of morphogenic competence in mature *Pinus sylvestris* L. buds in vitro. *Ann. Bot.* **2002**, *90*, 293–298. <https://doi.org/10.1093/aob/mcf176>.
- Andújar, I.; González, N.; García-Ramos, J.C.; Bogdanchikova, N.; Pestryakov, A.; Escalona, M.; Concepción, O. Argovit™ Silver nanoparticles reduce contamination levels and improve morphological growth in the in vitro culture of *Psidium friedrichsthalianum* (O. Berg) Nied. *SN Appl. Sci.* **2020**, *2*, 2110. <https://doi.org/10.1007/s42452-020-03948-9>.
- Astolfi, S.; Marianello, C.; Grego, S.; Bellarosa, R. Preliminary investigation of LED lighting as growth light for seedlings from different tree species in growth chambers. *Not. Bot. Horti. Agrobot. Cluj-Napoca* **2012**, *40*, 31–38. <https://doi.org/10.15835/nbha4028221>.
- Bairu, M.W.; Kane, M.E. Physiological and developmental problems encountered by in vitro cultured plants. *Plant Growth Regul.* **2011**, *63*, 101–103. <https://doi.org/10.1007/s10725-011-9565-2>.
- Bonga, J.M. The effect of various culture media on the formation of embryo-like structures in cultures derived from explants taken from mature *Larix decidua*. *Plant Cell Tissue Organ Cult.* **2004**, *77*, 43–48. <https://doi.org/10.1023/B:TICU.0000016488.79965.b7>.
- Bonga, J.M. Can explant choice help resolve recalcitrance problems in in vitro propagation, a problem still acute especially for adult conifers? *Trees Struct. Funct.* **2017**, *31*, 781–789. <https://doi.org/10.1007/s00468-016-1509-z>.
- Budiarto, K. Spectral quality affects morphogenesis on *Anthurium* plantlet during in vitro culture. *Agrivita* **2010**, *32*, 234–240. <https://doi.org/10.17503/agrivita.v32i3.20>.
- Calamar, A.; De Klerk, G.J. Effect of sucrose on adventitious root regeneration in apple. *Plant Cell Tissue Organ Cult.* **2002**, *70*, 207–212. <https://doi.org/10.1023/A:1016356123367>.
- Campalans, A.; Messeguer, R.; Goday, A.; Pagès, M. Plant responses to drought, from ABA signal transduction events to the action of the induced proteins. *Plant Physiol. Biochem.* **1999**, *37*, 327–340. [https://doi.org/10.1016/S0981-9428\(99\)80039-4](https://doi.org/10.1016/S0981-9428(99)80039-4).
- Castander-Olarieta, A.; Montalbán, I.A.; De Medeiros Oliveira, E.; Dell'Aversana, E.; D'Amelia, L.; Carillo, P.; Steiner, N.; Fraga, H.P.F.; Guerra, M.P.; Goicoa, T.; et al. Effect of thermal stress on tissue ultrastructure and metabolite profiles during initiation of radiata pine somatic embryogenesis. *Front. Plant Sci.* **2019**, *9*, 1-16. <https://doi.org/10.3389/fpls.2018.02004>.
- Castander-Olarieta, A.; Pereira, C.; Montalbán, I.A.; Mendes, V.M.; Correia, S.; Suárez-Álvarez, S.; Manadas, B.; Canhoto, J.; Moncaleán, P. Proteome-wide analysis of heat-stress in *Pinus radiata* somatic embryos reveals a combined response of sugar metabolism and translational regulation mechanisms. *Front. Plant Sci.* **2021**, *12*, 631239. <https://doi.org/10.3389/fpls.2021.631239>.
- Castander-Olarieta, A.; Montalbán, I. A.; Moncaleán, P. Multi-strategy approach towards optimization of maturation and germination in radiata pine somatic embryogenesis. *Plant Cell Tissue Organ Cult.* **2023**, *153*, 173-190. <https://doi.org/10.1007/s11240-023-02457-y>.

- Cézar, T.M.; Higa, A.R.; Koehler, H.S.; Ribas, L.L.F. Influence of culture medium, explant length and genotype on micropropagation of *Pinus taeda* L. *Ciênc. Florest.* **2015**, *25*, 13–22. <https://doi.org/10.1590/1980-509820152505013>.
- Coleman, G.D.; Ernst, S.G. In vitro shoot regeneration of *Populus deltoides*: Effect of cytokinin and genotype. *Plant Cell Rep.* **1989**, *8*, 459–462. <https://doi.org/10.1007/BF00269048>.
- Corrêa, L.D.R.; Paim, D.C.; Schwambach, J.; Fett-Neto, A.G. Carbohydrates as regulatory factors on the rooting of *Eucalyptus saligna* Smith and *Eucalyptus globulus* Labill. *Plant Growth Regul.* **2005**, *45*, 63–73. <https://doi.org/10.1007/s10725-004-6125-z>.
- Corredoira, E.; San-José, M.C.; Vieitez, A.M. Induction of somatic embryogenesis from different explants of shoot cultures derived from young *Quercus alba* Trees. *Trees Struct. Funct.* **2012**, *26*, 881–891. <https://doi.org/10.1007/s00468-011-0662-7>.
- Cortizo, M.; De Diego, N.; Moncaleán, P.; Ordás, R.J. Micropropagation of adult stone pine (*Pinus pinea* L.). *Trees Struct. Funct.* **2009**, *23*, 835–842. <https://doi.org/10.1007/s00468-009-0325-0>.
- de Almeida, M.; de Almeida, C.V.; Graner, E.M.; Brondani, G.E.; de Abreu-Tarazi, M.F. Pre-procambial cells are niches for pluripotent and totipotent stem-like cells for organogenesis and somatic embryogenesis in the peach palm: A histological study. *Plant Cell Rep.* **2012**, *31*, 1495–1515. <https://doi.org/10.1007/s00299-012-1264-6>.
- De Diego, N.; Montalbán, I.A.; Fernandez de Larrinoa, E.; Moncaleán, P. In vitro regeneration of *Pinus pinaster* adult trees. *Can. J. For. Res.* **2008**, *38*, 2607–2615. <https://doi.org/10.1139/X08-102>.
- De Diego, N.; Montalbán, I.A.; Moncaleán, P. In vitro regeneration of adult *Pinus sylvestris* L. Trees. *S. Afr. J. Bot.* **2010**, *76*, 158–162. <https://doi.org/10.1016/j.sajb.2009.09.007>.
- De Oliveira, L.F.; Ribas, L.L.F.; Quoirin, M.; Koehler, H.S.; Amano, E.; Higa, A.R. Micropropagation of *Pinus taeda* L. from juvenile material. *Tree For. Sci. Biotechnol.* **2012**, *6*, 96–101.
- Escandón, M.; Valledor, L.; Pascual, J.; Pinto, G.; Cañal, M.J.; Meijón, M. System-wide analysis of short-term response to high temperature in *Pinus radiata*. *J. Exp. Bot.* **2017**, *68*, 3629–3641. <https://doi.org/10.1093/jxb/erx198>.
- Farjon, A. The Kew review: Conifers of the world. *Kew Bull.* **2018**, *73*, 8. <https://doi.org/10.1007/s12225-018-9738-5>.
- Fehér, A. Why somatic plant cells start to form embryos? In *Somatic Embryogenesis*; Mujib, A., Šamaj, J., Eds.; Plant Cell Monographs; Springer: Berlin/Heidelberg, Germany, 2005; Volume 2, pp. 85–101. https://doi.org/10.1007/7089_019.
- Gibson, S.I. Plant sugar-response pathways. part of a complex regulatory web. *Plant Physiol.* **2000**, *124*, 1532–1539. <https://doi.org/10.1104/pp.124.4.1532>.
- Gupta, S.; Jatothu, B. Fundamentals and applications of light-emitting diodes (LEDs) in in vitro plant growth and morphogenesis. *Plant Biotechnol. Rep.* **2013**, *7*, 211–220. <https://doi.org/10.1007/s11816-013-0277-0>.
- Hildebrandt, T.M.; Nunes Nesi, A.; Araújo, W.L.; Braun, H.P. Amino acid catabolism in plants. *Mol. Plant.* **2015**, *8*, 1563–1579. <https://doi.org/10.1016/j.molp.2015.09.005>.

- Jankovský, L.; Palovčíková, D.; Dvořák, M.; Tomšovský, M. Records of Brown spot needle blight related to *Lecanosticta acicola* in the Czech Republic. *Plant Prot. Sci.* **2009**, *45*, 16–18. <https://doi.org/10.17221/39/2008-pps>.
- Jao, R.C.; Fang, W. An adjustable light source for photo-phyto related research and young plant production. *Appl. Eng. Agric.* **2003**, *19*, 601–608. <https://doi.org/10.13031/2013.15317>.
- Jiménez, V.M. Involvement of plant hormones and plant growth regulators on in vitro somatic embryogenesis. *Plant Growth Regul.* **2005**, *47*, 91–110. <https://doi.org/10.1007/s10725-005-3478-x>.
- Jo, E.A.; Tewari, R.K.; Hahn, E.J.; Paek, K.Y. Effect of photoperiod and light intensity on in vitro propagation of *Alocasia amazonica*. *Plant Biotechnol. Rep.* **2008**, *2*, 207–212. <https://doi.org/10.1007/s11816-008-0063-6>.
- Jo, E.A.; Tewari, R.K.; Hahn, E.J.; Paek, K.Y. In vitro sucrose concentration affects growth and acclimatization of *Alocasia amazonica* plantlets. *Plant Cell Tissue Organ Cult.* **2009**, *96*, 307–315. <https://doi.org/10.1007/s11240-008-9488-4>.
- Joshi, V.; Joung, J.G.; Fei, Z.; Jander, G. Interdependence of threonine, methionine and isoleucine metabolism in plants: Accumulation and transcriptional regulation under abiotic stress. *Amino acids* **2010**, *39*, 933–947. <https://doi.org/10.1007/s00726-010-0505-7>.
- Juan-Vicedo, J.; Serrano-Martinez, F.; Cano-Castillo, M.; Casas, J.L. In vitro propagation, genetic assessment, and medium-term conservation of coastal endangered species *Tetraclinis articulata* (Vahl) Masters (Cupressaceae) from adult trees. *Plants* **2022**, *11*, 187. <https://doi.org/10.3390/plants11020187>.
- Kalia, R.K.; Arya, S.; Kalia, S.; Arya, I.D. Plantlet regeneration from fascicular buds of seedling shoot apices of *Pinus roxburghii* Sarg. *Biol. Plant.* **2007**, *51*, 653–659. <https://doi.org/10.1007/s10535-007-0138-1>.
- Khanam, M.N.; Anis, M. Organogenesis and efficient in vitro plantlet regeneration from nodal segments of *Allamanda cathartica* L. using TDZ and ultrasound assisted extraction of quercetin. *Plant Cell Tissue Organ Cult.* **2018**, *134*, 241–250. <https://doi.org/10.1007/s11240-018-1416-7>.
- Krasensky, J.; Jonak, C. Drought, salt, and temperature stress-induced metabolic rearrangements and regulatory networks. *J. Exp. Bot.* **2012**, *63*, 1593–1608. <https://doi.org/10.1093/jxb/err460>.
- Kulus, D.; Woźny, A. Influence of light conditions on the morphogenetic and biochemical response of selected ornamental plant species under in vitro conditions: A mini-review. *Bio. Technologia* **2020**, *101*, 75–83. <https://doi.org/10.5114/bta.2020.92930>.
- Kumar, N.; Reddy, M.P. In vitro plant propagation: A review. *J. For. Environ. Sci.* **2011**, *27*, 61–72. <https://doi.org/10.7747/JFS.2011.27.2.1>.
- Larraburu, E.E.; Correa, G.S.; Llorente, B.E. In vitro development of yellow lapacho (Bignoniaceae) using high-power light emitting diode. *Rev. Arvore.* **2018**, *42*, e420508. <https://doi.org/10.1590/1806-90882018000500008>.
- Li, H.; Xu, Z.; Tang, C. Effect of light-emitting diodes on growth and morphogenesis of upland cotton (*Gossypium hirsutum* L.) plantlets in vitro. *Plant Cell Tissue Organ Cult.* **2010**, *103*, 155–163. <https://doi.org/10.1007/s11240-010-9763-z>.

- Lugan, R.; Niogret, M.F.; Leport, L.; Guégan, J.P.; Larher, F.R.; Savouré, A.; Kopka, J.; Bouchereau, A. Metabolome and water homeostasis analysis of *Thellungiella salsuginea* suggests that dehydration tolerance is a key response to osmotic stress in this halophyte. *Plant J.* **2010**, *64*, 215–229. <https://doi.org/10.1111/j.1365-313X.2010.04323.x>.
- Mingozi, M.; Morini, S.; Lucchesini, M.; Mensuali-Sodi, A. Effects of leaf soluble sugars content and net photosynthetic rate of quince donor shoots on subsequent morphogenesis in leaf explants. *Biol. Plant.* **2011**, *55*, 237–242. <https://doi.org/10.1007/s10535-011-0034-6>.
- Moncaleán, P.; Alonso, P.; Centeno, M.L.; Cortizo, M.; Rodríguez, A.; Fernández, B.; Ordás, R.J. Organogenic responses of *Pinus pinea* cotyledons to hormonal treatments: BA metabolism and cytokinin content. *Tree Physiol.* **2005**, *25*, 1–9. <https://doi.org/10.1093/treephys/25.1.1>.
- Montalbán, I.A.; De Diego, N.; Igartua, E.A.; Setién, A.; Moncaleán, P. A combined pathway of somatic embryogenesis and organogenesis to regenerate radiata pine plants. *Plant Biotechnol. Rep.* **2011a**, *5*, 177–186. <https://doi.org/10.1007/s11816-011-0171-6>.
- Montalbán, I.A.; De Diego, N.; Moncaleán, P. Testing novel cytokinins for improved in vitro adventitious shoots formation and subsequent ex vitro performance in *Pinus radiata*. *Forestry* **2011b**, *84*, 363–373. <https://doi.org/10.1093/forestry/cpr022>.
- Montalbán, I.A.; Novák, O.; Rolčík, J.; Strnad, M.; Moncaleán, P. Endogenous cytokinin and auxin profiles during in vitro organogenesis from vegetative buds of *Pinus radiata* Adult Trees. *Physiol. Plant.* **2013**, *148*, 214–231. <https://doi.org/10.1111/j.1399-3054.2012.01709.x>.
- Moon, H.K.; Park, S.Y.; Kim, Y.W.; Kim, C.S. Growth of Tsuru-Rindo (*Tripterospermum japonicum*) cultured in vitro under various sources of light-emitting diode (LED) irradiation. *J. Plant Biol.* **2006**, *49*, 174–179. <https://doi.org/10.1007/BF03031014>.
- Moradi, P.; Ford-Lloyd, B.; Pritchard, J. Metabolomic approach reveals the biochemical mechanisms underlying drought stress tolerance in thyme. *Anal. Biochem.* **2017**, *527*, 49–62. <https://doi.org/10.1016/j.ab.2017.02.006>.
- Muthuramalingam, P.; Krishnan, S.R.; Pandian, S.; Mareeswaran, N.; Aruni, W.; Pandian, S.K.; Ramesh, M. Global analysis of threonine metabolism genes unravel key players in rice to improve the abiotic stress tolerance. *Sci. Rep.* **2018**, *8*, 9270. <https://doi.org/10.1038/s41598-018-27703-8>.
- Narváez, I.; Martín, C.; Jiménez-Díaz, R.M.; Mercado, J.A.; Pliego-Alfaro, F. Plant regeneration via somatic embryogenesis in mature wild olive genotypes resistant to the defoliating pathotype of *Verticillium dahliae*. *Front. Plant Sci.* **2019**, *10*, 1471. <https://doi.org/10.3389/fpls.2019.01471>.
- Neale, D.B.; Wheeler, N.C. The Conifers. In *The Conifers: Genomes, Variation and Evolution*; Springer Nature: Cham, Switzerland, 2019; pp. 1–21. https://doi.org/10.1007/978-3-319-46807-5_1.
- Nuccio, M.L.; Rhodes, D.; McNeil, S.D.; Hanson, A.D. Metabolic engineering of plants for osmotic stress resistance. *Curr. Opin. Plant Biol.* **1999**, *2*, 128–134. [https://doi.org/10.1016/S1369-5266\(99\)80026-0](https://doi.org/10.1016/S1369-5266(99)80026-0).
- Nunes, S.; Sousa, D.; Pereira, V.T.; Correia, S.; Marum, L.; Santos, C.; Dias, M.C. Efficient protocol for in vitro mass micropropagation of slash pine. *In Vitro Cell. Dev. Biol. Plant.* **2018**, *54*, 175–183. <https://doi.org/10.1007/s11627-018-9891-4>.

Oliva, M.; Guy, A.; Galili, G.; Dor, E.; Schweitzer, R.; Amir, R.; Hacham, Y. Enhanced production of aromatic amino acids in tobacco plants leads to increased phenylpropanoid metabolites and tolerance to stresses. *Front. Plant Sci.* **2021**, *11*, 604349. <https://doi.org/10.3389/fpls.2020.604349>.

Parzymies, M. Nano-silver particles reduce contaminations in tissue culture but decrease regeneration rate and slows down growth and development of *Aldrovanda vesiculosa* explants. *Appl. Sci.* **2021**, *11*, 3653. <https://doi.org/10.3390/app11083653>.

Pastelín-Solano, M.C.; Ramírez-Mosqueda, M.A.; Bogdanchikova, N.; Castro-González, C.G.; Bello-Bello, J.J. Silver nanoparticles affect the micropropagation of vanilla (*Vanilla planifolia* Jacks. Ex Andrews). *Agrociencia* **2020**, *54*, 1–13.

Patel, A.K.; Lodha, D.; Shekhawat, N.S. An improved micropropagation protocol for the *ex situ* conservation of *Mitragyna parvifolia* (Roxb.)Korth (Rubiaceae): An endangered tree of pharmaceutical importance. *In Vitro Cell. Dev. Biol. Plant.* **2020**, *56*, 817–826. <https://doi.org/10.1007/s11627-020-10089-6>.

Pereira, C.; Montalbán, I.A.; Pedrosa, A.; Tavares, J.; Pestryakov, A.; Bogdanchikova, N.; Canhoto, J.; Moncaleán, P. Regeneration of *Pinus halepensis* (Mill.) through organogenesis from apical shoot buds. *Forests* **2021**, *12*, 363. <https://doi.org/10.3390/f12030363>.

Phillips, G.C. In vitro morphogenesis in plants—Recent advances. *In Vitro Cell. Dev. Biol. Plant.* **2004**, *40*, 342–345. <https://doi.org/10.1079/IVP2004555>.

Phillips, G.C.; Garda, M. Plant tissue culture media and practices: An overview. *In Vitro Cell. Dev. Biol. Plant.* **2019**, *55*, 242–257. <https://doi.org/10.1007/s11627-019-09983-5>.

Poudel, P.R.; Kataoka, I.; Mochioka, R. Effect of red- and blue-light-emitting diodes on growth and morphogenesis of grapes. *Plant Cell Tissue Organ Cult.* **2008**, *92*, 147–153. <https://doi.org/10.1007/s11240-007-9317-1>.

Quoirin, M.; Lepoivre, P. Études des milieux adaptés aux cultures in vitro de *Prunus*. *Acta Hortic.* **1977**, *78*, 437–442. <https://doi.org/10.17660/ActaHortic.1977.78.54>.

Ragonezi, C.; Klimaszewska, K.; Castro, M.R.; Lima, M.; de Oliveira, P.; Zavattieri, M.A. Adventitious rooting of conifers: Influence of physical and chemical factors. *Trees Struct. Funct.* **2010**, *24*, 975–992. <https://doi.org/10.1007/s00468-010-0488-8>.

Reed, B.M.; Sarasan, V.; Kane, M.; Bunn, E.; Pence, V.C. Biodiversity conservation and conservation biotechnology tools. *In Vitro Cell. Dev. Biol. Plant.* **2011**, *47*, 1–4. <https://doi.org/10.1007/s11627-010-9337-0>.

Riikonen, J.; Kettunen, N.; Gritsevich, M.; Hakala, T.; Särkkä, L.; Tahvonen, R. Growth and development of Norway spruce and Scots pine seedlings under different light spectra. *Environ. Exp. Bot.* **2016**, *121*, 112–120. <https://doi.org/10.1016/j.envexpbot.2015.06.006>.

Rojas-Vargas, A.; Castander-Olarieta, A.; Moncaleán, P.; Montalbán, I.A. Optimization of the micropropagation of elite adult trees of *Sequoia sempervirens*: Forest species of interest in the Basque Country, Spain. *Rev. Bionatura.* **2021**, *6*, 1511–1519. <https://doi.org/10.21931/RB/2021.01.01.11>.

Santosh, K.C.; Liu, M.; Zhang, Q.; Fan, K.; Shi, Y.; Ruan, J. Metabolic Changes of Amino acids and Flavonoids in Tea Plants in Response to Inorganic Phosphate Limitation. *Int. J. Mol. Sci.* **2018**, *19*, 3683. <https://doi.org/10.3390/ijms19113683>.

Shin, K.S.; Murthy, H.N.; Heo, J.W.; Hahn, E.J.; Paek, K.Y. The effect of light quality on the growth and development of in vitro cultured *Doritaenopsis* plants. *Acta Physiol. Plant.* **2008**, *30*, 339–343. <https://doi.org/10.1007/s11738-007-0128-0>.

Silva, N.C.B.; Macedo, A.F.; Lage, C.L.S.; Esquibel, M.A.; Sato, A. Developmental effects of additional ultraviolet a radiation, growth regulators and tyrosine in *Alternanthera brasiliana* (L.) Kuntze cultured in vitro. *Braz. Arch. Biol. Technol.* **2005**, *48*, 779–786. <https://doi.org/10.1590/S1516-89132005000600014>.

Soumare, A.; Boubekri, K.; Lyamlouli, K.; Hafidi, M.; Ouhdouch, Y.; Kouisni, L. Efficacy of phosphate solubilizing actinobacteria to improve rock phosphate agronomic effectiveness and plant growth promotion. *Rhizosphere* **2021**, *17*, 100284. <https://doi.org/10.1016/j.rhisph.2020.100284>.

Stephano-Hornedo, J.L.; Torres-Gutiérrez, O.; Toledano-Magaña, Y.; Gradilla-Martínez, I.; Pestryakov, A.; Sánchez-González, A.; García-Ramos, J.C.; Bogdanchikova, N. Argovit™ silver nanoparticles to fight Huanglongbing disease in Mexican limes (*Citrus aurantifolia* Swingle). *RSC Adv.* **2020**, *10*, 6146–6155. <https://doi.org/10.1039/c9ra09018e>.

Sumaryono, W.M.; Ratnadewi, D. Effect of carbohydrate source on growth and performance of in vitro sago palm (*Metroxylon sagu* Rottb.) plantlets. *HAYATI J. Biosci.* **2012**, *19*, 88–92. <https://doi.org/10.4308/hjb.19.2.88>.

Thorpe, T.; Stasolla, C.; Yeung, E.C.; de Klerk, G.J.; Roberts, A.; George, E.F. The components of plant tissue culture media II: Organic additions, osmotic and pH effects, and support systems. In *Plant Propagation by Tissue Culture*, 3rd ed.; George, E.F., Hall, M.A., de Klerk, G.J., Eds.; Springer: Dordrecht, The Netherlands, 2008; pp. 115–173.

Traore, A.; Xing, Z.; Bonser, A.; Carlson, J. Optimizing a protocol for sterilization and in vitro establishment of vegetative buds from mature douglas fir trees. *J. Am. Soc. Hortic. Sci.* **2005**, *40*, 1464–1468. <https://doi.org/10.21273/hortsci.40.5.1464>.

Tung, H.T.; Thuong, T.T.; Cuong, D.M.; Luan, V.Q.; Hien, V.T.; Hieu, T.; Nam, N.B.; Phuong, H.T.N.; Van The Vinh, B.; Khai, H.D.; et al. Silver nanoparticles improved explant disinfection, in vitro growth, runner formation and limited ethylene accumulation during micropropagation of strawberry (*Fragaria* × *Ananassa*). *Plant Cell Tissue Organ Cult.* **2021**, *145*, 393–403. <https://doi.org/10.1007/s11240-021-02015-4>.

Valledor, L.; Escandón, M.; Meijón, M.; Nukarinen, E.; Cañal, M.J.; Weckwerth, W. A universal protocol for the combined isolation of metabolites, DNA, long RNAs, small RNAs, and proteins from plants and microorganisms. *Plant J.* **2014**, *79*, 173–180. <https://doi.org/10.1111/tpj.12546>.

Valverde-Cerdas, L.; Rojas-Vargas, A.; Hine-Gómez, A. In vitro propagation of *Albizia guachapele*, *Cedrela odorata*, *Platymiscium pinnatum* and *Guaicum sanctum*. *Plant Tissue Cult. Biotechnol.* **2008**, *18*, 151–156. <https://doi.org/10.3329/ptcb.v18i2.3397>.

Wendling, I.; Trueman, S.J.; Xavier, A. Maturation and related aspects in clonal forestry-part II: Reinvigoration, rejuvenation and juvenility maintenance. *New For.* **2014**, *45*, 473–486. <https://doi.org/10.1007/s11056-014-9415-y>.

Woods, A.J.; Martín-García, J.; Bulman, L.; Vasconcelos, M.W.; Boberg, J.; La Porta, N.; Peredo, H.; Vergara, G.; Ahumada, R.; Brown, A.; et al. *Dothistroma* needle blight, weather and possible climatic triggers for the disease's recent emergence. *For. Pathol.* **2016**, *46*, 443–452. <https://doi.org/10.1111/efp.12248>.

Xu, J.J.; Fang, X.; Li, C.Y.; Yang, L.; Chen, X.Y. General and Specialized Tyrosine Metabolism Pathways in Plants. *Abiotech* **2020**, *1*, 97–105. <https://doi.org/10.1007/s42994-019-00006-w>.

Yaseen, M.; Ahmad, T.; Sablok, G.; Standardi, A.; Hafiz, I.A. Review: Role of carbon sources for in vitro plant growth and development. *Mol. Biol. Rep.* **2013**, *40*, 2837–2849. <https://doi.org/10.1007/s11033-012-2299-z>.

Zavattieri, A.; Lima, M.; Sobral, V.; Oliveira, P.; Costa, A. Effects of carbon source, carbon concentration and culture conditions on in vitro rooting of *Pinus pinea* L. microshoots. *Acta Hort.* **2009**, *812*, 173–180. <https://doi.org/10.17660/ActaHortic.2009.812.19>.

Zhang, J.X.; Ma, L.Q.; Yu, H.S.; Zhang, H.; Wang, H.T.; Qin, Y.F.; Shi, G.L.; Wang, Y.N. A tyrosine decarboxylase catalyzes the initial reaction of the salidroside biosynthesis pathway in *Rhodiola sachalinensis*. *Plant Cell Rep.* **2011**, *30*, 1443–1453. <https://doi.org/10.1007/s00299-011-1053-7>

CHAPTER 4

Adult trees *Cryptomeria japonica* (Thunb. ex L.f.) D. Don Micropropagation: Factors Involved in the Success of the Process

The content of this chapter corresponds to the published article “Rojas-Vargas, A.; Montalbán, I.A.; Moncaleán, P. 2023. Adult Trees *Cryptomeria japonica* (Thunb. ex L.f.) D. Don Micropropagation: Factors Involved in the Success of the Process. *Forests* 14, 743. <https://doi.org/10.3390/f14040743>”.

ABSTRACT

Cryptomeria japonica (Thunb. ex L.f.) D. Don is a commercial tree native to Japan and is one of the most important forest species in that country and the Azores (Portugal). Because of the quality of *C. japonica* timber, several genetic improvement programs have been performed. Recently, some studies focusing on *C. japonica* SE have been carried out (Maruyama et al., 2021). However, in this species, this process uses immature seeds as initial explants, and for this reason, it is not possible to achieve the maximum genetic gain (100% genetic of the donor plant). Although some studies have been made applying organogenesis to this species, the success of the process in adult trees is low. For this reason, our main goal was to optimize the micropropagation method by using trees older than 30 years as a source of plant material. In this sense, in a first experiment, we studied the effect of different types of initial explants and three basal culture media on SI; then, two sucrose concentrations and two light treatments (LEDs versus FL) were tested for the improvement of rooting. In a second experiment, the effects of different PGRs (BA, m-T, TDZ) on SI and the subsequent phases of the organogenesis process were analyzed. The cultures produced the highest number of shoots when QL medium (Quoirin and Lepoivre, 1977) and long basal explants (>1.5 cm) were used; the shoots obtained produced a higher number of roots when they were grown under red LEDs. Moreover, RI was significantly higher in shoots previously induced with m-T.

RESUMEN

Cryptomeria japonica (Thunb. ex L.f.) D. Don es un árbol originario de Japón y es una de las especies forestales más importantes de ese país y de las Azores (Portugal). Debido a la calidad de la madera de *C. japonica*, se han llevado a cabo varios programas de mejora genética. Recientemente, también se ha abordado el desarrollo de estudios centrados en la embriogénesis somática (Maruyama et al., 2021). Sin embargo, en esta especie, este proceso utiliza semillas inmaduras como explantos iniciales, por lo que no es posible lograr la máxima ganancia genética (100% genotipo del árbol donador). Aunque se han realizado algunos estudios aplicando organogénesis a esta especie, el éxito del proceso en árboles adultos es bajo. Por esta razón, nuestro principal objetivo fue optimizar el método de micropropagación utilizando árboles mayores de 30 años como fuente inicial de material vegetal. En este sentido, en un primer experimento estudiamos el efecto de diferentes tipos de explantos iniciales y tres medios de cultivo basales sobre la inducción de brotes; luego, se probaron dos concentraciones de sacarosa y dos tratamientos de luz (LEDs versus FL) para mejorar el proceso de enraizamiento. En un segundo experimento, se analizaron los efectos de diferentes reguladores del crecimiento vegetal (BA, m-T, TDZ) sobre la inducción de brotes y las fases posteriores del proceso de organogénesis. El mayor número de brotes se obtuvo cuando los explantos basales largos (>1,5 cm) se desarrollaron en el medio de cultivo QL (Quoirin y Lepoivre, 1977); los brotes obtenidos produjeron un mayor número de raíces cuando se cultivaron bajo LEDs rojas. Además, la inducción de raíces fue significativamente mayor en los brotes previamente inducidos con m-T.

1. INTRODUCTION

The Japanese cedar, *Cryptomeria japonica* (Thunb. ex L.f.) D. Don, subfamily Taxodioideae, family Cupressaceae, is a monoecious conifer distributed across East Asia (Kusumi et al., 2000; Onuma et al., 2023). *C. japonica* covers approximately 4.5 million ha, representing 44% of the total reforested area (Forestry Agency, 2020). This conifer is one of the most important timber tree species in Japan, and it is traditionally used for construction wood and for obtaining biomass (Taniguchi et al., 2020).

Tsubomura and Taniguchi (2008) mentioned that Japanese cedar is clonally propagated by cuttings, but this type of propagation requires many hours of manual labor, therefore it is difficult to establish a short-term propagation protocol. For the abovementioned reasons, biotechnological approaches for *C. japonica* clonal propagation, such as *in vitro* methods including SE or shoot organogenesis, are valuable tools for the propagation of this conifer (Phillips, 2004). In this sense, SE is a recognized technique for the large-scale propagation of conifers (Park et al., 1998). However, SE is a complex, multistage process initiated from immature seeds, so it is not possible to reproduce the genotype of the donor plant. In contrast, micropropagation by nodal tissue culture is faster, and higher multiplication rates are possible (Tsubomura and Taniguchi, 2008; Sahu and Sahu, 2013). Micropropagation starting from nodal segments uses smaller explants than conventional techniques, and selection of juvenile explants is possible (Sahu and Sahu, 2013). Regarding explants coming from mature trees, they are less likely to dedifferentiate and reprogram (Narváez et al., 2019), but the use of explants from mature trees for conifer micropropagation has also been reported (Rojas-Vargas et al., 2021; Juan-Vicedo et al., 2022). However, because these trees are selected after reaching maturity, there has been limited success reported in the vegetative propagation of mature conifers, and procedures should be improved. Although the benefits of tissue culture for the propagation of forest trees have been recognized, the success of such methods is still highly dependent on the species, the explant quality, the age of the donor plant, the culture medium, PGRs, and/or the interaction among all these different factors. As a result, morphogenesis determines the growth and development of plant tissues, and it is influenced by several physico-chemical factors (Rai et al., 2010). In relation to culture media, DCR (Gupta and Durzan, 1985), MS (Murashige and Skoog, 1962), and QL (Quoirin and Lepoivre, 1977) are commonly used basal media for *in vitro* regeneration of conifers (De Diego et al., 2010; Arab et al., 2014). Furthermore, the CKs added to the culture medium have a direct effect on the endogenous phytohormone balance, provoking a response to the induction of axillary shoot buds and affecting the organogenesis of the culture (Phillips, 2004; De Almeida et al., 2012).

Carbohydrates in plants are basic elements; they constitute substrates for respiration and are essential for many other processes related to plant development or gene expression, and in many species they favor rooting, acting mainly as a source of energy (Gibson, 2000; Calamar and De Klerk, 2002; Dewir et al., 2016).

Plant growth and development are also influenced by different physical factors, with light being one of the most important (Alallaq et al., 2020). The traditional light source used in *in vitro* culture in the growth chambers is FL, with irradiances between 25 and 150 $\mu\text{molm}^{-2}\text{s}^{-1}$ for a 16 h photoperiod (Larraburu et al., 2018). FL emit a broad light spectrum, and their physiological effects on plants are not specific (Rocha et al., 2010). Furthermore, the power consumption of FL is high as the heat emitted needs to be removed from growth chambers using air conditioners, making the process expensive (Kulus and Woźny, 2020). LEDs are available today as an alternative to conventional light sources for *in vitro* plant growth (Marín-Martínez et al., 2022), and they present advantages over FL such as small size, longer lifespan, less power consumption, high energy conversion efficiency, and adjustable light spectra (Río-Alvarez et al., 2014).

The analysis of the combined effects morphological (type of explants), chemical (culture media, sucrose concentrations), and physical factors (LEDs) at different micropropagation stages of adult trees has not been carried out in *Cryptomeria japonica*.

Moreover, the use of ventilated culture containers is not widely used for conifer species. For these reasons, with the main objective of improving the micropropagation protocol for adult Japanese cedar, we focused on optimizing (1) the shoot induction stage using different types of explants, CKs, and culture media as well as (2) the rooting stage using different sucrose concentrations, Ecobox containers[®], and light treatments (FL versus LEDs).

2. MATERIALS AND METHODS

2.1. Plant Material

Actively growing *C. japonica* twigs were collected in October 2019 from two healthy >30-year-old adult trees located in Arkaute (Spain; 42°51'9.35" N, 2°37'30.55" W) to carry out Experiment 1 (Figure 1A). In October 2020, actively growing twigs were collected from four healthy >30-year-old adult trees located in Urnieta (Spain; 43°13'13.03" N, 1°57'40.91" W), and one adult tree (Spain; 43°13'50.32" N, 1°59'25.37" W) was chosen to carry out Experiment 2.




Figure 1. Plant material at different stages of *Cryptomeria japonica* micropropagation process: **(A)** actively growing twigs used as starting explants; arrows show the explants used for *in vitro* culture establishment, bar=30 mm; **(B)** explant type; apical explants >1.5 cm (AT), apical explants <1.0 cm (AS), basal explants >1.5 cm (BT) and basal explants <1.0 cm (BS), bar=10 mm; **(C)** basal explants of 2.0 cm length after 4 weeks cultured in QL medium (Quoirin and Lepoivre, 1977) supplemented with 8.8 μM BA, bar=30 mm; **(D)** elongated shoots after 6 weeks in hormone-free QL medium supplemented with 2 gL^{-1} activated charcoal, bar=30 mm; **(E)** rooted shoots after six weeks in hormone-free QL medium supplemented with 2 gL^{-1} activated charcoal and under red LEDs, bar=30 mm; **(F)** plantlets after four weeks in *ex vitro* conditions in the greenhouse, bar=30 mm.

2.2. Sterilization

The plant material was first washed with commercial detergent, then rinsed under running water for 5 min immersed in 70% ethanol for 2 min, and then washed two times with sterile distilled water in the laminar flow unit. Finally, the actively growing twigs were disinfected in commercial bleach (30% v/v) (active chlorine 37 gL⁻¹ sodium hypochlorite) for 20 min and rinsed three times in sterile distilled water for 5 min each.

After the sterilization protocol, four types of explants were tested: apical explants >1.5 cm (AT), apical explants <1.0 cm (AS), basal explants >1.5 cm (BT), and basal explants <1.0 cm (BS) (Figure 1B).

2.3. Micropropagation Process

2.3.1. Experiment 1

After sterilization, the four types of explants were cultured vertically on 25×145 mm test tubes with polypropylene caps (Lab Associates, Oudenbosch, Netherlands) containing 15 mL of bud IM medium (Supplementary Table S1). Three basal media were assayed: DCR (Gupta and Durzan, 1985), MS (Murashige and Skoog, 1962), and QL (Quoirin and Lepoivre, 1977). All media were supplemented with 3% (w/v) sucrose, BA 8.8 μM (Duchefa Biochemie, Haarlem, Netherlands) and 8 gL⁻¹ Difco Agar® (Becton and Dickinson, Madrid, España) granulated. The pH of all media was adjusted to 5.8, and then they were autoclaved at 121 °C for 20 min. All cultures were placed in the growth chamber at a photoperiod of 16 h with 120 μmolm⁻²s⁻¹ light intensity provided by cool FL tubes (TLD 58 W/33; Philips, France) and a temperature of 21±1 °C.

As soon as SI was observed (after four weeks) (Figure 1C), four to five explants were transferred to baby food glass jars with Magenta™ b-cap lids filled with 25 mL of EM (Supplementary Table S1). EM was composed of hormone-free DCR, MS, or QL supplemented with 2 gL⁻¹ activated charcoal, 3% (w/v) sucrose and solidified with 8.5 gL⁻¹ Difco Agar® granulated; pH and autoclaving conditions were those mentioned for IM.

The shoots were transferred to fresh EM every six weeks. Shoots were cultivated individually in a fresh EM when they reached 10-15 mm (Figure 1D). The conditions in the growth chamber were the same as those described above.

2.3.2. Root Induction and Acclimatization of Rooted Plants

Elongated shoots of at least 20-25 mm long were used for RI. Based on the results of Experiment 1, QL basal medium was selected. The explants were transferred to Ecoboxes (Eco2box/green filter, consisting of a polypropylene oval vessel with a “breathing” hermetic cover (Duchefa®) with 100 mL of RIM (Supplementary Table S1), which was composed of half-strength macronutrient QL medium with 50 μM NAA (Duchefa Biochemie, Haarlem, Netherlands), 8 gL⁻¹ Difco Agar® and 1.5% or 3% (w/v) sucrose. The pH and autoclaving conditions were those previously described. The shoots were placed under dim light for eight days. Then, two different light treatments were tested for four weeks: (A) FL (color temperature 4000 K), 120 μmolm⁻²s⁻¹ light intensity provided by cool LF (TLD 58 W/33; Philips, France); and (B) red light (peak wavelength 630 nm), 60 μmolm⁻²s⁻¹ light intensity provided by adjustable LEDs (RB4K Grow Light LEDs). The photoperiod and the temperature of the growth chamber were the same as previously described.

After five weeks of culture in RIM, shoots were cultured for six weeks in Ecoboxes with 100 mL of REM (Supplementary Table S1); this medium consisted of half-strength macronutrient QL medium supplemented with 2 gL⁻¹ activated charcoal, 3% sucrose or 1.5% (w/v) sucrose and 8.5 gL⁻¹ Difco Agar®. Then, the rooted plants (Figure 1E) were planted *ex vitro*, transferring them to moist peat moss (Pindstrup, Aarhus, Denmark) with vermiculite at a proportion of 8:2 (v/v); acclimatization was carried out in a

greenhouse under controlled conditions at a temperature of $21\pm 1^{\circ}\text{C}$ and progressively decreasing the humidity during a month from 95 to 80% (Figure 1F).

2.3.3. Experiment 2

Based on the results of Experiment 1, basal explants of >1.5 cm length and QL medium were selected to perform this experiment (Supplementary Table S2). The medium was supplemented with one of these three types of CK: BA, m-T, or TDZ (Duchefa Biochemie, Haarlem, Netherlands), all of them at $8.8\ \mu\text{M}$. The explants were placed in the growth chamber at the same conditions described above (Section 2.3.1. Experiment 1).

As soon as SI was observed (after four weeks), four to five shoots were transferred into baby food glass jars with MagentaTM b-cap lids and 25 mL of EM (Supplementary Table S2), and they were subcultured every six weeks. Shoots that reached 10-15 mm were separated and individually cultivated in fresh EM. The growth chamber temperature and photoperiod were the same as those previously described (Section 2.3.1. Experiment 1).

2.3.4. Root Induction and Acclimatization of Rooted Plants

Shoots at least 20-25 mm long from the EM were employed for RI. Based on the results of Experiment 1, the shoots were cultivated in Ecoboxes filled with half-strength macronutrient QL basal medium supplemented with $50\ \mu\text{M}$ NAA, 1.5% (w/v) sucrose and $8\ \text{g}\cdot\text{L}^{-1}$ Difco Agar[®] (Supplementary Table S2). In addition, based on the results from Experiment 1, red LED was selected for the rooting stage; the shoots were placed under dim light for eight days, followed by four weeks under a 16 h photoperiod with red light (peak wavelength 630 nm) and $60\ \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ light intensity provided by adjustable LEDs (RB4K Grow Light LEDs). After these five weeks, shoots were cultured for six weeks in Ecoboxes with 100 mL of REM (Supplementary Table S2). The photoperiod and the temperature of the growth chamber were the same as those described in previous sections. Then, rooted plants were acclimatized as described above (Section 2.3.2).

2.4. Data collection and Statistical Analysis

2.4.1. Experiment 1

Twenty-four to forty-eight test tubes and one explant per test tube (AT, AS, BT, or BS) per each tree (two trees) were cultured in each culture medium. After two months of culture, the contamination, survival, and SI percentage (%) for each condition tested were measured. In the case of the SI (%) and the NS/E, these were calculated with respect to the non-contaminated explants. The effect of the explant type and culture medium on survival and SI was analyzed using a logistic regression model, and when necessary, differences were assessed by Tukey's post hoc test ($\alpha=0.05$).

Data for the NS/E were analyzed by ANOVA. When necessary, multiple comparisons were made using Tukey's post hoc test ($\alpha=0.05$).

After the root expression stage, data for the RI (%), the mean of NR/E and the LLR (cm) were recorded. A completely randomized design using seven to twenty-four plantlets per sucrose concentration and light treatment was performed.

The effect of the sucrose concentration and light treatment on RI (%) was analyzed with a logistic regression model. Data for NR/E and LLR were analyzed by ANOVA, and when necessary, differences were assessed by Tukey's post hoc test ($\alpha=0.05$). To evaluate the effect of the sucrose concentration and light treatment on the acclimatization percentage, a logistic regression

model was applied to plantlets after four weeks of growth in the greenhouse. Data processing was done using R Core Team software® (version 4.2.1, Vienna, Austria).

2.4.2. Experiment 2

Forty test tubes and one explant per test tube (BT) were cultured in each culture medium per tree (five trees). After two months of culture, the contamination, survival and SI percentages were recorded for each condition tested.

The SI (%) and the NS/E were calculated with respect to the non-contaminated explants after the elongation stage. The effect of the CK type on survival and SI (%) was analyzed using a logistic regression model; when necessary, differences were assessed by Tukey's post hoc test ($\alpha=0.05$).

Data for the NS/E were analyzed by ANOVA, and when necessary, differences were assessed by Tukey's post hoc test ($\alpha=0.05$). After the root expression stage, data for the RI (%), the mean of NR/E, and the LLR (cm) were recorded. A completely randomized design using forty-one to fifty plantlets per CK applied during the SI stage was performed.

The effect of the CK applied during the SI stage on the RI was analyzed using a logistic regression model. Data for NR/E and LLR were analyzed by ANOVA, and when necessary, differences were assessed by Tukey's post hoc test ($\alpha=0.05$). To evaluate the effect of the CK type applied during the SI stage on the acclimatization percentage, a logistic regression was used. As mentioned above, all data were processed using R Core Team software®.

3. RESULTS

3.1. Micropropagation Process

3.1.1. Experiment 1

The rates of contamination were registered eight weeks after sterilization, showing general values of 39%. Explants' survival was significantly affected by the explant type used; AS explants showed significantly higher survival rates (90%) than AT (62%) and BT (32%; Figure 2, Supplementary Table S3).




Figure 2. Survival (%) in different explant types of *Cryptomeria japonica* cultured in DCR medium (Gupta and Durzan, 1985), MS medium (Murashige and Skoog, 1962) and QL medium (Quoirin and Lepoivre, 1977). Apical explants >1.5 cm (AT), apical explants <1.0 cm (AS), basal explants >1.5 cm (BT), and basal explants <1.0 cm (BS). Data are presented as mean values \pm S.E. Significant differences are indicated by different letters according to Tukey's post hoc test ($p < 0.05$).

The basal medium and the interaction between explant type and basal medium did not show significant differences for survival (%) (Supplementary Table S3). The survival percentage ranged from 57% in explants grown in MS and QL media to 63% in explants cultured in DCR medium.

When the variables explant type and basal medium and the interaction between them were analyzed after the induction stage, statistically significant differences were only found for the percentage of SI depending on the explant type (Figure 3 and Supplementary Table S4). A significantly higher SI percentage was obtained in AS (69%) and AT (56%) compared to the rest of the explants tested (Figure 3).




Figure 3. Shoot induction (%) in different explant types of *Cryptomeria japonica* cultured in DCR medium (Gupta and Durzan, 1985), MS medium (Murashige and Skoog, 1962) and QL medium (Quoirin and Lepoivre, 1977). Apical explants >1.5 cm (AT), apical explants <1.0 cm (AS), basal explants >1.5 cm (BT), and basal explants <1.0 cm (BS). Data are presented as mean values \pm S.E. Significant differences are indicated by different letters according to Tukey's post hoc test ($p < 0.05$).

The SI percentage ranged from 38% in explants grown in DCR medium to 46% in explants cultured in QL medium. Explants developed in MS medium showed an intermediate value of SI (43%).

Regarding the NS/E, significant differences were found for the variables explant type and basal medium and the interaction between them (Figure 4, Supplementary Table S4). BT explants cultured in QL medium produced a significantly higher response than the other explant and medium combinations tested (Figure 4). Based on the results of Experiment 1, basal explants of >1.5 cm length and QL medium were selected to carry out the micropropagation process in Experiment 2.




Figure 4. Number of shoots per explant in different explant types of *Cryptomeria japonica* cultured in DCR medium (Gupta and Durzan, 1985), MS medium (Murashige and Skoog, 1962) and QL medium (Quoirin and Lepoivre, 1977) supplemented with 6-benzyladenine (BA, 8.8 μ M). Apical explants >1.5 cm (AT), apical explants <1.0 cm (AS), basal explants >1.5 cm (BT), and basal explants <1.0 cm (BS). Data are presented as mean values \pm S.E. Significant differences are indicated by different letters according to Tukey's post hoc test ($p < 0.05$).

3.1.2. Root Induction and Acclimatization of Rooted Plants

When the effect of the sucrose concentration, the light treatment, or the interaction between them on the RI (%) was analyzed, no statistically significant differences were observed (Supplementary Table S5). The RI percentage ranged from 36% in shoots cultured in 3% sucrose concentration under FL to 54% in shoots, independently of the sucrose concentration in the culture medium and exposure to red LEDs.

The root number was significantly affected by the light treatment applied in the rooting phase (Figure 5 and Supplementary Table S5). In this sense, explants exposed to red LEDs showed significantly higher NR/E (6.5 ± 0.5) than those under FL (2.7 ± 0.4) (Figure 5). The sucrose concentration and the interaction between sucrose concentration and light treatment did not show statistically significant differences (Supplementary Table S5). The NR/E ranged from 4.5 ± 0.6 for shoots grown with 1.5% sucrose to 5.4 ± 0.7 for shoots cultured in QL medium supplemented with 3% sucrose.




Figure 5. Number of roots per explant in shoots of *Cryptomeria japonica* cultured in QL medium (Quoirin and Lepoivre, 1977), supplemented with sucrose [1.5 or 3 (% w/v)] under light treatments [fluorescent light (F) and red LEDs (R)]. Data are presented as mean values ± S.E. Significant differences are indicated by different letters according to Tukey's post hoc test ($p < 0.05$).

The different sucrose concentrations (1.5% or 3.0%) and light treatments tested for SI showed a statistically significant effect on LLR, whereas the interaction between them did not have a significant effect (Figure 6 and Supplementary Table S5). A significantly higher LLR was observed in shoots cultured in the presence of 3% (w/v) sucrose compared with shoots grown at the lowest sucrose concentration (Figure 6A). In the same way, shoots exposed to FL showed significantly higher LLR than those under red LEDs (Figure 6B).




Figure 6. Length of the longest root in shoots of *Cryptomeria japonica* cultured in QL medium (Quoirin and Lepoivre, 1977), according to sucrose concentration [1.5 or 3 (% w/v)] (A) and light treatments [fluorescent light (F) and red LEDs(R)] (B). Data are presented as mean values ± S.E. Significant differences are indicated by different letters according to Tukey's post hoc test ($p < 0.05$).

When the effect of sucrose concentration and light treatment applied during the rooting phase on the acclimatization percentage of rooted shoots was analyzed, no statistically significant differences were observed (Supplementary Table S6). Statistically significant differences were found in the acclimatization percentage for the interaction between sucrose concentration and light treatment (Supplementary Table S6). Nevertheless, as the p-value was bordering on significance, Tukey's post hoc test could not

detect them. The acclimatization percentage ranged from 30% in shoots cultured in medium supplemented with 3% (w/v) and exposed to FL to 80% in those growing in the same sucrose concentration supplemented the culture medium under red LEDs.

3.2. Experiment 2

Four weeks after sterilization, the contamination rates were at 28%. No statistically significant differences were observed for the explant survival percentage considering the CK tested (Supplementary Table S7). Explant survival percentages ranged from 66% in explants cultured in medium with m-T to 76% in explants grown with BA treatment.

The CK type showed a significant effect on the SI (%) (Figure 7, Supplementary Table S8). A significantly higher SI percentage was observed in explants induced with BA and m-T treatments (50% and 48%, respectively) than in explants grown on QL medium supplemented with TDZ.




Figure 7. Shoot induction (%) in explants of *Cryptomeria japonica* cultured on QL medium (Quoirin and Lepoivre, 1977), supplemented with 6-benzyladenine (BA), meta-topolin (m-T), and thidiazuron (TDZ) (8.8 μ M). Data are presented as mean values \pm S.E. Significant differences are indicated by different letters according to Tukey's post hoc test ($p < 0.05$).

No statistically significant differences were observed when the effect of CK type on the NS/E was analyzed (Supplementary Table S8). It was not possible to obtain shoots from explants induced with TDZ treatment due to tissue necrosis. Explants induced with m-T and BA treatments produced 2.3 ± 0.1 and 2.4 ± 0.1 NS/E, respectively.

Regarding the RI percentage, significant differences were observed depending on the CK used for SI (Figure 8 and Supplementary Table S9). A significantly higher RI percentage was recorded in shoots induced with m-T (54%) than in shoots grown with BA (33%) during the SI stage (Figure 8).




Figure 8. Root induction (%) in shoots of *Cryptomeria japonica* cultured in QL medium (Quoirin and Lepoivre, 1977), supplemented with 50 μ M 1-naphthaleneacetic acid (NAA), according to cytokinin type (6-benzyladenine (BA) and meta-topolin (m-T) (8.8 μ M) use during shoot induction stage. Data are presented as mean values \pm S.E. Significant differences are indicated by different letters according to Tukey's post hoc test ($p < 0.05$).

When the effect of the CK type utilized for SI was evaluated for the NR/E and LLR parameters, no statistically significant differences were observed (Supplementary Table S9). The NR/E was 3.1 ± 0.5 for shoots previously induced with BA and 3.9 ± 0.6 for those from m-T treatment.

Explants cultured with BA and m-T treatment during the induction stage showed a LLR of 3.3 ± 0.5 cm and 3.7 ± 0.3 cm, respectively. No statistically significant differences were observed for shoots coming from different CK treatments when acclimatization percentage was analyzed (Supplementary Table S10). The *ex vitro* survival rate was 92% for shoots induced with BA and 94% for those induced with m-T treatment.

4. DISCUSSION

As was reviewed by Payghamzadeh and Kazemitabar (2011), contamination is considered a crucial obstacle that prohibits the successful establishment of an aseptic *in vitro* culture. In this study, contamination rates ranging from 28% to 39% were obtained using sodium hypochlorite. In this sense, our work obtained lower contamination rates than those recorded by (Hine-Gómez and Valverde, 2003) for *C. japonica*, who obtained 46% using calcium hypochlorite for surface sterilization of apical explants from adult trees. The sterilization protocol applied in Experiment 1 and Experiment 2 resulted in an optimal *in vitro* establishment of *C. japonica*. Therefore, our results suggest that it can be used for the establishment of cultures from other mature conifer explants (De Diego et al., 2010).

Factors such as the type, size, or age of the explant, the physiological state of the donor plant, the type of disinfectant and its concentration, can influence the effectiveness of the sterilization protocol (Da Silva et al., 2016). In our case, the survival percentage was significantly higher in AS explants in Experiment 1. A similar tendency was observed in *Cedrus libani*, where shoot apices (<1.0 cm) from adult trees showed the best survival response (Renau-Morata et al., 2005). Similarly, in *Taxus mairei*, the highest survival percentage was obtained in small stems (<1.0 cm) from cutting (Chang et al., 2001). In our study, the smaller size and lower exposure of the explants' surface could have favored the culture establishment (Sathyagowri and Seran, 2011), as supported by George and Debergh (2008), who mentioned that smaller explants may be easier plant material to sterilize from microorganisms.

In our experiments, the efficiency of the organogenic response was assessed by considering the SI percentage and the NS/E. In Experiment 1, the results showed statistically significant differences for AS and AT explants for SI percentage. In this sense, our study confirmed that the explant size is an important factor affecting axillary bud proliferation (Renau-Morata et al., 2005), and probably, the larger explants have more mineral nutrient reserves and endogenous hormones to support the culture (Desai et al., 2015). Additionally, George and Debergh (2008) explained that bigger explants from extensive parts of the shoot apex or stem segments with lateral buds could have advantages over smaller explants. Furthermore, it is probable that the morphogenesis process of the explants tested that is under the influence of exogenous PGRs and its interactions with endogenous phytohormones (De Almeida et al., 2012).

CKs play several recognized roles in plant development, through the suppression of apical dominance and promotion of the development of axillary buds, the promotion of cell division, or the stimulation of plant protein synthesis (Arab et al., 2014; Rodríguez et al., 2022). BA is the most used plant growth regulator; it is applied alone or in combination with other CK to promote *in vitro* SI due to its effectiveness and affordability (Bairu et al., 2007; De Diego et al., 2008; Phillips and Garda 2019). During Experiment 2, explants cultured with BA or m-T showed an efficient organogenic response regarding SI percentage, and although not significant, a slightly higher response in BT explants induced with BA was observed. Analogous results were obtained in several organogenesis protocols from adult trees of *P. pinea*, *P. radiata*, *S. sempervirens* and *P. halepensis*, where, to obtain an *in vitro* shoot response, BA at concentrations ranging from 4.4 to 50 μM was utilized (Cortizo et al., 2009; Montalbán et al., 2013; Pereira et al., 2021, Rojas-Vargas et al., 2021; Rojas-Vargas et al., 2022).

The highest NS/E was obtained in BT explants cultured in QL medium (Experiment 1). In accordance with this result, in previous experiments in our lab, the highest number of NS/E was obtained in *S. sempervirens* when explants bigger than 1.5 cm in length were used (Rojas-Vargas et al., 2021). Similarly, Hine-Gómez and Valverde (2003) and Rafi and Salehi (2018) developed an *in vitro* protocol for *C. japonica* and *Cedrus deodara*, using explants of mature trees from 1.0 to 2.5 cm in length, respectively. On the other hand, NS/E was higher on basal QL medium than on basal MS and DCR medium. According to Maruyama et al. (2021), this may be attributable to a higher level of inorganic nitrogen present in those media when compared to QL medium. This hypothesis is supported by Tuskan et al. (1990) who explained that a nitrate excess could have a negative impact on the organogenic response. Similar to our experiment on *C. japonica*, the lower nitrogen content of QL medium promoted organogenesis in *C. deodara* and *Pinus ponderosa* (Rafi and Salehi, 2018; Rojas-Vargas et al., 2023). Recently, for improving the micropropagation protocol of plants, artificial neural networks algorithms have been used to build models to determine the effect of mineral nutrients, vitamins, and PGRs on several growth and quality parameters of micropropagated plants (Arteta et al., 2022).

Several studies attributed the improvement in multiplication rates or rooting percentages and the alleviation of physiological disorders to the use of topolins in plant tissue culture (Aremu et al., 2012). However, in our study, no significant differences for the effect of CK on NS/E were found, but a slightly higher response in explants cultured with BA was observed (Experiment 2). Analogous results were obtained in *P. radiata*, *S. sempervirens*, and *P. ponderosa*, where the induction of axillary buds was not improved when applying m-T instead of BA (Montalbán et al., 2011; Montalbán et al., 2013; Rojas-Vargas et al., 2021; Rojas-Vargas et al., 2023).

TDZ has received more attention due to its ability to aid *in vitro* regeneration of woody plants (Cortizo et al., 2009; De Diego et al., 2010; Zarei et al., 2020), and it has been recommended in explants coming from adult trees to induce regeneration via axillary shoot proliferation and adventitious shoot organogenesis (Vinoth and Ravindhran, 2018). However, in our work, it was not possible to obtain shoots coming from explants induced with TDZ due to tissue necrosis. The reason for this result may be due to

an inadequacies concentration of TDZ, as supported by Dewir et al. (2018), who mentioned that the TDZ concentration and exposure time at their optimum depends on the species. Nevertheless, TDZ at low concentrations, in pulse treatments or during short exposure periods can be effective in circumventing TDZ induced abnormalities such as tissue necrosis (Dewir et al., 2018). Adventitious rooting can be influenced by physical and chemical factors: PGRs, light quality, temperature, medium composition and carbohydrates (Zarei et al., 2020). When RI percentage was analyzed, no significant effect of sucrose concentration (1.5 or 3.0%) and light treatment was found. A similar result was observed in the organogenesis protocols of *C. japonica*, where the culture medium was supplemented with sucrose 3.0% to obtain rooting (Hine-Gómez and Valverde, 2003; Koguta et al., 2017). As mentioned above, in the last few years, m-T has been proven as an alternative to conventional CK for *in vitro* propagation of plants (Gantait and Mitra, 2021). In this sense, the advantages of m-T to promote rhizogenesis have been described (Aremu et al., 2012). In Experiment 2, when we focused on the study of the effect of the CK type on RI, we found that shoots induced with m-T promoted the highest rooting percentage. Similar to our results, Naaz et al. (2019), studying *Syzygium cumini*, reported that shoots induced with m-T increased rhizogenic competency compared with shoots coming from KIN, 2-isopentyl adenine or BA. Similarly, in *Caralluma umbellata*, shoots derived from a culture medium supplemented with m-T and NAA showed the highest *in vitro* rooting activity (Jayaprakash et al., 2023). In contrast to our work, in *S. sempervirens* and *Juniperus drupacea*, growing shoots with m-T did not improve the rooting response (Rojas-Vargas et al., 2021; Ioannidis et al., 2023). Summarizing, the effect of m-T to promote rhizogenesis at different concentrations may be species-specific and depends on the starting material used to establish the *in vitro* culture.

Plant growth and development are strongly influenced by the quality of the light in their environment (Sarropoulou et al., 2023). In the last few years, LEDs have shown a favorable response in *in vitro* culture when compared with the results obtained using fluorescent tubes (Chen et al., 2020; Gupta and Jatothu, 2013). Additionally, Ragonezi et al., (2010) mentioned that the light type and wavelength specificity influence adventitious rooting. In this sense, statistically significant differences in NR/E were found when shoots were cultured under red LEDs (Experiment 1). Similarly, in *C. japonica* and *Populus sieboldii* × *Populus grandidentata*, red LEDs showed a higher response to *in vitro* rooting (Tsubomura and Tanaguchi, 2008; Kobayashi et al., 2022). In *Pinus pseudostrobus* shoots exposed to red LEDs, the best rooting rates were observed at 30 days of evaluation (Marín-Martínez et al., 2022). Contrary to our results, *P. radiata* and *P. ponderosa* shoots grown under white FL and white LEDs, respectively, displayed the highest rooting responses (Rojas-Vargas et al., 2022; Rojas-Vargas et al., 2023). Regarding LLR, shoots under FL showed longer primary roots. Analogous results were observed by Ishii et al. (2011) in *C. japonica* and Rojas-Vargas et al. (2023) in *P. ponderosa*, where shoots growing under FL showed the longest root length. Contrary to our result, the longest roots were recorded in plants of *P. radiata* and *P. pseudostrobus* exposed to red LEDs (Marín-Martínez et al., 2022; Rojas-Vargas et al., 2022). Summarizing, different plant species respond differently, even when using the same light treatment, which may be due to the genotypic characteristics of the plants or the physiological state of the explants (Da Silva et al., 2016; Lai et al., 2022). Regarding the acclimatization stage, no significant differences were observed for the effect of CK on acclimatization percentage, but a slightly higher response in explants cultured with m-T was observed. These results agreed with those observed in *Aloe polyphylla*, *S. cumini* and *C. umbellata*, where the plantlets induced with m-T were successfully acclimatized (Bairu et al., 2007; Naz et al., 2019; Jayaprakash et al., 2023). The reason for this may be due to the rapid uptake and transport of m-T into the plant system and the production of reversibly sequestered metabolites (Jayaprakash et al., 2023).

5. CONCLUSIONS

The regeneration of *C. japonica* through micropropagation of adult trees using basal explants of >1.5 cm length was achieved and it depends on physico-chemical factors. The optimal result in terms of shoot induction was obtained when basal explants were cultured in QL medium supplemented with BA treatment.

Our results suggest that the use of m-T and a 1.5% sucrose concentration favored root induction. In this sense, the use of red LEDs was better for the number of roots per explant. Finally, in the greenhouse, the shoots, independently of the CKs used in the shooting stage, showed high acclimatization success. To optimize the micropropagation efficiency in *C. japonica*, our results suggest the use of fluorescent light for the shoot induction stage and red LEDs for the rooting stage.

6. REFERENCES

- Alallaq, S.; Ranjan, A.; Brunoni, F.; Novák, O.; Lakehal, A.; Bellini, C. Red light controls adventitious root regeneration by modulating hormone homeostasis in *Picea abies* seedlings. *Front. Plant Sci.* **2020**, *11*, 586140. <https://doi.org/10.3389/fpls.2020.586140>.
- Arab, M.M.; Yadollahi, A.; Shojaeiyan, A.; Shokri, S.; Ghoghah, S.M. Effects of nutrient media, different cytokinin types and their concentrations on in vitro multiplication of G N15 (hybrid of almond peach) vegetative rootstock. *J. Genet. Eng. Biotechnol.* **2014**, *12*, 81–87. <https://doi.org/10.1016/j.jgeb.2014.10.001>.
- Aremu, A.O.; Bairu, M.W.; Doležal, K.; Finnie, J.F.; Van Staden, J. Topolins: A panacea to plant tissue culture challenges? *Plant Cell Tissue Organ Cult.* **2012**, *108*, 1–16. <https://doi.org/10.1007/s11240-011-0007-7>.
- Arteta, T.A.; Hameg, R.; Landin, M.; Gallego, P.P.; Barreal, M.E. Artificial Neural Networks Elucidated the Essential Role of Mineral Nutrients versus Vitamins and Plant Growth Regulators in Achieving Healthy Micropropagated Plants. *Plants* **2022**, *11*, 1284. <https://doi.org/10.3390/plants11101284>.
- Bairu, M.W.; Stirk, W.A.; Dolezal, K.; Van Staden, J. Optimizing the micropropagation protocol for the endangered *Aloe polyphylla*: Can meta-topolin and its derivatives serve as replacement for benzyladenine and zeatin? *Plant Cell Tissue Organ Cult.* **2007**, *90*, 15–23. <https://doi.org/10.1007/s11240-007-9233-4>.
- Calamar, A.; De Klerk, G.J. Effect of sucrose on adventitious root regeneration in apple. *Plant Cell Tissue Organ Cult.* **2002**, *70*, 207–212. <https://doi.org/10.1023/A:1016356123367>.
- Chang, S.H.; Ho, C.K.; Chen, Z.Z.; Tsay, J.Y. Micropropagation of *Taxus mairei* from mature trees. *Plant Cell Rep.* **2001**, *20*, 496–502. <https://doi.org/10.1007/s002990100362>.
- Chen, L.; Zhang, K.; Gong, X.; Wang, H.; Gao, Y.; Wang, X.; Zeng, Z.; Hu, Y. Effects of different LEDs light spectrum on the growth, leaf anatomy, and chloroplast ultrastructure of potato plantlets in vitro and minituber production after transplanting in the greenhouse. *J. Integr. Agric.* **2020**, *19*, 108–119. [https://doi.org/10.1016/S2095-3119\(19\)62633-X](https://doi.org/10.1016/S2095-3119(19)62633-X).
- Cortizo, M.; De Diego, N.; Moncaleán, P.; Ordás, R.J. Micropropagation of adult stone pine (*Pinus pinea* L.). *Trees Struct. Funct.* **2009**, *23*, 835–842. <https://doi.org/10.1007/s00468-009-0325-0>.
- Da Silva, J. A. T.; Winarto, B.; Dobránszki, J.; Cardoso, J. C.; Zeng, S. Tissue disinfection for preparation of *Dendrobium* in vitro culture. *Folia Hortic.* **2016**, *28*, 57–75. <https://doi.org/10.1515/fhort-2016-0008>.

- De Almeida, M.; de Almeida, C.V.; Graner, E.M.; Brondani, G.E.; de Abreu-Tarazi, M.F. Pre-procambial cells are niches for pluripotent and totipotent stem-like cells for organogenesis and somatic embryogenesis in the peach palm: A histological study. *Plant Cell Rep.* **2012**, *31*, 1495–1515. <https://doi.org/10.1007/s00299-012-1264-6>.
- De Diego, N.; Montalbán, I.A.; Fernandez de Larrinoa, E.; Moncaleán, P. In vitro regeneration of *Pinus pinaster* adult trees. *Can. J. For. Res.* **2008**, *38*, 2607–2615. <https://doi.org/10.1139/X08-102>.
- De Diego, N.; Montalbán, I.A.; Moncaleán, P. In vitro regeneration of adult *Pinus sylvestris* L. trees. *S. Afr. J. Bot.* **2010**, *76*, 158–162. <https://doi.org/10.1016/j.sajb.2009.09.007>.
- Desai, C.; Inghalihalli, R.; Krishnamurthy, R. Micropropagation of *Anthurium andraeanum* an important tool in floriculture. *J. Pharmacogn. Phytochem.* **2015**, *4*, 112–117.
- Dewir, Y.H.; Murthy, H.N.; Ammar, M.H.; Alghamdi, S.S.; Al-Suhaibani, N.A.; Alsadon, A.A.; Paek, K.Y. In vitro rooting of leguminous plants: Difficulties, alternatives, and strategies for improvement. *Hortic. Environ. Biotechnol.* **2016**, *57*, 311–322. <https://doi.org/10.1007/s13580-016-0060-6>.
- Dewir, Y.H.; Nurmansyah; Naidoo, Y.; da Silva, J.A.T. Thidiazuron-induced abnormalities in plant tissue cultures. *Plant Cell Rep.* **2018**, *37*, 1451–1470. <https://doi.org/10.1007/s00299-018-2326-1>.
- Forestry Agency. *Statistical Handbook of Forest and Forestry*; Forestry Agency, Ministry of Agriculture, Forestry and Fisheries: Tokyo, Japan, 2020; pp. 8–9. (In Japanese).
- Gantait, S.; Mitra, M. Role of Meta-topolin on in vitro shoot regeneration: An insight. In: *Me-Tatoplin: A Growth Regulator for plant Biotechnology and Agriculture*; Ahmad, N., Strnad, M., Eds.; Springer: Singapore, 2021; pp 143–168. <https://doi.org/10.1007>.
- George, E.F.; Debergh, P.C. Micropropagation: Uses and Methods. In *Plant Propagation by Tissue Culture*; George, E.F., Hall, M.A., De Klerk, G.J., Eds.; Springer: Dordrecht, The Netherlands, 2008; pp. 29–64. https://doi.org/doi.org/10.1007/978-1-4020-5005-3_1.
- Gibson, S.I. Plant sugar-response pathways. part of a complex regulatory web. *Plant Physiol.* **2000**, *124*, 1532–1539. <https://doi.org/10.1104/pp.124.4.1532>.
- Gupta, P.K.; Durzan, D.J. Shoot multiplication from mature trees of Douglas-fir (*Pseudotsuga menziesii*) and sugar pine (*Pinus lambertiana*). *Plant Cell Rep.* **1985**, *4*, 177–179. <https://doi.org/10.1007/BF00269282>.
- Gupta, S.; Jatothu, B. Fundamentals and applications of light-emitting diodes (LEDs) in in vitro plant growth and morphogenesis. *Plant Biotechnol. Rep.* **2013**, *7*, 211–220. <https://doi.org/10.1007/s11816-013-0277-0>.
- Hine-Gómez, A.; Valverde-Cerdas, L. Establecimiento in vitro de *Cryptomeria japonica*(Taxocidaceae). *Rev. Biol. Trop.* **2003**, *51*, 683–690. (In Spanish with English Abstract).
- Ioannidis, K.; Tomprou, I.; Panayiotopoulou, D.; Boutsios, S.; Daskalaku, E.N. Potential and Constraints on In Vitro Micropropagation of *Juniperus drupacea* Labill. *Forests* **2023**, *14*, 142. <https://doi.org/10.3390/f14010142>.

Ishii, K.; Hosoi, Y.; Taniguchi, T.; Tsubomura, M.; Kondo, T.; Yamada, H.; Saito, M.; Suda, T.; Fukisawa, T.; Tanaka, K. In vitro culture of various genotypes of male sterile Japanese cedar (*Cryptomeria japonica* D. Don). *Plant Biotechnol.* **2011**, *28*, 103–106. <https://doi.org/10.5511/plantbiotechnology.10.0906a>.

Jayaprakash, K.; Manokari, M.; Cokulraj, M.; Dey, Abhijit.; Faisal, M.; Alatar, A.A.; Joshee, N.; Shekhawat, M.S. Improved organogenesis and micro-structural traits in micropropagated plantlets of *Caralluma umbellata* Haw. in response to Meta-Topolin. *Plant Cell Tissue Organ Cult.* **2023**. <https://doi.org/10.1007/s11240-023-02447-0>.

Juan-Vicedo, J.; Serrano-Martinez, F.; Cano-Castillo, M.; Casas, J.L. In vitro propagation, genetic assessment, and, medium-term conservation of coastal endangered species *Tetraclinis articulata* (Vahl) Masters (Cupressaceae) from adult trees. *Plants* **2022**, *11*, 187. <https://doi.org/10.3390/plants11020187>.

Kobayashi, H.; Hashimoto, K.; Ohba, E.; Kurata, Y. Light-emitting Diode–Induced Root Photomorphogenesis and Root Retention in *Populus*. *HortScience* **2022**, *57*, 872–876. <https://doi.org/10.21273/HORTSCI16537-22>.

Koguta, K.V.; Flores, P.C.; Alcantara, G.B.; Higa, A.R. Serial micropropagation and cuttings as rejuvenation methods for *Cryptomeria japonica* (L.F) D.Don. *Rev. Espac.* **2017**, *38*, 7.

Kulus, D.; Woźny, A. Influence of light conditions on the morphogenetic and biochemical response of selected ornamental plant species under in vitro conditions: A mini-review. *BioTechnologia* **2020**, *101*, 75–83. <https://doi.org/10.5114/bta.2020.92930>.

Kusumi, J.; Tsumura, Y.; Yoshimaru, H.; Tachida, H. Phylogenetic relationships in Taxodiaceae and Cupressaceae sensu stricto based on matK gene, chlL gene, trnLtrnF IGS region, and trnL intron sequences. *Am. J. Bot.* **2000**, *87*, 1480–1488. <https://doi.org/10.2307/2656874>.

Lai, C.S.; Kho, Y.H.; Chew, B.L.; Raja, P.B.; Subramaniam, S. Organogenesis of *Cucumis metuliferus* plantlets under the effects of LEDs and silver nanoparticles. *S. Afr. J. Bot.* **2022**, *148*, 78–87. <https://doi.org/10.1016/j.sajb.2022.04.020>.

Larraburu, E.E.; Correa, G.S.; Llorente, B.E. In vitro development of yellow lapacho (Bignoniaceae) using high-power lightemitting diode. *Rev. Arvore.* **2018**, *42*, e420508. <https://doi.org/10.1590/1806-90882018000500008>.

Marín-Martínez, L.A.; Iglesias-Andreu, L.G. Effect of LED lights on the in vitro growth of *Pinus pseudostrobus* Lindl., plants. *J. For. Sci.* **2022**, *68*, 311–317. <https://doi.org/10.17221/43/2022-JFS>.

Maruyama, T.E.; Ueno, S.; Mori, H.; Kaneeda, T.; Moriguchi, Y. Factors influencing somatic embryo maturation in sugi (Japanese Cedar, *Cryptomeria japonica* (Thunb. ex L.f.) D. Don). *Plants* **2021**, *10*, 874. <https://doi.org/10.3390/plants10050874>.

Montalbán, I.A.; De Diego, N.; Moncaleán, P. Testing novel cytokinins for improved in vitro adventitious shoots formation and subsequent ex vitro performance in *Pinus radiata*. *Forestry* **2011**, *84*, 363–373. <https://doi.org/10.1093/forestry/cpr022>.

Montalbán, I.A.; Novák, O.; Rolčík, J.; Strnad, M.; Moncaleán, P. Endogenous cytokinin and auxin profiles during in vitro organogenesis from vegetative buds of *Pinus radiata* Adult Trees. *Physiol. Plant.* **2013**, *148*, 214–231. <https://doi.org/10.1111/j.1399-3054.2012.01709.x>.

- Murashige, T.; Skoog, F. A revised medium for rapid growth and bio assays with *Tobacco* tissue cultures. *Physiol. Plant.* **1962**, *15*, 474–497. <https://doi.org/10.1111/j.1399-3054.1962.tb08052.x>.
- Naaz, A.; Hussain, S.A.; Anis, M.; Alatar, A.A. Meta-topolin improved micropropagation in *Syzygium cumini* and acclimatization to ex vitro conditions. *Biol. Plant.* **2019**, *63*, 174–182. <https://doi.org/10.32615/bp.2019.020>.
- Narváez, I.; Martín, C.; Jiménez-Díaz, R.M.; Mercado, J.A.; Pliego-Alfaro, F. Plant regeneration via somatic embryogenesis in mature wild olive genotypes resistant to the defoliating pathotype of *Verticillium dahliae*. *Front. Plant Sci.* **2019**, *10*, 1471. <https://doi.org/10.3389/fpls.2019.01471>.
- Onuma, Y.; Uchiyama, K.; Kimura, M.; Tsumura, Y. Genetic analysis distinguished new natural population and old plantations of *Cryptomeria japonica*. *Trees For. People* **2023**, *11*, 100365. <https://doi.org/10.1016/j.tfp.2022.100365>.
- Park, J.S.; Barret, J.D.; Bonga, J.M. Application of somatic embryogenesis in high-value clonal forestry: Deployment, genetic control, and stability of cryopreserves clones. *Cell. Dev. Biol. Plant.* **1998**, *34*, 231–239. <https://doi.org/10.1007/BF02822713>.
- Payghamzadeh, K.; Kazemitabar, S.K. In vitro propagation of walnut—A review. *Afr. J. Biotechnol.* **2011**, *10*, 290–311. <https://doi.org/10.5897/AJB10.324>.
- Pereira, C.; Montalbán, I.A.; Pedrosa, A.; Tavares, J.; Pestryakov, A.; Bogdanchikova, N.; Canhoto, J.; Moncaleán, P. Regeneration of *Pinus halepensis* (Mill.) through organogenesis from apical shoot buds. *Forests* **2021**, *12*, 363. <https://doi.org/10.3390/f12030363>.
- Phillips, G.C. In vitro morphogenesis in plants—recent advances. *In Vitro Cell. Dev. Biol. Plant.* **2004**, *40*, 342–345. <https://doi.org/10.1079/IVP2004555>.
- Phillips, G.C.; Garda, M. Plant tissue culture media and practices: An overview. *In Vitro Cell. Dev. Biol. Plant.* **2019**, *55*, 242–257. <https://doi.org/10.1007/s11627-019-09983-5>.
- Quoirin, M.; Lepoivre, P. Études des milieux adaptés aux cultures in vitro de *Prunus*. *Acta Hort.* **1977**, *78*, 437–442. <https://doi.org/10.17660/ActaHortic.1977.78.54>.
- Rafi, Z.N.; Salehi, H. Factors affecting in-vitro propagation of some genotypes of Himalayan cedar [*Cedrus deodara* (Roxb. ex Lamb) G. Don.]. *Adv Hort. Sci.* **2018**, *32*, 479–48. <https://doi.org/10.13128/ahs-21833>.
- Ragonezi, C.; Klimaszewska, K.; Castro, M.R.; Lima, M.; de Oliveira, P.; Zavattieri, M.A. Adventitious rooting of conifers: Influence of physical and chemical factors. *Trees Struct. Funct.* **2010**, *24*, 975–992. <https://doi.org/10.1007/s00468-010-0488-8>.
- Rai, M.K.; Asthana, P.; Jaiswal, V.S.; Jaiswa, U. Biotechnological advances in guava (*Psidium guajava* L.): Recent developments and prospects for further research. *Trees* **2010**, *24*, 1–12. <https://doi.org/10.1007/s00468-009-0384-2>.
- Renau-Morata, B.; Ollero, J.; Arrillaga, I.; Segura, J. Factors influencing axillary shoot proliferation and adventitious budding in cedar. *Tree Physiol.* **2005**, *25*, 477–486. <https://doi.org/10.1093/treephys/25.4.477>.

Río-Álvarez, I.; Rodríguez-Herva, J.J.; Martínez, P.M.; González-Melendi, P.; García-Casado, G.; Rodríguez-Palenzuela, P.; López-Solanilla, E. Light regulates motility, attachment, and virulence in the plant pathogen *Pseudomonas syringae* pv tomato DC3000. *Environ. Microbiol.* **2014**, *16*, 2072–2085. <https://doi.org/10.1111/1462-2920.12240>.

Rocha, P.S.G.; Oliveira, R.P.; Scivittaro, W.B.; Santos, U.L. Diodos emissores de luz e concentrações de BAP na multiplicação in vitro de morangueiro. *Ciênc. Rural* **2010**, *40*, 1922–1928. <https://doi.org/10.1590/S0103-84782010000900011>.

Rodríguez, S.M.; Ordás, R.J.; Alvarez, J.M. Conifer Biotechnology: An Overview. *Forests* **2022**, *13*, 1061. <https://doi.org/10.3390/f13071061>.

Rojas-Vargas, A.; Castander-Olarieta, A.; Moncaleán, P.; Montalbán, I.A. Optimization of the micropropagation of elite adult trees of *Sequoia sempervirens*: Forest species of interest in the Basque Country, Spain. *Rev. Bionatura.* **2021**, *6*, 1511–1519. <https://doi.org/10.21931/RB/2021.01.01.11>.

Rojas-Vargas, A.; Castander-Olarieta, A.; Montalbán, I.A.; Moncaleán, P. Influence of physico-chemical factors on the efficiency and metabolite profile of adult *Pinus radiata* D. Don bud organogenesis. *Forests* **2022**, *13*, 1455. <https://doi.org/10.3390/f13091455>.

Rojas-Vargas, A.; Castander-Olarieta, A.; do Nascimento, A.M.M.; Vélez, M.L.; Pereira, C.; Martins, J.; Zuzarte, M.; Canhoto, J.; Montalbán, I.A.; Moncaleán, P. Testing Explant Sources, Culture Media, and Light Conditions for the Improvement of Organogenesis in *Pinus ponderosa* (P. Lawson and C. Lawson). *Plants* **2023**, *12*, 850. <https://doi.org/10.3390/plants12040850>.

Sahu, J.; Sahu, R.K. A review on low cost methods for in vitro micropropagation of plant through tissue culture technique. *UK J. Pharm. Biosci.* **2013**, *1*, 38–41. <https://doi.org/10.20510/ukjpb/1/i1/91115>.

Sarropoulou, V.; Sperdoulis, I.; Adamakis, I.D.; Grigoriadou, K. The use of different LEDs wavelength and light intensities for in vitro proliferation of cherry rootstock: Influence on photosynthesis and photomorphogenesis. *Plant Cell Tissue Organ Cult.* **2023**, *152*, 317–330. <https://doi.org/10.1007/s11240-022-02408-z>.

Sathyagowri, S.; Seran, T.H. In vitro plant regeneration of ginger (*Zingiber officinale* Rosc.) with emphasis on initial culture establishment. *J. Med. Aromat. Plants* **2011**, *1*, 195–202.

Taniguchi, T.; Konagaya, K.; Nanasato, Y. Somatic embryogenesis in artificially pollinated seed families of 2nd generation plus trees and cryopreservation of embryogenic tissue in *Cryptomeria japonica* D. Don (sugi). *Plant Biotechnol.* **2020**, *37*, 239–245. <https://doi.org/10.5511/plantbiotechnology.20.0220a>.

Tsubomura, M.; Taniguchi, T. Micropropagation of the male sterile ‘Soushun’ Japanese cedar. *Comb. Proc. Int. Plant. Prop. Soc.* **2008**, *58*, 421–427.

Tuskan, G.A.; Sargent, W.A.; Rensema, T.; Walla, J.A. Influence of plant growth regulators, basal media and carbohydrate levels on the in vitro development of *Pinus ponderosa* (Dougl. Ex Law.) cotyledon explants. *Plant Cell Tissue Organ Cult.* **1990**, *20*, 47–52. <https://doi.org/10.1007/BF00034756>.

Chapter 4

Vinoth, A.; Ravindhran, R. In vitro morphogenesis of woody plants using thidiazuron. In *Thidiazuron: From Urea Derivative to Plant Growth Regulator*; Ahmad, N., Faisal, M., Eds.; Springer: Singapore, 2018; pp. 211–230. <https://doi.org/10.1007/978-981-10-8004-3>.

Zarei, M.; Salehi, H.; Jowkar, A. Controlling the barriers of cloning mature *Picea abies* (L.) H. Karst. via tissue culture and co-cultivation with *Agrobacterium rhizogenes*. *Trees* **2020**, *34*, 637–647. <https://doi.org/10.1007/s00468-019-01945-z>

CHAPTER 5

Testing Explant Sources, Culture Media, and Light Conditions for the Improvement of Organogenesis in *Pinus ponderosa* P.

Lawson and C. Lawson

The content of this chapter corresponds to the published article “Rojas-Vargas, A.; Castander-Olarieta, A.; do Nascimento, A.M.M.; Vélez, M.L.; Pereira, C.; Martins, J.; Zuzarte, M.; Canhoto, J.; Montalbán, I.A.; Moncaleán, P. 2023. Testing Explant Sources, Culture Media, and Light Conditions for the Improvement of Organogenesis in *Pinus ponderosa* P. Lawson and C. Lawson. *Plants* 12, 850. <https://doi.org/10.3390/plants12040850>”.

ABSTRACT

Pinus ponderosa P. Lawson and C. Lawson is a commercial tree and one of the most important forest species in North America. Ponderosa pine suffers hardship when going through vegetative propagation and, in some cases, 15-30 years are needed to achieve full reproductive capacity. Based on previous works on *P. ponderosa* regeneration through *in vitro* organogenesis and trying to improve the published protocols, our objective was to analyze the influence of different types of explants, basal culture media, CKs, auxins, and light treatments on the success of shoot multiplication and rooting phases. Whole zygotic embryos and 44 μM BA showed the best results in terms of explants survival. For shoot organogenesis, whole zygotic embryos and half LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] macronutrients were selected. A significant positive interaction between whole zygotic embryos and half LP macronutrients was found for the percentage of EFS. Regarding the light treatments applied, a significantly higher percentage of shoots elongated enough to be rooted was detected in shoots growing under blue LED at a light intensity of $61.09 \mu\text{molm}^{-2}\text{s}^{-1}$. However, the acclimatization percentage was higher in shoots previously cultivated under FL at a light intensity of $61.71 \mu\text{molm}^{-2}\text{s}^{-1}$. Anatomical studies using light microscopy and SEM showed that light treatments promoted differences in anatomical aspects in *in vitro* shoots; in this sense, needles of plantlets exposed to red and blue LEDs revealed less stomata compared with needles from plantlets exposed to FL.

RESUMEN

Pinus ponderosa P. Lawson y C. Lawson es un árbol comercial y una de las especies forestales más importantes de América del Norte. El pino Ponderosa presenta dificultades en cuanto a propagación vegetativa y, en algunos casos, se necesitan entre 15 y 30 años para alcanzar la plena capacidad reproductiva. Basándonos en trabajos previos sobre regeneración de *P. ponderosa* a través de organogénesis *in vitro* y tratando de mejorar los protocolos publicados, nuestro objetivo fue analizar la influencia de diferentes tipos de explantos, medios de cultivo basal, CKs, auxinas y tratamientos de luz en el éxito de las fases de multiplicación y enraizamiento de tallos. La utilización de embriones cigóticos completos y una concentración de BA de $44 \mu\text{M}$ fue la que mejores resultados mostró en términos de supervivencia de los explantos. Para el proceso de organogénesis se seleccionaron embriones cigóticos completos y el medio de cultivo LP [Quoirin y Lepoivre, 1977 modificado por Aitken-Christie et al. (1988)] a la mitad de la concentración de macronutrientes. Se encontró una interacción positiva estadísticamente significativa entre los embriones cigóticos completos y el medio de cultivo LP la mitad de macronutrientes para la variable porcentaje de explantos que forman brotes. Con respecto a los tratamientos de luz aplicados, se detectó un porcentaje significativamente mayor de brotes lo suficientemente alargados como para enraizarse, en brotes que crecieron bajo LED azul a una intensidad de luz de $61,09 \mu\text{molm}^{-2}\text{s}^{-1}$. Sin embargo, el porcentaje de aclimatación fue mayor en los brotes previamente cultivados bajo luces fluorescentes a una intensidad lumínica de $61,71 \mu\text{molm}^{-2}\text{s}^{-1}$. Los estudios anatómicos que usaron microscopía óptica y microscopía electrónica de barrido mostraron que los tratamientos con luz promovieron diferencias en aspectos anatómicos en brotes *in vitro*; las acículas de plántulas expuestas a LED rojos y azules presentaban menos estomas en comparación con las acículas de plántulas expuestas a luces fluorescentes.

1. INTRODUCTION

Conifers cover approximately 39% of the world forests and are the best known and most economically important species among gymnosperms; with a total of around 615 species, including pines (Armenise et al., 2012; Farjon, 2018).

Ponderosa pine (Western Yellow Pine) is the largest of the Western pine species and is a major lumber tree in the Western North America; its natural range includes every state West of the Great Plains and North into Western Canada and South into Mexico (Ellis and Bilderback, 1991; Raish et al., 1997). In the West of the United States, it is considered one of the most fire-resistant conifers. This species develops a protective outer corky bark early in life and presents long needles with high moisture content surrounding the terminal buds, which helps to protect the apical meristems, allowing branch tips to refoliate (Hall, 1980; Miller, 2000; Fitzgerald, 2005).

The species is a major source of timber; ponderosa pine forests are also important as wildlife habitats, for recreational use and for esthetic values (Ellis and Bilderback, 1991; Raish et al., 1997). However, climate change conditions can provoke problems because this species is susceptible to diseases such as Diplodia tip blight (*Diplodia pinea*), Dothistroma blight (*Dothistroma pini*) and Elytroderma needle blight (*Elytroderma deformans*); although some authors consider ponderosa pine to be relatively disease resistant (Ellis and Bilderback, 1984, 1991).

The time taken for conifers to reach sexual maturity is long, often 20-30 years; therefore, considerable time is required before superior offspring appear (Ellis and Bilderback, 1984; 1991). Grafting, rooting cuttings, and tissue culture are vegetative propagation methods used to improve this process (Ellis and Bilderback, 1991). In this sense, *in vitro* methods, including SE or shoot organogenesis followed by rooting, provide valuable tools that can be used for the propagation of ponderosa pine (Phillips, 2004). Quite often, *in vitro* culture has low success rates of propagation due to the multitude of physico-chemical conditions influencing plant growth. Factors, such as the type and concentration of PGRs, basal media, solidifying agents, type of explant, among others, can be manipulated to induce or optimize the whole process (Phillips, 2004; Kulus and Woźny, 2020).

Among these physico-chemical factors, light is one of the most important factors controlling plant growth and development (Alallaq et al., 2020). In general, FL has been used as the main source of lighting, with irradiances between 25 and 150 $\mu\text{molm}^{-2}\text{s}^{-1}$ for a 16 h photoperiod (Larraburu et al., 2018). However, the power consumption of this kind of lamp is high, making the process expensive, and culture rooms need to remove the heat emitted by them using air conditioners together with FL efficiency (Kulus and Woźny, 2020). For these reasons, the use of LEDs is recommended as they present advantages over FL, such as durability, low power consumption, the possibility to fix specific wavelengths, smaller size and negligible heat production (Fang and Jao, 2000).

Previous works dealing with the micropropagation *P. ponderosa* are scarce. Ellis and Bilderback (1984) studied multiple bud formations culturing zygotic embryos. Years later, Ellis and Judd (1987) analyzed protein profiles on bud-forming cotyledons. In 1990, Tuskan et al. (1990) investigated the influence of PGRs, basal media and carbohydrate levels. Other authors investigated *in vitro* formation of axillary buds from immature shoots (Lin et al., 1991), and the influence of different culture media on bud induction in whole embryos and cotyledons (Ellis and Bilderback, 1991). Finally, the effect of CKs on plastid development and photosynthetic polypeptides during organogenesis has also been investigated (Mazari and Camm, 1993).

In our lab, several studies have been developed to optimize SE (Castander-Olarieta et al., 2022a) and organogenesis (Moncaleán et al., 2005; Montalbán et al., 2011) in many species of *Pinus*. Moreover, as a result of priming, somatic cells in conifers have showed different metabolic, epigenetic and proteomic profiles related to abiotic stress tolerance (Pereira et al., 2021a; Castander-Olarieta et al., 2022b; Do Nascimento et al., 2022). However, as far as we know, no work has been done in order to

improve the micropropagation of *P. ponderosa* by testing the chemical composition of the culture media or physical factors, such as type of lights (FL versus LEDs) and their effect on morphogenesis, tissue micromorphology, and the overall success of the process.

Considering the abovementioned information, the main objective of our study was to optimize the organogenesis process of *P. ponderosa* to develop an efficient protocol for *in vitro* propagation. To carry out this objective, we focused on improving (1) the multiplication phase using different types of explants, CKs, culture media, and light treatments, and (2) the rooting stage using different auxins and light treatments.

2. MATERIALS AND METHODS

2.1. Plant Material

P. ponderosa P. Lawson and C. Lawson seeds were obtained from 14-year-old trees grown in a clonal seed orchard of the Rotonda, Trevelin Experimental Field, (Argentina; 43°06'05" N, 71°33'30" W) (batch number: 25 U7519 JP) and they were also purchased from the Instituto Nacional de Tecnología Agropecuaria (Trevelin, Argentina).

2.2. Sterilization

Seeds were rinsed under running water for 5 min and sterilized following two different protocols: 1) commercial bleach 5% (active chlorine 37 gL⁻¹ sodium hypochlorite) plus one drop of Tween 20® for 20 min, followed by three rinses in sterile distilled water for 5 min each; 2) 10% H₂O₂ (v/v) (30% active H₂O₂) plus one drop of Tween 20® for 20 min, followed by three rinses in sterile distilled water for 5 min each. The sterilization protocols were performed under sterile conditions in a laminar flow unit. Seeds were stored in moistened sterile filter paper at 4 °C in darkness. After four days, seed coats and megagametophytes were removed aseptically and two explant types (a) cotyledons (Figure 1A) or whole zygotic embryos (Figure 1B) were used.




Figure 1. Plant material at different stages of *Pinus ponderosa* organogenic process: **(A)** cotyledon cultured for 2 weeks in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with 22 μM 6-benzyladenine (BA), bar=1.5 mm; **(B)** whole zygotic embryo cultured for 1 week in LP supplemented with 22 μM BA, bar=1.5 mm; **(C)** developing shoots from zygotic embryo cultured for 4 weeks in half LP macronutrients supplemented with 13.1 μM BA and exposed to red LEDs, bar=3.0 mm; **(D)** shoots after 6 weeks in elongation medium, which consisted of hormone-free LP supplemented with 2 gL^{-1} activated charcoal, bar=5.0 mm; **(E)** rooted shoots after 2 weeks in root induction medium, which consisted of half LP macronutrients supplemented with a mixture of 5 μM 1-naphthaleneacetic acid (NAA) and 5 μM indole-3-butyric acid (IBA), bar=5.0 mm; **(F)** acclimatized shoots exposed to fluorescent light in *in vitro* conditions after 4 weeks in *ex vitro* conditions in the greenhouse, bar=20 mm.

2.3. Organogenic Process

2.3.1. Experiment 1

After sterilization, explants (cotyledons or whole zygotic embryos) were cultivated on Petri dishes (90 × 15 mm) containing 20 mL of bud IM (Figure 1A, B). Two basal media were assayed: LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] and half LP macronutrients (HLP), which consisted of LP with half macronutrients. Both media were supplemented with 3% (w/v) sucrose and solidified with 8 gL^{-1} Difco Agar granulated. Moreover, three concentrations of BA (4.4, 22 and 44 μM) were evaluated. The pH of all media was adjusted to 5.8 before autoclaving (121 °C, 20 min). Cotyledons were excised from embryos and placed horizontally onto the IM; whole zygotic embryos were cultured in an inverted position with the cotyledons immersed in the IM.

As soon as bud induction was observed (after four weeks), the explants were transferred to Petri dishes containing 20 mL of EM. EM consisted of hormone-free LP or HLP supplemented with 2 gL^{-1} activated charcoal, 3% (w/v) sucrose, and was solidified with 8.5 gL^{-1} Difco Agar granulated. After 30 days in culture (when elongating needle fascicles were evident), explants were

subcultured into baby food jars with Magenta TM b-cap lids containing 25 mL of LP or HLP elongation medium. The shoots were subcultured every six weeks into the same medium. When shoots were 10-15 mm long, they were separated and cultivated individually in a fresh medium.

All the cultures were laid on the growth chamber at a temperature of 21 ± 1 °C, at a 16 h photoperiod with $120 \mu\text{molm}^{-2}\text{s}^{-1}$ light intensity provided by cool FL tubes (TLD 58 W/33; Philips, France).

2.3.2. Experiment 2

Based on the results of Experiment 1, sterilization protocol B and whole zygotic embryos were used as initial explant and HLP was selected to carry out this experiment. Two types of CKs were tested: BA and m-T at the same concentration ($13.1 \mu\text{M}$). Three different light treatments were also tested for the conditions named above: A) blue light (peak wavelength 470 nm), $61.09 \mu\text{molm}^{-2}\text{s}^{-1}$; B) red Light (peak wavelength 630 nm), $61.09 \mu\text{molm}^{-2}\text{s}^{-1}$; and C) white light (color temperature 4000 K), $61.09 \mu\text{molm}^{-2}\text{s}^{-1}$, all of them provided by adjustable LEDs (RB4K Grow LEDs). As a control, whole zygotic embryos were cultured under cool white FL (TLD 58 W/33; color temperature 4100 K) at $61.71 \mu\text{molm}^{-2}\text{s}^{-1}$ light intensity. The cultures were laid on the growth chamber at a temperature of 21 ± 1 °C for five weeks at a 16 h photoperiod.

As soon as bud induction was observed (after four weeks) (Figure 1C), the explants were transferred to Petri dishes with HLP elongation medium. After 30 days in culture (when elongating needle fascicles were evident), explants were subcultured into baby food jars with Magenta TM b-cap lids containing 25 mL of the same medium (Figure 1D). The shoots were subcultured every six weeks into the same medium. When shoots were 10-15 mm long, they were separated and cultivated individually. The different lighting conditions described previously were maintained in this phase.

2.3.3. Rooting and Adaptation

After the elongation phase, shoots of at least 20-25 mm long were used for RI. The explants were cultivated into baby food jars with Magenta TM b-cap lids containing 25 mL of RIM, which consisted of HLP supplemented with three different auxins: A) $5 \mu\text{M}$ NAA; B) $10 \mu\text{M}$ NAA; and C) a mixture of $5 \mu\text{M}$ NAA and $5 \mu\text{M}$ IBA; all media were supplemented with 3% (w/v) sucrose and solidified with 8.5 gL^{-1} Plant Agar®. The different light conditions were the same as described above. After four weeks of culture in medium A or two weeks in medium B and C, explants were cultured in baby food jars with Magenta TM b-cap lids containing 25 mL of REM, which consisted of hormone-free HLP supplemented with 2 gL^{-1} activated charcoal, 3% (w/v) sucrose, and 8.5 gL^{-1} Plant Agar® for six weeks. Growth chamber temperature and photoperiod were the same as those described above.

After six weeks in REM, explants with visible roots (Figure 1E) were transferred to wet peat moss (Pindstrup, Ryomgård, Denmark): vermiculite (8:2, v/v) and acclimatized in the greenhouse under controlled conditions at 21 ± 1 °C with progressively decreasing humidity for one month from 95% to 80% (Figure 1F). Prior to acclimatization and following the procedure described in Castander et al. (2020), the plants that developed a poor root system were transferred to Ecoboxes (Eco2box/green filter: a polypropylene vessel with a “breathing” hermetic cover (Duchefa®, Duchefa Biochemie, Haarlem, Netherlands) containing perlite:peat (1:1, v/v) moistened with liquid HLP.

3. Morphological Characterization of Needles

3.1. Light Microscopy Analysis

For structural analysis after eighteen weeks in the elongation medium, 16 needles were used following the procedure described in Giacomolli et al. (2022). Two needles per CK and light treatment were fixed in 0.2 M phosphate buffer (pH 7.2) and 2.5% paraformaldehyde for 48 h at 4 °C. Then, the samples were washed twice with 0.1 M phosphate buffer (pH 7.2) for 15 min and then dehydrated in ethanolic series (30%, 50%, 70%, 80%, 90%, 95%, and 100% v/v) (1 h each).

Subsequently, the samples were embedded in paraffin by means of dehydration with ethanol and Clear Rite™ and the consecutive immersion in liquid paraffin at 65 °C, as described in Rossi et al. (2006). The dehydration process was the following: ethanol 70% (v/v) for 120 min; ethanol 90% (v/v) for 90 min (2×); ethanol 95% (v/v) for 90 min; ethanol 100% (v/v) for 90 min (×2); ethanol: Clear Rite™ (1:1 v/v) and pure Clear Rite™ for 90 min (2×). Then, samples were embedded in paraffin wax at 65 °C for 120 min (2×) before the inclusion in paraffin molds. After paraffin inclusion (HistoDream EW, Milestone Medical Sorisole, Italy), sections of 8-10 µm were obtained in a rotary microtome (Microm HM 340E, Thermo Scientific, Waltham, USA) and transferred to microscope slides previously prepared with albumin glycerol and kept at 30 °C for 12 h. Next, the deparaffinization was performed with Clear Rite™ for 20 min, 100% ethanol for 4 min, and washing under running water. The samples were stained with an Astra Blue (0.15%) and Safranin (0.04%) water solution for 10 min; and finally, they were observed in an optical microscope (Leica DM 4000 B, Mannheim, Germany) and photos were taken with a Leica camera (Leica Application Suite version 4.13).

3.2. Scanning Electron Microscopy

For SEM, the analyses were performed following the procedure described in Giacomolli et al. (2022). Two needles per CK and light treatment and two from adult trees were fixed in 0.1 M phosphate buffer (pH 7.2) and 2.5% paraformaldehyde for 24 h at 4 °C. After that, the samples were washed twice with 0.1 M phosphate buffer (pH 7.2) for 15 min and then dehydrated in a series of ethanol at different concentrations (30%, 50%, 70%, 80%, 90%, 95%, and 100% v/v) for 15 min each.

All the samples were cut into small pieces and prepared following the procedure described in Marques et al. (2021). Then, the samples were placed on carbon stickers above metallic stubs, observed without further preparations in freeze conditions (-20 °C) at 10.0 kV, and were analyzed and photographed using a variable pressure scanning electron microscope (Flex SEM 1000, Hitachi, Tokyo, Japan).

4. Data Collection and Statistical Analysis

4.1. Experiment 1

Three Petri dishes and eight to ten explants per Petri dish (cotyledons or whole zygotic embryos) per sterilization protocol, were cultured in each culture medium and BA concentration. Contamination, survival and EFS percentages for each condition tested were measured after two months of culture. When the axillary shoots were isolated and cultured individually in an elongation medium, the EFS and the mean number of shoots were calculated with respect to the non-contaminated explants. A logistic regression model was used to analyze the effect of the sterilization protocol, explant type, culture medium, and BA concentration on survival and EFS (%). When necessary, Tukey's post hoc test ($\alpha=0.05$) was used for multiple comparisons.

4.2. Experiment 2

Five Petri dishes and eight whole zygotic embryos per Petri dish, CK type and light treatment were cultured in HLP. When the axillary shoots were isolated and cultured individually in the EM medium, the EFS, the NS/E and percentage of shoots elongated sufficiently to be rooted (PSR) out of the total NS/E were calculated. A logistic regression model was used to analyze the effect of the CK type and light treatment on survival and EFS. When necessary, Tukey's post hoc test ($\alpha=0.05$) was used for multiple comparisons.

The confirmation of the homogeneity of variances and normality of the data on the NS/E and PSR were performed, and PSR was \sqrt{x} transformed to meet homocedasticity. Data for the NS/E and PSR were analyzed by ANOVA. When necessary, multiple comparisons were made using Tukey's post hoc test ($\alpha=0.05$).

A completely randomized design was carried out using six to twenty plantlets per CK type and light treatment per each auxin treatment. The RI percentage, the mean NR/E, and the LLR (cm) were recorded after six weeks of culture in a REM.

To assess the effect of the CK and light treatment on the RI, a logistic regression was performed. Data for NR/E and LLR were analyzed by ANOVA. When necessary, multiple comparisons were made using Tukey's post hoc test ($\alpha=0.05$). The acclimatization percentage was calculated after four weeks under *ex vitro* conditions. The data were analyzed using R Core Team software® (version 4.2.1, Vienna, Austria).

5. RESULTS

5.1. Induction of Organogenesis

5.1.1. Experiment 1

Contamination rates were registered after sterilization protocols A (21%) and B (16%). The sterilization protocol did not significantly affect the survival percentage. Based on this result, protocol B was selected as the preferred sterilization method in Experiment 2.

When the type of initial explant was analyzed, a significantly higher survival percentage was observed when using whole zygotic embryos (89%) instead of isolated cotyledons (77%).

Explants' survival was also significantly affected by BA concentration used (Supplementary Tables S1 and S2); explants cultured in 44 μM BA showed significantly higher survival rates (86%) than those induced at 4.4 μM BA (Figure 2).

The interaction between BA concentration and culture medium, and between explant type and sterilization protocol did not reveal statistically significant differences (Supplementary Tables S1 and S2).




Figure 2. Percentage of survival of *Pinus ponderosa* explants cultured in LP or half LP macronutrients medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with 6-benzyladenine (BA). Data are presented as mean values±S.E. Different letters indicate significant differences according to Tukey's post hoc test ($p<0.05$).

When considering all the variables analyzed (sterilization protocol, explant type, culture medium, and BA concentration) and the interaction between them, after four weeks of IM, statistically significant differences were only found for the percentage of EFS depending on the culture medium, the explant type and the interaction between them (Supplementary Table S3). Whole zygotic embryos cultured in HLP showed significantly higher EFS than both the embryos cultured in LP and cotyledons, regardless of the culture medium employed (Figure 3). When we used cotyledon as initial explants, buds developed slowly and failed to elongate. Based on these results, whole zygotic embryos and HLP were selected as the best conditions for shoot organogenesis in Experiment 2.




Figure 3. Percentage of explants forming shoots of *Pinus ponderosa* cotyledons or whole zygotic embryos cultured in LP or half LP macronutrients (HLP) [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with 6-benzyladenine (BA). Data are presented as mean values±S.E. Different letters indicate significant differences according to Tukey's post hoc test ($p<0.05$).

The highest number of shoots (3.31 ± 2.8) were found in whole zygotic embryos cultured in HLP supplemented with $22 \mu\text{M}$ BA.

5.1.2. Experiment 2

In this experiment, the contamination rate registered was 1%. As shown in Supplementary Table S4, statistically significant differences were found for the survival percentage according to CK type and light treatment. However, Tukey's post hoc test did not detect these differences because the p-value was bordering on significance. The survival percentage was 93% in whole zygotic embryos cultured with m-T in the culture medium and growing under blue LEDs. On the other hand, 100% survival was found in whole zygotic embryos growing under the remaining treatments tested.

After five weeks of culture in IM, no significant differences were found for the percentage of EFS according to the CK type, the light treatment, or the interaction between them (Supplementary Table S5). The EFS ranged from 43% in whole zygotic embryos induced with m-T under white LEDs to 60% in whole zygotic embryos cultured with BA under blue and red LEDs, and whole zygotic embryos induced with m-T under FL.

Regarding the NS/E, no significant differences were observed for the variables studied (CK type and light treatment), or the interaction between them (Supplementary Table S6). The NS/E produced ranged from 5.40 ± 1.07 in whole zygotic embryos cultured with $13.1 \mu\text{M}$ m-T under FL to 9.11 ± 1.55 in whole zygotic embryos induced with $13.1 \mu\text{M}$ BA and exposed to red LEDs. However, at the end of the elongation stage of shoot organogenesis, a significantly higher percentage of PSR was obtained in explants developed under blue LEDs (48%) than in those exposed to red LEDs (16%); explants grown under white LEDs or FL showed intermediate values (Figure 4).




Figure 4. Percentage of shoots elongated sufficiently to be rooted (PSR) of *Pinus ponderosa* growing on half LP macronutrients medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with 6-benzyladenine (BA) under light treatments (blue LEDs (B), fluorescent light (F), red LEDs (R), and white LEDs (W)). Data are presented as mean values \pm S.E. Different letters indicate significant differences according to Tukey's post hoc test ($p < 0.05$).

On the other hand, the CK type and the interaction between CK type and light treatment did not have a significant effect on the PSR (Supplementary Table S7).

No statistically significant differences were observed in shoots exposed to different light treatments for RI, NR/E or the LLR (Supplementary Table S8). The RI percentage ranged from 3% in shoots cultured with 5 μ M NAA to 18% in shoots grown with a mixture of 5 μ M NAA and 5 μ M IBA. Shoots induced with 10 μ M NAA showed an intermediate value of RI (12%).

The highest response for RI was obtained in shoots exposed to white LEDs and the lowest response was recorded in shoots under blue LEDs (Table 1). When shoots were exposed to white LEDs, the highest NR/E was obtained, and the lowest response was found in shoots under red LEDs. The longest primary roots (ranging from 2.78 to 2.15) were recorded in shoots exposed to white LEDs and FL, respectively. Shoots growing under red LEDs displayed the lowest response (Table 1).

Table 1. Root induction (%), number of roots per explant, and length of the longest root of *Pinus ponderosa* shoots cultured in half LP macronutrients medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] according to light treatment (blue LEDs, red LEDs, white LEDs, and fluorescent light).

Light treatment	Root induction (%)	Number of roots per explant	Length of the longest root (cm)
Blue LEDs	7 \pm 30	1.71 \pm 0.29	1.60 \pm 0.77
Red LEDs	8 \pm 40	1.25 \pm 0.25	0.98 \pm 0.34
White LEDs	19 \pm 80	2.40 \pm 0.87	2.78 \pm 1.64
Fluorescent	8 \pm 40	2.00 \pm 0.71	2.15 \pm 1.35

Data are presented as mean values \pm S.E.

Acclimatized shoots were successfully obtained from shoots exposed to FL (50%), shoots under white LED (20%), and shoots exposed to blue LEDs (14%); it was not possible to acclimatize shoots grown under red LED.

5.2. Morphological Characterization of Needles

The effect of the chemical composition of the culture media and the physical factors on tissue micromorphology from needles of *in vitro* plantlets was studied, and then it was compared to the micromorphology from needles of *ex vitro* plants. In this way, adaxial and cross sections of *P. ponderosa* needles from plantlets cultured with different CK types and light treatment were analyzed. Regardless of the CK type employed, needles from plantlets did not show microscopic alterations in the internal structure and organization.

Adaxial needle surfaces from *in vitro* plantlets indicated the presence of stomata arranged in parallel and uniseriate bands, while the margin areas were serrated with spinose teeth (Figure 5A).

The longitudinal section of needles showed the parenchyma mesophyll cells and the large intercellular spaces between them. No structural changes in the mesophyll cell of needles from plantlets exposed to light treatments were found (Figure 5B). Furthermore, in the center of the needle cross section, differentiated vascular bundles with phloem and xylem were identified (Figure 5C).

Histological analysis of needles from *in vitro* plantlets showed that the epidermis presented a single-cell layer and numerous tannin-rich cells were spread among the mesophyll, where large intercellular spaces were common (Figure 5D). No clear differentiation of the endodermis and the vascular bundle was observed (Figure 5D).




Figure 5. Scanning electron microscopy and histological observations of needles from *in vitro* *Pinus ponderosa* plantlets: **(A)** adaxial surface of a needle showing teeth (t) and two rows of stomata (st); **(B)** longitudinal section of a needle showing the parenchyma mesophyll cells (p) and the large intercellular spaces (ip) between them; **(C)** SEM observation of a needle cross section; **(D)** histological observation of needle cross section. There is poor differentiation of the vascular bundle (vb), phloem (ph), and xylem (xy) as well as of the endodermis (en). Tannin-rich cells (t) are spread among the mesophyll and large intercellular spaces (ip) are common. ab-abaxial surface; ad-adaxial surface; e-epidermis.

As a general trend, needles of plantlets exposed to blue and red LEDs showed less stomata compared with the stomata of needles from plantlets grown under FL (Figures 6A, C, H). In terms of arrangement and number of stomata, when the adaxial surface of a needle growing under FL was analyzed, a regular pattern in the needle was found (Figure 6H), such as mentioned above (Figure 5A).

The morphology of the stomata of needles from shoots grown under white LEDs (Figure 6E) was similar in the pattern of arrangement to those coming from needles of shoots grown under FL (Figure 6H). Regarding their shape, stomata were ovoidal in all light conditions. The scanning electron micrograph of adaxial needle surfaces showed spinose teeth at the margins (Figures 6A, C, E, F, H). No microscopic alterations in the morphological characterization of the spinose teeth of needles from plantlets grown under different light treatments were observed (Figures 6A, C, E, H).

The internal structure and organization of needles from plantlets was similar for all light treatments (Figures 6B, D, G, I).




Figure 6. Scanning electron microscopy observations of needles from *in vitro* *Pinus ponderosa* plantlets exposed to different light treatments: **(A)** adaxial surface of a needle showing teeth (t) and two rows of stomata (st); **(B)** needle cross section; **(C)** adaxial surface of a needle showing teeth (t) and two rows of stomata (st); **(D)** needle cross section; **(E)** adaxial surface of a needle showing teeth (t) and two rows of stomata; **(F)** close view of the stomata. A tooth can also be observed; **(G)** needle cross section; **(H)** adaxial surface of a needle showing teeth (t) and two rows of stomata (st); **(I)** needle cross section. ab-abaxial surface; ad-adaxial surface.

SEM analysis showed similar anatomical aspects when needles from field-growing plants were compared with needles of *in vitro* plantlets (Figure 7A). The characterization of needles from field-growing plants revealed that the epidermal cells were arranged in a tightly packed single layer and the presence of a thin layer of cuticle on their outer surfaces was also observed (Figure 7A).




Figure 7. Scanning electron microscopy details of the organization of *Pinus ponderosa* needle from field-growing plants: **(A)** middle zone showing the organization of the vascular and surrounding tissues; **(B)** higher magnification of a resin duct (rc); **(C)** starch grains (s) in the parenchyma cells (transfusion tissue) surrounding the vascular bundles; **(D)** ridge-like invaginations (w) typical of the cell walls of the mesophyll parenchyma cells. ab-abaxial surface; ad-adaxial surface; c-cuticle; e-epidermis; en-endodermis; h-hypodermis; ph-phloem; rc-resin duct; t-transfusion tissue; xy-xylem.

The middle zone showed the organization of the vascular bundle and surrounding tissues; xylem, phloem, endodermis, and transfusion tissue were recognizable (Figure 7A).

The resin duct was generally localized adjacent to the epidermis and partially immersed in the mesophyll cells (Figure 7B). The presence of starch grains in the parenchyma cells (transfusion tissue) surrounding the vascular bundles was also distinguished (Figure 7C). The cell's morphology revealed the presence of ridge-like invaginations typical of the cell walls of the mesophyll parenchyma cells and the hypodermis was also identified (Figure 7D).

6. DISCUSSION

The establishment of an aseptic *in vitro* culture is one of the most important factors of plant tissue culture and the effective elimination of contamination contributes to the better development of the explants (Payghamzadeh and Kazemitabar, 2011; Pereira et al., 2021b). When introducing *P. ponderosa* material *in vitro*, contamination rates ranging from 1% to 16% were obtained using hydrogen peroxide (protocol B). In *P. radiata* (Montalbán et al., 2011; 2012), hydrogen peroxide was an effective agent for surface sterilization of seeds, obtaining contamination percentages below 7%. Our results suggest that the protocol used in this work was adequate for this species and can be used routinely, being a valid procedure for the propagation of other *Pinus* species as well. The most important initial step in the micropropagation process is an adequate sterilization protocol that minimizes tissue damage to achieve the greatest number of aseptic explants ready to be used in the next phases of micropropagation (De Souza et al., 2017; Pereira et al., 2021b).

As has been recently reviewed, factors such as explant source and PGRs influence the *in vitro* regeneration of plants via organogenesis in different species (Parzymies, 2021; Rojas-Vargas et al., 2021; Tung et al., 2021). In this regard, other authors mentioned that cotyledon explants had several advantages over embryonic explants (Aitken et al., 1981). First, when using cotyledons, the explant area that is exposed to the culture medium is bigger than when culturing whole zygotic embryos. Second, most seeds have at least eight cotyledons, which could be distributed among the various treatments (Saborio et al., 1997). On the contrary, in *P. halepensis* when using whole embryos or cotyledons as initial explants, over 90% of the embryos gave rise to adventitious buds, whereas isolated cotyledons slowly developed adventitious buds and these buds failed to elongate (Lambardi et al., 1993). These results are in accordance with our study. When we used cotyledon as initial explants, buds developed slowly and failed to elongate. Additionally, De Diego et al. (2011) selected embryos as initial explants for *P. pinaster* and mentioned that this would increase the genetic diversity as well as considerably decrease the manual labor cost.

In our study, the capacity of explants to form shoots was significantly higher when whole zygotic embryos were cultured in HLP. Following the trend observed in previous studies (Sul and Korban, 2004; Zhang et al., 2006), significant differences in SI rates were found when different culture media were tested for induction in *Pinus* species. Furthermore, several studies in *P. halepensis*, *P. pinaster*, *P. ayacahuite*, and *P. ponderosa* reported that salt concentration in the culture medium had an influence on morphogenesis (Ellis and Judd, 1987; Lambardi et al., 1993; Saborio et al., 1997; De Diego et al., 2011).

In woody species, it has been recognized that CKs are necessary to promote *in vitro* multiple shoot formation (Valverde-Cerdas et al., 2008; De Diego et al., 2011). Among CKs, BA is the most used in plant tissue culture due to its effectiveness and affordability (Bairu et al., 2007).

In our experiments, BA at the concentrations tested (4.4, 22, and 44 μM) showed efficient organogenic response for number of shoots, and, although not significant, a slightly higher response in whole zygotic embryos cultured in HLP supplemented with 22 μM BA was observed. A similar tendency was observed in several micropropagation protocols in *Pinus* where the culture medium was supplemented with BA (from 1 to 50 μM) to obtain bud induction (Sul and Korban, 2004; Moncaleán et al., 2005; Alonso et al., 2006; Montalbán et al., 2013; Pereira et al., 2021b; Rojas-Vargas et al., 2022).

When EFS and NS/E were analyzed, no significant differences were found for the effect of BA or m-T, but a slightly higher response in explants cultured with BA was observed. BA in *in vitro* culture can cause hyperhydricity, shoot-tip necrosis, and histogenic instability (Chen et al., 2020). This was not the case in our work, where BA did not provoke *in vitro* abnormalities. In this sense, in the last years, many reports have described the use of topolins in plant tissue culture, because topolins improve parameters, such as multiplication rate, alleviating physiological disorders and increasing the rooting percentage (Aremu et al., 2012). In

contrast to our study, in *Aloe polyphylla*, the use of m-T resulted in a high multiplication rate (Bairu et al., 2007). Likewise, Werbrouck et al. (1996) found better shoot/root balance when m-T was used in *Spathiphyllum floribundum*. In the same way, in *P. sylvestris* and *P. pinaster*, a higher rate of organogenic response was obtained by using m-T instead of BA (De Diego et al., 2010, 2011). However, in other studies of our group, no improvement was found when using m-T instead of BA for the induction of axillary shoots in *P. radiata* and *S. sempervirens* (Montalbán et al., 2011, 2013; Rojas-Vargas et al., 2021). Summarizing, the effect of different CKs at different concentrations depends on the species and the initial explant used to initiate the *in vitro* culture.

The influence of light in regulating growth, development, and adventitious root formation has been confirmed in different studies (Chen et al., 2020; dos Santos et al., 2022). LED showed positive effects in *in vitro* plants responses, when used as an alternative to conventional lighting (Gupta and Jatothu, 2013; Chen et al., 2020). Several studies have indicated the stimulatory effect of red or blue LEDs on shoot organogenesis (Gupta and Agarwal, 2017). In this work, no statistically significant difference was found in EFS when whole zygotic embryos were cultured under different light treatments. In contrast to our work, *Gerbera jamesonii in vitro* shoots growing under red LEDs displayed the greatest elongation rates (Pawłowska et al., 2018). Chen et al. (2020) in *Passiflora edulis* reported an increase in chlorophyll content and plantlet quality when plants were grown under red LEDs. Additionally, the highest stem length was observed in *in vitro* potato plantlets under red LEDs (Chen et al., 2020).

When studying the effect of the light treatment on PSR, shoots growing under blue LEDs showed the highest response. These results agree with those observed in *Curculigo orchoides* and *Zingiber officinale* where blue LEDs promoted *in vitro* shoot formation (Gupta and Sahoo, 2015; Gnasekaran et al., 2021). Furthermore, Chen et al. (2020) observed the greatest stem diameter and the highest health index in *Solanum tuberosum* when cultured under blue LEDs.

Studies performed in *Z. officinale* and *Phalaenopsis pulcherrima* indicated that LED treatments promoted *in vitro* RI and development (Gnasekaran et al., 2021; Chashmi et al., 2022). In our experiments, when the LED was tested, no significant differences were found for the RI, NR/E and LLR but a slightly higher response in shoots under white LEDs was found. In contrast, in *Z. officinale* and *Citronella mucronata* a higher rooting response was found when cultured under red or the combination of red and blue LEDs in a ratio of 1:2 (Gnasekaran et al., 2021; Guerra et al., 2022).

In *Cucumis metuliferus*, explants showed high RI rates when exposed to white LEDs and in a mixed circuit of blue and red (Lai et al., 2022). Furthermore, *Cedrela odorata in vitro* cultures grown under white LEDs presented a higher number of roots formed (dos Santos et al., 2022). In this regard, Lai et al. (2022) explained that one reason for this result might be that white light provides with a wider variety of minor lights, which are also required by plants for optimal growth.

Contrary to our results, in *P. radiata*, *in vitro* shoots growing under FL showed high NR/E (Rojas-Vargas et al., 2022). Regarding LLR, shoots under white LEDs and FL light showed longer primary roots. A similar pattern was reported in *Handroanthus ochraceus* where plants showed higher root length values when exposed to low FL irradiances (Larraburu et al., 2018). In contrast, longer roots were obtained in plants of *P. radiata* and *Vitis ficifolia* growing under red LEDs (Poudel et al., 2008; Rojas-Vargas et al., 2022).

The light treatments tested promoted differences in anatomical aspects of the stomata in the needles of *P. ponderosa in vitro* shoots. The morpho-anatomic characteristics of needles from *in vitro* and *ex vitro* plants were similar and are in accordance with the study described for *P. halepensis* by Boddi et al. (2002).

The teeth structures on the needle surface were morphologically like those found in primary needles of *P. halepensis* and *P. nigra* (Boddi et al., 2002; Mitić et al., 2017). In this sense, Boddi et al. (2002) described teeth or spine-like structures in transverse sections as more elongated cells with cytoplasmic content differing from the adjacent epidermic cells for thicker walls; however,

the relevance of these structures is unknown. A scanning electron micrograph revealed stomata arranged in parallel and as uniseriate bands, which is in accordance with the work described for *P. nigra* by Mitić et al. (2017). Similar to that observed in *P. halepensis* (Boddi et al., 2002), large and ovoidal stomata were detected in this study. In contrast, stomata in primary needles of *P. canariensis* presented a volcanic shape (Stabentheiner et al., 2004).

SEM analysis revealed the presence of hypodermis in needles from field-growing plants. Contrary to our results, in *P. halepensis* seedlings, primary needles were characterized by the absence of a hypodermal cell layer and the epidermis being in direct contact with the mesophyll tissue, but these differences could be attributed to the fact that these analyses were performed in seedlings 22–24 weeks after emergence (Boddi et al., 2002).

LED sources improve the leaf anatomy of *in vitro* plantlets in *Musa* spp.; the abaxial epidermis was thinner than the adaxial epidermis (Vendrame et al., 2022). In *Chrysanthemum* leaves, the thickness of the adaxial epidermal cells increased under white LED; blue LED favored the anatomical development of the palisade parenchyma layer; and red LED reduced this parenchyma (Zheng and Van Labeke, 2018). Red LED also affected the epidermis morphology in *Solanum tuberosum* plantlets (Chen et al., 2020). Epidermises with greater thickness were induced and palisade parenchyma and spongy parenchyma were arranged neatly (Chen et al., 2020). However, LED illumination did not improve morpho-anatomic characteristics, including vascular bundles from *in vitro* cultures of *P. ponderosa*.

In *Pfaffia glomerata in vitro* plantlets, combinations of red and blue LEDs increased the size of vascular bundles and vessel elements (Silva et al., 2020). In the same way, the central vascular bundle of plants from *Epidendrum fulgens* grown with natural ventilation under combinations of white/high blue, deep red/white/medium blue, and deep red/white/far red/medium blue LED had a higher content of sclerenchyma (Fritsche et al., 2022). Furthermore, Gnasekaran et al. (2022) found that LED spectral quality alters plant chloroplast ultrastructure through the effects on starch accumulation in *Z. officinale*. Summarizing, the effect of different LEDs sources at different concentrations in morpho-anatomic characteristics seems to depend on the species cultured *in vitro*.

All these aspects could explain the results obtained in our study, where the basal media, CKs, the explant type and the light quality influenced the organogenic process and root formation of *P. ponderosa* and their *ex vitro* acclimatization. Furthermore, our results demonstrated that different LEDs can be strategically used to improve micropropagation efficiency and reduce the costs of *in vitro* ponderosa pine plant production.

7. CONCLUSIONS

The regeneration of *P. ponderosa* through organogenesis using whole zygotic embryos as explants was achieved. The best shoot induction percentage was obtained when whole zygotic embryos were cultured in half LP macronutrients. Following the novel use of LEDs for organogenesis, a higher number of rootable shoots were obtained when blue light was used. However, our results suggest that white LED could be more beneficial for rooting stages. Finally, in the greenhouse, the shoots cultured under FL showed the highest acclimatization success (50%). This may indicate that the use of different kinds of lights could be necessary during the process to optimize the organogenic process in *P. ponderosa*. Concerning the anatomy studies, this was the first microscopy assessment of *in vitro* plantlets exposed to LEDs treatments. The needles from *in vitro* and *ex vitro* plants showed similar morpho-anatomic characteristics.

It would be interesting for future experiments to test other combinations and concentrations of auxins for rooting induction, and other LEDs, such as a combination of red and blue LEDs, to improve the rooting and acclimatization of shoots.

8. REFERENCES

- Aitken, J.; Horgan, K.J.; Thorpe, T.A. Influence of explant selection on shoot forming capacity of juvenile tissue of *Pinus radiata*. *Can. J. For. Res.* **1981**, *11*, 112–117. <https://doi.org/10.1139/x81-015>.
- Aitken-Christie, J.; Singh, A.P.; Davis, H. Multiplication of meristematic tissue: A new tissue culture system for radiata pine. In *Genetic Manipulation of Woody Plants*; Hanover, J.W., Keathley, D.E., Eds.; Plenum Publishing Corp: New York, NY, USA, 1988; pp. 413–432. https://doi.org/10.1007/978-1-4613-1661-9_26.
- Alallaq, S.; Ranjan, A.; Brunoni, F.; Novák, O.; Lakehal, A.; Bellini, C. Red light controls adventitious root regeneration by modulating hormone homeostasis in *Picea abies* seedlings. *Front. Plant Sci.* **2020**, *11*, 586140. <https://doi.org/10.3389/fpls.2020.586140>.
- Alonso, P.; Moncaleán, P.; Fernández, B.; Rodríguez, A.; Centeno, M.L.; Ordás, R.J. An Improved micropropagation protocol for Stone Pine (*Pinus pinea* L.). *Ann. For. Sci.* **2006**, *63*, 879–885. <https://doi.org/10.1051/forest:2006071>.
- Aremu, A.O.; Bairu, M.W.; Doležal, K.; Finnie, J.F.; Van Staden, J. Topolins: A panacea to plant tissue culture challenges? *Plant Cell Tissue Organ Cult.* **2012**, *108*, 1–16. <https://doi.org/10.1007/s11240-011-0007-7>.
- Armenise, L.; Simeone, M.C.; Piredda, R.; Schirone, B. Validation of DNA barcoding as an efficient tool for taxon identification and detection of species diversity in Italian conifers. *Eur. J. For. Res.* **2012**, *131*, 1337–1353. <https://doi.org/10.1007/s10342-012-0602-0>.
- Bairu, M.W.; Stirk, W.A.; Dolezal, K.; Van Staden, J. Optimizing the micropropagation protocol for the endangered *Aloe polyphylla*: Can meta-topolin and its derivatives serve as replacement for benzyladenine and zeatin? *Plant Cell Tissue Organ Cult.* **2007**, *90*, 15–23. <https://doi.org/10.1007/s11240-007-9233-4>.
- Boddi, S.; Bonzi, L.M.; Calamassi, R. Structure and ultrastructure of *Pinus halepensis* primary needles. *Flora* **2002**, *197*, 10–23. <https://doi.org/10.1078/0367-2530-00008>.
- Castander-Olarieta, A.; Moncaleán, P.; Montalbán, I.A. *Pinus canariensis* plant regeneration through somatic embryogenesis. *For. Syst.* **2020**, *29*, 1–6. <https://doi.org/10.5424/fs/2020291-16136>.
- Castander-Olarieta, A.; Moncaleán, P.; Montalbán, I.A. Somatic Embryogenesis in Pines. In *Somatic Embryogenesis. Methods in Molecular Biology*; Ramírez-Mosqueda, M.A., Ed.; Humana: New York, NY, USA, 2022a; Volume 2527, pp. 41–46. https://doi.org/10.1007/978-1-0716-2485-2_4.
- Castander-Olarieta, A.; Pereira, C.; Mendes, V.M.; Correia, S.; Manadas, B.; Canhoto, J.; Montalbán, I.A.; Moncaleán, P. Thermopriming-associated proteome and sugar content responses in *Pinus radiata* embryogenic tissue. *Plant Sci.* **2022b**, *321*, 111327. <https://doi.org/10.1016/j.plantsci.2022.111327>.
- Chashmi, K.A.; Omran, V.O.G.; Ebrahimi, R.; Moradi, H.; Abdosi, V. Light quality affects protocorm-like body (PLB) formation, growth and development of in vitro plantlets of *Phalaenopsis pulcherrima*. *Biol. Bull. Acad. Sci.* **2022**, *49*, 476–484. <https://doi.org/10.1134/S1062359022050107>.

- Chen, L.; Zhang, K.; Gong, X.; Wang, H.; Gao, Y.; Wang, X.; Zeng, Z., Hu, Y. Effects of different LEDs light spectrum on the growth, leaf anatomy, and chloroplast ultrastructure of potato plantlets in vitro and minituber production after transplanting in the greenhouse. *J. Integr. Agric.* **2020**, *19*, 108–119. [https://doi.org/10.1016/S2095-3119\(19\)62633-X](https://doi.org/10.1016/S2095-3119(19)62633-X).
- Chen, Y.C.; Chang, C.; Lin, H.L. Topolins and red light improve the micropropagation efficiency of passion fruit (*Passiflora edulis* Sims) “Tainung No. 1”. *HortScience* **2020**, *55*, 1337–1344. <https://doi.org/10.21273/HORTSCI15078-20>.
- De Diego, N.; Montalbán, I.A.; Moncaleán, P. In vitro regeneration of adult *Pinus sylvestris* L. trees. *S. Afr. J. Bot.* **2010**, *76*, 158–162. <https://doi.org/10.1016/j.sajb.2009.09.007>.
- De Diego, N.; Montalbán, I.A.; Paloma, M. Improved micropropagation protocol for maritime pine using zygotic embryos. *Scand. J. For. Res.* **2011**, *26*, 202–211. <https://doi.org/10.1080/02827581.2011.559174>.
- De Souza, R.A.; Padilha Dantas, P.V.; Cavalcante, P.D.F.; Tenório, R.R.; Houllou, L.M. Basic procedure for the in vitro propagation of brazilian trees for reforestation purposes. *J. Environ. Anal. Prog.* **2017**, *2*, 107–114. <https://doi.org/10.24221/jeap.2.2.2017.1139.107-114>.
- Do Nascimento, A.M.M.; Montalbán, I.A.; Llamazares De Miguel, D.; Goicoa, T.; Ugarte, M.D.; Moncaleán, P. High temperature and water deficit cause epigenetic changes in somatic plants of *Pinus radiata* D. Don. *Plant Cell Tissue Organ Cult.* **2022**, *151*, 107–121. <https://doi.org/10.1007/s11240-022-02336-y>.
- dos Santos Junior, C.F.; Rech, T.D.; Costa, M.D.; Navroski, M.C.; Boff, P.; Boff, M.I.C. Efeito da qualidade de luz na multiplicação e enraizamento in vitro de *Cedrela odorata* L. *Ciênc. Florest.* **2022**, *32*, 2419–2432. <https://doi.org/10.5902/1980509866513>.
- Ellis, D.D.; Bilderback, D.E. Multiple bud formation by cultured embryos of *Pinus ponderosa*. *J. Plant Physiol.* **1984**, *115*, 201–204. [https://doi.org/10.1016/S0176-1617\(84\)80120-0](https://doi.org/10.1016/S0176-1617(84)80120-0).
- Ellis, D.; Bilderback, D.E. Ponderosa pine (*Pinus ponderosa* P. Lawson & C. Lawson). In *Biotechnology in Agriculture and Forestry 16, Trees III*; Bajaj, Y., Ed.; Springer: Berlin, Germany, 1991; pp. 339–357. https://doi.org/10.1007/978-3-662-13231-9_21.
- Ellis, D.D.; Judd, R.C. SDS-PAGE analysis of bud-forming cotyledons of *Pinus ponderosa*. *Plant Cell Tissue Organ Cult.* **1987**, *11*, 57–65. <https://doi.org/10.1007/BF00036576>.
- Fang, W.; Jao, R.C. A review on artificial lighting of tissue cultures and transplants. In *Transplant Production in the 21st Century*; Kubota, C., Chun, C., Eds.; Springer: Dordrecht, Netherlands, 2000; pp. 108–113. https://doi.org/10.1007/978-94-015-9371-7_17.
- Farjon, A. The kew review conifers of the world. *Kew Bull.* **2018**, *73*, 8. <https://doi.org/10.1007/s12225-018-9738-5>.
- Fitzgerald, S. Fire ecology of Ponderosa pine and the rebuilding of fire-resilient ponderosa pine ecosystems. In *Proceedings of the Symposium on Ponderosa Pine: Issues, Trends, and Management, Klamath Falls, Oregon, USA, 18–21 October 2005*; Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture: California, USA, 2005; pp. 197–225. <https://doi.org/10.2737/PSW-GTR-198>.

Fritsche, Y.; Pinheiro, M.V.M.; Guerra, M.P. Light quality and natural ventilation have different effects on protocorm development and plantlet growth stages of the in vitro propagation of *Epidendrum fulgens* (Orchidaceae). *S. Afr. J. of Bot.* **2022**, *146*, 864–874. <https://doi.org/10.1016/j.sajb.2022.02.019>.

Giacomolli Polesi, L.; Pacheco de Freitas Fraga, H.; Goeten, D.; Panato Back, F.; de Medeiros Oliveira, E.; Steiner, N.; Guerra, M.P. Morphohistological and biochemical features of the *Guadua chacoensis* (Bambusoideae; Poaceae) somatic embryogenesis. *Plant Cell Tissue Organ Cult.* **2022**, *148*, 479–499. <https://doi.org/10.1007/s11240-021-02199-9>.

Gnasekaran, P.; Rahman, Z.A.; Chew, B.L.; Appalasamy, S.; Mariappan, V.; Subramaniam, S. Development of micropropagation system of *Zingiber officinale* var. *rubrum* Theilade using different spectrum light-emitting diode (LED) irradiation. *Ind. Crops Prod.* **2021**, *170*, 113748. <https://doi.org/10.1016/j.indcrop.2021.113748>.

Gnasekaran, P.; Rahman, Z.A.; Chew, B.L.; Uddain, J.; Mariappan, V.; Appalasamy, S.; Subramaniam, S. Regulation of chloroplast ultrastructure, adventitious bud anatomy and clonal fidelity of in vitro *Zingiber officinale* var. *rubrum* Theilade plantlets in response to LED spectral quality. *Ind. Crops Prod.* **2022**, *181*, 114825. <https://doi.org/10.1016/j.indcrop.2022.114825>.

Guerra, F.; Badilla, L.; Cautin, R.; Castro, M. Micropropagation of *Citronella mucronata* D. Don, a vulnerable chilean endemic tree species. *Plants* **2022**, *11*, 2425. <https://doi.org/10.3390/plants11182425>.

Gupta, S.; Jatothu, B. Fundamentals and applications of light-emitting diodes (LEDs) in in vitro plant growth and morphogenesis. *Plant Biotechnol. Rep.* **2013**, *7*, 211–220. <https://doi.org/10.1007/s11816-013-0277-0>.

Gupta, S.; Sahoo, T.K. Light emitting diode (LED) induced alteration of oxidative events during in vitro shoot organogenesis of *Curculigo orchioides* Gaertn. *Acta Physiol. Plant.* **2015**, *37*, 233. <https://doi.org/10.1007/s11738-015-1990-9>.

Gupta, S.; Agarwal, A. Influence of LED lighting on in vitro plant regeneration and associated cellular redox balance. In *Light Emitting Diodes for Agriculture: Smart Lighting*; Gupta, S., Ed.; Springer: Singapore, 2017; pp. 273–298.

Hall, F.C. Fire history-blue mountains, Oregon. In *Proceedings of the Fire History Workshop, Tucson, Arizona, USA, 20–24 October 1980*; Stokes, M.A., Dieterich, J.H., Eds.; Rocky Mountain Forest and Range Experiment Station, Forest Service, U.S. Department of Agriculture: Colorado, USA, 1980; pp. 75–81. <https://doi.org/10.2737/RM-GTR-81>.

Kulus, D.; Woźny, A. Influence of light conditions on the morphogenetic and biochemical response of selected ornamental plant species under in vitro conditions: A mini-review. *Bio Technol.* **2020**, *101*, 75–83. <https://doi.org/10.5114/bta.2020.92930>.

Lai, C.S.; Kho, Y.H.; Chew, B.L.; Raja, P.B.; Subramaniam, S. Organogenesis of *Cucumis metuliferus* plantlets under the effects of LEDs and silver nanoparticles. *S. Afr. J. Bot.* **2022**, *148*, 78–87. <https://doi.org/10.1016/j.sajb.2022.04.020>.

Lambardi, M.; Sharma, K.K.; Thorpe, T.A. Optimization of in vitro bud induction and plantlet formation from mature embryos of Aleppo Pine (*Pinus halepensis* Mill). *Vitr. Cell. Dev. Biol. Plant.* **1993**, *29*, 189–199. <https://doi.org/10.1007/BF02632034>.

Larraburu, E.E.; Correa, G.S.; Llorente, B.E. In vitro development of yellow lapacho (Bignoniaceae) using high-power light emitting diode. *Rev. Arvore.* **2018**, *42*, e420508. <https://doi.org/10.1590/1806-90882018000500008>.

- Lin, Y.; Wagner, M.R.; Heidmann, L.J. In vitro formation of axillary buds by immature shoots of Ponderosa Pine. *Plant Cell Tissue Organ Cult.* **1991**, *26*, 161–166. <https://doi.org/10.1007/BF00039938>.
- Marques, M.P.; Martins, J.; de Carvalho, L.A.E.B.; Zuzarte, M.R.; Costa, R.D.; Canhoto, C. Study of physiological and biochemical events leading to vitrification of *Arbutus unedo* L. cultured in vitro. *Trees* **2021**, *35*, 241–253. <https://doi.org/10.1007/s00468-020-02036-0>.
- Mazari, A.; Camm, E.L. Effect of cytokinins on plastid development and photosynthetic polypeptides during organogenesis of *Pinus ponderosa* Dougl. cotyledons cultured in vitro. *Plant Cell Tissue Organ Cult.* **1993**, *33*, 81–89. <https://doi.org/10.1007/BF01997602>.
- Miller, M. Fire Autecology. In *Wildland Fire in Ecosystems: Effects of Fire on Flora*; Tech. Rep. RMRS-GTR-42; Brown, J., Smith, J., Eds.; Rocky Mountain Research Station, Forest Service, U.S. Department of Agriculture: Ogden, Utah, USA, 2000; volume 2, pp 257. <https://doi.org/10.2737/RMRS-GTR-42-V2>.
- Mitić, Z.S.; Zlatković, B.K.; Miljković, M.S.; Jovanović, S.Č.; Marin, P.D.; Stojanović, G.S. First insights into micromorphology of needle epicuticular waxes of south-eastern european *Pinus nigra* J. F. Arnold populations. *Iheringia Sér. Bot.* **2017**, *72*, 373–379. <https://doi.org/10.21826/2446-8231201772306>.
- Moncaleán, P.; Alonso, P.; Centeno, M.L.; Cortizo, M.; Rodríguez, A.; Fernández, B.; Ordás, R.J. Organogenic responses of *Pinus pinea* cotyledons to hormonal treatments: BA metabolism and cytokinin content. *Tree Physiol.* **2005**, *25*, 1–9. <https://doi.org/10.1093/treephys/25.1.1>.
- Montalbán, I.A.; De Diego, N.; Moncaleán, P. Testing novel cytokinins for improved in vitro adventitious shoots formation and subsequent ex vitro performance in *Pinus radiata*. *Forestry* **2011**, *84*, 363–373. <https://doi.org/10.1093/forestry/cpr022>.
- Montalbán, I.A.; De Diego, N.; Moncaleán, P. Enhancing initiation and proliferation in Radiata Pine (*Pinus radiata* D. Don) somatic embryogenesis through seed family screening, zygotic embryo staging and media adjustments. *Acta Physiol. Plant.* **2012**, *34*, 451–460. <https://doi.org/10.1007/s11738-011-0841-6>.
- Montalbán, I.A.; Novák, O.; Rolčík, J.; Strnad, M.; Moncaleán, P. Endogenous cytokinin and auxin profiles during in vitro organogenesis from vegetative buds of *Pinus radiata* adult trees. *Physiol. Plant.* **2013**, *148*, 214–231. <https://doi.org/10.1111/j.1399-3054.2012.01709.x>.
- Parzymies, M. Nano-silver particles reduce contaminations in tissue culture but decrease regeneration rate and slows down growth and development of *Aldrovanda vesiculosa* explants. *Appl. Sci.* **2021**, *11*, 3653. <https://doi.org/10.3390/app11083653>.
- Pawłowska, B.; Żupnik, M.; Szewczyk-Taranek, B.; Cioć, M. Impact of LED light sources on morphogenesis and levels of photosynthetic pigments in *Gerbera jamesonii* grown in vitro. *Hortic. Environ. Biotechnol.* **2018**, *59*, 115–123. <https://doi.org/10.1007/s13580-018-0012-4>.
- Payghamzadeh, K.; Kazemitabar, S.K. In vitro propagation of walnut—A review. *Afr. J. Biotechnol.* **2011**, *10*, 290–311. <https://doi.org/10.5897/AJB10.324>.

Pereira, C.; Castander-Olarieta, A.; Sales, E.; Montalbán, I.A.; Canhoto, J.; Moncaleán, P. Heat stress in *Pinus halepensis* somatic embryogenesis induction: Effect in DNA methylation and differential expression of stress-related genes. *Plants* **2021a**, *10*, 2333. <https://doi.org/10.3390/plants10112333>.

Pereira, C.; Montalbán, I.A.; Pedrosa, A.; Tavares, J.; Pestryakov, A.; Bogdanchikova, N.; Canhoto, J.; Moncaleán, P. Regeneration of *Pinus halepensis* (Mill.) through organogenesis from apical shoot buds. *Forests* **2021b**, *12*, 363. <https://doi.org/10.3390/f12030363>.

Phillips, G.C. In vitro morphogenesis in plants—Recent advances. *In Vitro Cell. Dev. Biol. Plant.* **2004**, *40*, 342–345. <https://doi.org/10.1079/IVP2004555>.

Poudel, P.R.; Kataoka, I.; Mochioka, R. Effect of red- and blue-light-emitting diodes on growth and morphogenesis of grapes. *Plant Cell Tissue Organ Cult.* **2008**, *92*, 147–153. <https://doi.org/10.1007/s11240-007-9317-1>.

Quoirin, M.; Lepoivre, P. Études des milieux adaptés aux cultures in vitro de *Prunus*. *Acta Hortic.* **1977**, *78*, 437–442. <https://doi.org/10.17660/ActaHortic.1977.78.54>.

Raish, C.; Yong, W.; Marzluff, J. Contemporary human use of south western ponderosa pine forests. In *Songbird Ecology in South Western Ponderosa Pine Forests: A Literature Review*; Block, W.M., Finch, D.M., Eds.; Rocky Mountain Research Station, Forest Service, U.S. Department of Agriculture: Colorado, USA, 1997; pp. 28–42. <https://doi.org/10.2737/RM-GTR-292>.

Rojas-Vargas, A.; Castander-Olarieta, A.; Moncaleán, P.; Montalbán, I.A. Optimization of the micropropagation of elite adult trees of *Sequoia sempervirens*: Forest species of interest in the Basque Country, Spain. *Rev. Bionatura.* **2021**, *6*, 1511–1519. <https://doi.org/10.21931/RB/2021.01.01.11>.

Rojas-Vargas, A.; Castander-Olarieta, A.; Montalbán, I.A.; Moncaleán, P. Influence of physico-chemical factors on the efficiency and metabolite profile of adult *Pinus radiata* D. Don bud organogenesis. *Forests* **2022**, *13*, 1455. <https://doi.org/10.3390/f13091455>.

Rossi, S.; Anfodillo, T.; Menardi, R. Trephor: A new tool for sampling microcores from tree stems. *IAWA J.* **2006**, *27*, 89–97. <https://doi.org/10.1163/22941932-9000013>.

Saborio, F.; Dvorak, W.S.; Donahue, J.K.; Thorpe, T.A. In vitro regeneration of plantlets from mature embryos of *Pinus ayacahuite*. *Tree Physiol.* **1997**, *17*, 787–796. <https://doi.org/10.1093/treephys/17.12.787>.

Silva, T.D.; Batista, D.S.; Fortini, E.A.; Castro, K.M.; Felipe, S.H.S.; Fernandes, A.M.; Sousa, R.M.J.; Chagas, K.; Silva, J.V.S.D.; Correia, L.N.F.; et al. Blue and red light affects morphogenesis and 20-hydroxyecdysone content of in vitro *Pfaffia glomerata* accessions. *J. Photochem. Photobiol. B.* **2020**, *203*, 111761. <https://doi.org/10.1016/j.jphotobiol.2019.111761>.

Stabentheiner, E.; Pfeifhofer, H.W.; Juliane Peters, M.; Jiménez, S.; Morales, D.; Grill, D. Different surface characteristics of primary and secondary needles of *Pinus canariensis*. *Flora* **2004**, *199*, 90–99. <https://doi.org/10.1078/0367-2530-00138>.

Sul, I.W.; Korban, S.S. Effects of salt formulations, carbon sources, cytokinins, and auxin on shoot organogenesis from cotyledons of *Pinus pinea* L. *Plant Growth Regul.* **2004**, *43*, 197–205. <https://doi.org/10.1023/B:GROW.0000046013.47892.4f>.

Tung, H.T.; Thuong, T.T.; Cuong, D.M.; Luan, V.Q.; Hien, V.T.; Hieu, T.; Nam, N.B.; Phuong, H.T.N.; Van The Vinh, B.; Khai, H.D.; et al. Silver nanoparticles improved explant disinfection, in vitro growth, runner formation and limited ethylene accumulation during micropropagation of strawberry (*Fragaria* × *Ananassa*). *Plant Cell Tissue Organ Cult.* **2021**, *145*, 393–403. <https://doi.org/10.1007/s11240-021-02015-4>.

Tuskan, G.A.; Sargent, W.A.; Rensema, T.; Walla, J.A. Influence of plant growth regulators, basal media and carbohydrate levels on the in vitro development of *Pinus ponderosa* (Dougl. Ex Law.) cotyledon explants. *Plant Cell Tissue Organ Cult.* **1990**, *20*, 47–52. <https://doi.org/10.1007/BF00034756>.

Valverde-Cerdas, L.; Rojas-Vargas, A.; Hine-Gómez, A. In vitro propagation of *Albizia guachapele*, *Cedrela odorata*, *Platymiscium pinnatum* and *Guaiacum sanctum*. *Plant Tissue Cult. Biotechnol.* **2008**, *18*, 151–156. <https://doi.org/10.3329/ptcb.v18i2.3397>.

Vendrame, W.A.; Feuille, C.; Beleski, D.; Bueno, P.M.C. In vitro growth responses of ornamental Bananas (*Musa* sp.) as affected by light sources. *Horticulturae* **2022**, *8*, 92. <https://doi.org/10.3390/horticulturae8020092>.

Werbrouck, S.P.O.; Strnad, M.; Van Onckelen, H.A.; Debergh, P.C. Meta-topolin, an alternative to benzyladenine in tissue culture?. *Physiol. Plant.* **1996**, *98*, 291–297. <https://doi.org/10.1034/j.1399-3054.1996.980210.x>.

Zhang, Y.; Wei, Z.M.; Xi, M.L.; Shi, J.S. Direct organogenesis and plantlet regeneration from mature zygotic embryos of masson pine (*Pinus massoniana* L.). *Plant Cell Tissue Organ Cult.* **2006**, *84*, 119–123. <https://doi.org/10.1007/s11240-005-9004-z>.

Zheng, L.; Van Labeke, M.C. Effects of different irradiation levels of light quality on *Chrysanthemum*. *Sci. Hortic.* **2018**, *233*, 124–131. <https://doi.org/10.1016/j.scienta.2018.01.033>.

CHAPTER 6

Optimization of the micropropagation of elite adult trees of *Sequoia sempervirens*: forest species of interest in the Basque Country, Spain

The content of this chapter corresponds to the published article “Rojas-Vargas, A.; Castander-Olarieta, A.; Montalbán, I.A.; Moncaleán, P. 2021. Optimization of the micropropagation of elite adult trees of *Sequoia sempervirens*: forest species of interest in the Basque Country, Spain. *Rev. Bionatura* 6, 1511–1519. <https://doi.org/10.21931/RB/2021.01.01.11>”.

ABSTRACT

Forest trees are renewable sources of timber and other valuable non-timber products. Nowadays, the increase of population and demand of forestry products results in over exploitation of forestry. Therefore, there is an urgent need to produce elite plants with higher productivity under stress derived from climate change to have available to afforestation. For this reason, propagation methods should be improved to be more efficient in terms of quality and productivity.

The main species planted in the Basque Country is *Pinus radiata*; during the last three years *P. radiata* plantations have suffered a fungus attack affecting mainly needles until the death of the tree. This crisis is caused by *Lecanosticta acicola* and the combined action of the fungi *Dothistroma septosporum* and *D. pini*, whose expansion seems to have been enhanced by weather-related factors, such as humid and hot summers. Although we have evidence of the presence of this disease in our mountains since 1942, for reasons that are not scientifically known today, the disease has had a very fast expansion with an aggressive effect. For the above, Basque Country forestry sector is looking for alternative species to be used in its plantations. Part of the forestry sector considers that *Sequoia sempervirens* could be a good choice for plantations. In addition, its high-quality wood, and its tolerance to the attack of several pathogens and other diseases derived of climate change are characteristics that could confer some advantage over other forest species.

The main goal of this study was to optimize the micropropagation of adult elite trees of *Sequoia sempervirens*. To carry out this objective, the effect of BA, m-T and KIN, and 4 types of explants in the multiplication stage were analyzed. Furthermore, the effect of two types of auxins: NAA, IBA and a mixture of both were evaluated on the induction of roots and their subsequent effect on the acclimatization process. The best multiplication index was obtained when 4.4 μM BA and apical explants longer than 1.5 cm of length were used. The RI percentage was 75% in the most responsive genotype analyzed when 4.4 μM BA was used on induction stage and 50 μM NAA was used for rooting. Finally, after 3 months in the greenhouse, the explants cultured with KIN and rooted in culture medium with IBA showed highest acclimatization success (94%).

RESUMEN

Los árboles forestales son fuentes de madera y otros valiosos productos no maderables. En la actualidad, el aumento de la población y la demanda de productos forestales se traduce en una sobreexplotación de los bosques. Por lo tanto, existe una necesidad urgente de producir plantas élite con mayor productividad bajo el estrés derivado del cambio climático para ser utilizadas en plantaciones forestales y programas de reforestación. Por esta razón, se deben mejorar los métodos de propagación tratando de que sean más eficientes en términos de calidad y productividad.

La principal especie plantada en el País Vasco es *Pinus radiata*; durante los últimos tres años las plantaciones de *P. radiata* se han visto afectadas por varias enfermedades que afectan principalmente a las acículas hasta la muerte del árbol. Esta crisis está provocada *Lecanosticta acicola* y la acción combinada de los hongos *Dothistroma septosporum* and *D. pini*, cuya expansión parece haberse visto favorecida por factores relacionados con el clima, como veranos húmedos y calurosos. Aunque tenemos constancia de la presencia de esta enfermedad en nuestras montañas desde 1942, por razones que hoy no se conocen científicamente, la enfermedad ha tenido una expansión muy rápida con un efecto agresivo. Por lo anterior, el sector forestal del País Vasco está buscando especies alternativas para utilizar en sus plantaciones. Parte del sector forestal considera que *Sequoia sempervirens* podría ser una buena opción para las plantaciones. Además, su madera de alta calidad y su tolerancia al ataque de diversos patógenos y otras enfermedades derivadas del cambio climático son características que podrían conferirle alguna ventaja frente a otras especies forestales.

El objetivo principal de este estudio fue optimizar la micropropagación de árboles adultos élitos de *S. sempervirens*. Para llevar a cabo este objetivo se analizó el efecto de distintas CKs (BA, m-T, KIN) y 4 tipos de explantos en la etapa de multiplicación. Además, se evaluó el efecto de dos tipos de auxinas: NAA, IBA y una mezcla de ambos sobre la inducción de raíces y su posterior efecto sobre el proceso de aclimatación. El mejor índice de multiplicación se obtuvo cuando se utilizaron 4,4 μM BA y explantos apicales de más de 1,5 cm de longitud. El porcentaje de inducción de raíces en el genotipo de mayor respuesta analizado fue de 75% cuando se utilizó 4.4 μM BA en la etapa de inducción y NAA (50 μM) para el enraizamiento. Finalmente, después de 3 meses en invernadero, los explantos cultivados con KIN y enraizados en medio de cultivo con IBA mostraron el mayor éxito de aclimatación (94%).

1. INTRODUCTION

The Coast redwood or California redwood *Sequoia sempervirens* (D. Don.) Endl. (Taxodiaceae) is a valuable forest species and occurs naturally in Western North America, especially in California (Mihaljević et al., 1999; Meneguzzi et al., 2019). This species is the tallest tree on earth with a high volume of standing biomass, in some stands exceeding 3,500 metric tons/hectare (Taha et al., 2014).

This conifer has been introduced and domesticated in European countries such as: Romania, Spain, France, Great Britain, Russia and Turkey, and it can be used for reforestation due to the quality of its wood as well as for ornamental purposes. In Spain, this species has been used by foresters for its productivity, its tolerance to the attack of several pathogens and other diseases. Moreover, its reforestation is recommended on valley bottoms and foothills (Otazua, 2016).

Nowadays, redwood shows seed reproduction difficulties, displaying low germination and rooting rates, dormancy of the shoot and low seedling viability (Olson et al., 1990; Ozudogru et al., 2011). Therefore, biotechnological *in vitro* techniques of plant tissue culture as micropropagation, organogenesis, adventitious or axillary shoot/bud regeneration, shoot tip culture, micrografting and SE emerged as useful tools for the propagation and conservation of germoplasm (Vengadesan et al., 2002; Bonga, 2016; De Souza et al., 2017).

In the specific case of *S. sempervirens*, there are few investigations on micropropagation, either through organogenesis or SE. Arnaud et al. (1993) developed a protocol for micropropagation and rejuvenation of the species using direct organogenesis and SE. Years later, Mihaljević et al. (1999) investigated the root formation in micropropagated shoots of *S. sempervirens* using *Agrobacterium*. On the 21st century, Sul and Korban (1994) developed a procedure for the micropropagation of juvenile and adult material and Lui et al. (2006) worked on the shoot regeneration and SE from needles of redwood. The medium-term conservation of *S. sempervirens* was investigated by Ozudogru et al. (2011), and Meneguzzi et al. (2019) evaluated shoot multiplication of two *S. sempervirens* genotypes with addition of KIN.

Considering all the abovementioned information, the objective of our study was to optimize the micropropagation process of elite selected adult trees of *S. sempervirens*. We focused on improving 1) the multiplication phase using different types of explants and CKs 2) the rooting using different auxins and the subsequent acclimatization phase under *ex vitro* conditions.

2. MATERIALS AND METHODS

2.1. Plant material

Stem sections were collected from the basal parts of three different mother trees (7, 11 and 12) of *S. sempervirens* located in Ataun, Gipuzkoa, Basque Country (Spain), at the geographic coordinates: 42° 58' 41" N and 2° 10' 53" O.

2.2. Shoot induction

In vitro microshoots growing in half strength ARN medium (Arnaud et al., 1993) were selected to carry out the experiments. Four types of explants were used: A) apical explants of 1.5 cm in length (AG), B) apical explants of 1.0 cm length (AP), C) basal explants of 1.5 cm length (BG) and D) basal explants of 1.0 cm in length (BP). Explants were cultured in glass jars (Merck) with 25 mL of ARN multiplication medium supplemented with 3% sucrose and 8 gL⁻¹ Difco Agar®. The effect of different CKs for the induction of shoots was evaluated: A) BA, B) m-T and D) KIN at a concentration of 4.4 µM. The pH of the medium was adjusted to 5.8 before autoclaving (121°C, 20 min). The explants were cultivated in the different induction media for four weeks in the growth chamber

at a temperature of $21\pm 1^{\circ}\text{C}$, under 16-h photoperiod with $120\ \mu\text{molm}^{-2}\text{s}^{-1}$ of light intensity provided by cool FL tubes (TLD 58 W/33; Philips, France).

2.3. Shoot elongation

The explants induced in the multiplication phase were cultivated for six weeks in shoot EM. This medium was half strength ARN (Arnaud et al., 1993) supplemented with $2\ \text{gL}^{-1}$ activated charcoal and solidified with $9\ \text{gL}^{-1}$ Difco Agar[®]. The pH of the medium was adjusted to 5.8 before autoclaving (121°C , 20 min).

2.4. Root induction and acclimatization of plants

After the elongation phase, stems with at least 2.5 cm in length were used for RI. Traceability of the explant was maintained according to its origin (genotype and CK treatment used along the SI stage). Subsequently, the stems were grown in RIM, which consisted of 1/3 strength ARN basal medium supplemented with three different auxin treatments: A) $50\ \mu\text{M}$ NAA, B) $50\ \mu\text{M}$ IBA and C) a mixture of $40\ \mu\text{M}$ NAA μM and $10\ \mu\text{M}$ IBA. All the different media were solidified with $9\ \text{gL}^{-1}$ Difco Agar[®]. The pH of the culture medium was adjusted to 5.8 before autoclaving (121°C , 20 min).

The stems were placed in the dark at $21\pm 1^{\circ}\text{C}$ for 8 days. After this period, the stems were cultured in REM, which consisted of 1/3 strength ARN basal medium (Arnaud et al., 1993) without plant growth regulators and solidified with $9\ \text{gL}^{-1}$ Difco agar[®]. The pH of the medium was adjusted to 5.8 before autoclaving (121°C , 20 min). The cultures were placed in the growth chamber at a temperature of $21\pm 1^{\circ}\text{C}$, under 16-h photoperiod at $120\ \mu\text{molm}^{-2}\text{s}^{-1}$. After six weeks of culture in REM, explants with visible roots were transferred to wet peat: vermiculite mixture (2:1, v/v) and acclimatized in the greenhouse under controlled conditions at $21\pm 1^{\circ}\text{C}$ and decreasing humidity progressively along one month from 95 to 80%.

2.5. Data collection and statistical analysis

To assess the effect of the genotype on each of the variables of this study an analysis of ANOVA was performed, followed by a Tukey's post hoc test ($\alpha=0.05$). The genotype had a significant effect in all variables studied. For this reason and to obtain robust conclusions, the genotype factor was introduced into all the models as a block variable to reduce variability and analyze the effect of the other variable factors (type of explant, CK treatment and auxin treatment) more accurately.

The SI and the NS/E were recorded after six weeks. The experiment used a completely a randomized design with 25 explants per treatment. To assess the effect of the type of explant and the CK treatment on the SI percentage a logistic regression was performed, followed by a Tukey's post hoc test ($\alpha=0.05$) for multiple comparisons. The NS/E variable did not fulfill homocedasticity and normality assumptions for ANOVA, then a Kruskal-Wallis was performed.

The RI (%), the NR/E and the root length was recorded after six weeks in REM. A completely randomized design was carried out using ten stems per CK treatment. As exception, in genotype 7, four stems were employed when KIN was used in induction stage. A logistic regression model was used to analyze the effect of the auxin and CK treatment on the RI percentage. Tukey's post hoc test ($\alpha=0.05$) was used for multiple comparisons. The NR/E variable did not fulfill homocedasticity and normality assumptions for ANOVA, then a Kruskal-Wallis was performed.

Data on the LLR root were square root transformed, and an ANOVA was performed, multiple comparisons were made using the Tukey's post hoc test ($\alpha=0.05$)

After twelve weeks under *ex vitro* conditions the survival percentage was calculated. Survival percentage variable did not fulfill homocedasticity and normality assumptions for ANOVA, then a Kruskal-Wallis was performed.

3. RESULTS

3.1. Shoot induction

When the effect of the genotype on the SI percentage was evaluated, genotypes 12 and 11 showed significantly higher percentages (98.98%) than genotype 7 (79%, Table 1). As shown in Table 1, the number of shoots obtained in genotype 12 (5.04) was significantly higher than those obtained in genotypes 11 and 7 (3.21 and 3.28, respectively).

Table 1. Shoot induction (%) and number shoots per explant from *Sequoia sempervirens* elite adult trees cultured in ARN medium (Arnaud et al., 1993).

Variable	Genotype		
	7	11	12
Shoot induction (%)	79.00±0.03 ^b	98.98±0.01 ^a	98.98±0.01 ^a
N° shoots /explant	3.28±0.12 ^b	3.21±0.07 ^b	5.04±0.16 ^a

Data are presented as mean values±S.E. Different letters within a row indicate significant differences ($p<0.05$).

The analysis of deviance of the factors studied showed a significant effect on both the SI and the NS/E developed from each explant and CK type. In the case of the NS/E there was a significant interaction between the explant type and the CK type (Table 2).

Table 2. Analysis of deviance for shoot induction (%) and number shoots per explant from *Sequoia sempervirens* elite adult trees cultured in ARN medium (Arnaud et al., 1993).

Source	df	Shoot induction (%)		N° shoots /explant	
		X ² Test	p-Value	X ² Test	p-Value
Explant type (E)	3	382.39	0.00011*	23.63084	2.98290 ^{5*}
Cytokinin type (C)	2	403.24	0.01861*	142.0222	<2.0 ^{-16*}
E x C	6	374.86	n.s	188.8565	<2.0 ^{-16*}

*Significant differences at $p<0.05$, n.s Non-significant at $p<0.05$, df Degrees of freedom.

Significantly higher SI percentages were obtained in AP, BG and BP explants (between 91% and 94%) than in AG explants (84%) (Table 3).

Table 3. Shoot induction (%) in different explant types from *Sequoia sempervirens* elite adult trees cultured in ARN medium (Arnaud et al.; 1993) (apical explants of 1.5 cm in height (AG), apical explants of 1.0 cm height, C) basal explants of 1.5 cm height (BG) and basal explants of 1.0 cm in height (BP).

Explant type	Shoot induction (%)
AG	84.00±0.02 ^b
AP	94.00±0.02 ^a
BG	91.00± 0.02 ^a
BP	94.00±0.02 ^a

Data are presented as mean values±S.E. Different letters indicate significant differences ($p<0.05$).

When considering the effect of the CK type on SI percentages, shoots induced with m-T showed significantly higher induction percentages than shoots induced with KIN. Shoots induced in the presence of BA displayed intermediate values (92%) (Table 4).

Table 4. Shoot induction (%) in different cytokinin type [6-benzyladenine (BA), kinetin (KIN) and meta-topolin (m-T)], from *Sequoia sempervirens* elite adult trees cultured in ARN medium (Arnaud et al., 1993).

Cytokinin type	Shoot induction (%)
BA	92.00±0.02 ^{ab}
KIN	87.00±0.02 ^b
m-T	93.00±0.01 ^a

Data are presented as mean values±S.E. Different letters indicate significant differences ($p<0.05$).

When evaluating the NS/E, a significant interaction between explant type and CK was observed. The best results were obtained in AG, AP and BG explants induced with m-T and in AG explants induced with BA (Figure 1). When BA or m-T were used for induction, the worst results were obtained in BP explants. The lowest response was observed in treatment KIN, independently of the explant type.




Figure 1. Number of shoots per explant in different explant types from *Sequoia sempervirens* elite adult trees cultured in ARN medium (Arnaud et al., 1993) supplemented with different cytokinins [6-benzyladenine (BA), kinetin (KIN) and meta-topolin (m-T)], (apical explants of 1.5 cm in length (AG), apical explants of 1.0 cm length (AP), basal explants of 1.5 cm length (BG) and basal explants of 1.0 cm in length (BP)). Data are presented as mean values±S.E. Different letters indicate significant differences ($p<0.05$).

Despite there were no statistically significant differences between BG/m-T, AP/m-T, and AG/BA, it is important to mention that the appearance of induced shoots was different (Figures 2A-C). The shoots induced in presence of BA and KIN were greener and more vigorous (Figure 2A) than the explants cultured in m-T (Figures 2B, C). These explants from m-T treatments were more yellowish and less vigorous (Figures 2B and C). Therefore, AG explants cultured in presence of BA showed the best results for SI.




Figure 2. Shoot induction from *Sequoia sempervirens* elite adult trees cultured in ARN multiplication medium (Arnaud et al., 1993) cultured for 6 weeks: **(A)** apical explant >1.5 cm of length (AG) of genotype 7 on ARN medium supplemented with 4.4 μ M 6-benzyladenine (BA), bar=3 mm ; **(B)** basal explant >1.5 cm of length (BG) of genotype 7 on ARN medium supplemented with 4.4 μ M meta-topolin (m-T), bar=5 mm; **(C)** apical explant <1.5 cm of length (AP) explant of genotype 7 on ARN medium supplemented with 4.4 μ M m-T, bar=3 mm.

Since AG explants showed the best appearance, a comparative figure was done to describe the aspect of regenerated shoots in the presence of CKs (Figure 3). The most vigorous shoots were observed in the presence of BA and KIN (Figures 3A-C, G-H).




Figure 3. Shoot induction from *Sequoia sempervirens* elite adult trees cultured in ARN multiplication medium (Arnaud et al., 1993) cultured for 6 weeks: **(A)** apical explant >1.5 cm of length (AG) of genotype 7 on ARN medium supplemented with 4.4 μ M 6-benzyladenine (BA), bar=3 mm; **(B)** AG of genotype 11 on ARN medium supplemented with 4.4 μ M BA, bar=3 mm; **(C)** AG explant of genotype 12 on ARN medium supplemented with 4.4 μ M BA, bar=3 mm; **(D)** AG explant of genotype 7 on ARN medium supplemented with 4.4 μ M meta-topolin (m-T), bar=3 mm; **(E)** AG explant >1.5 of genotype 11 on ARN supplemented with 4.4 μ M m-T, bar=5 mm; **(F)** AG explant of genotype 12 on ARN medium supplemented with 4.4 μ M m-T, bar=5 mm; **(G)** AG of genotype 7 on ARN medium supplemented with 4.4 μ M kinetin (KIN), bar=3 mm; **(H)** AG explant of genotype 11 on ARN medium supplemented with 4.4 μ M KIN, bar=5 mm; **(I)** AG of genotype 12 on ARN medium supplemented with 4.4 μ M KIN, bar=3 mm.

3.2. Root induction

A significantly higher RI percentage was obtained in genotype 12 (93%) (Table 5). The lowest response was shown by genotype 7 (31%) and intermediate values were recorded for genotype 11 (55%). Also, significantly higher NR/E was observed in genotype 12 (4.73) when compared with genotype 7 and 11 (Table 5).

Table 5. Root induction (%), number roots per explant and length of longest root from *Sequoia sempervirens* elite adult trees cultured in ARN medium (Arnaud et al.,1993).

Variable	Genotype		
	7	11	12
Root induction (%)	31.25±0.04 ^c	55.42±0.04 ^b	93.47±0.02 ^a
N ^o roots /explant	2.14±0.27 ^b	1.69±0.11 ^b	4.73±0.25 ^a
Length of longest root (cm)	2.95±0.21 ^a	3.73±0.33 ^a	2.18±0.12 ^b

Data are presented as mean values±S.E. Different letters within a row indicate significant differences ($p<0.05$).

On the contrary, significantly longer primary roots were recorded for genotype 11 (3.73 cm, Figure 4A) and genotype 7 (2.95 cm, Figure 4B) than for genotype 12 (2.18 cm, Figure 4C).




Figure 4. Root induction from *Sequoia sempervirens* elite adult trees micro shoots cultivated *in vitro* for 6 weeks on root expression medium (REM) (Arnaud et al., 1993): **(A)** explant of genotype 7 induced in root inducing medium (RIM) (Arnaud et al.,1993) supplemented with 50 μ M 1-naphthaleneacetic acid (NAA) (6-benzyladenine (BA) applied in shoot induction stage), bar=2 cm; **(B)** explant of genotype 11 induced in RIM medium supplemented with 50 μ M NAA (BA applied in shoot induction stage), bar=2 cm; **(C)** explant of genotype 12 induced in RIM medium supplemented with 50 μ M NAA (BA applied in shoot induction stage), bar=2 cm.

The auxin type and CK type used in multiplication stage showed a significant effect on the RI percentage. Further, the CK type showed a significant effect on the LLR (Table 6). Regarding the NR/E, no significant differences were found for the variables tested.

Table 6. Analysis of deviance for root induction (%), number roots per explant and length of longest root from *Sequoia sempervirens* elite adult trees cultured in ARN medium (Arnaud et al., 1993).

Source	df	Root induction		N° roots /explant		Length of longest root	
		X ² Test	p-Value	X ² Test	p-Value	F Test	p-Value
Auxin type (A)	2	510.96	0.01099 *	3.42422	n.s	2.703	0.06913
Cytokinin type (C)	2	503.59	0.02503*	1.15589	n.s	5.469	0.00479*
A x C	4	496.71	n.s	11.81255	n.s	1.042	n.s

*Significant differences at $p < 0.05$, n.s: Non-significant at $p < 0.05$, df Degrees of freedom.

As shown in Table 7, a significantly higher RI percentage was obtained in the treatment with NAA (70%) than in treatments with IBA or IBA/NAA mixture. As mentioned before, there was no effect of the auxin treatment on the NR/E (ranging from 2.98 to 3.94) or on the length of the primary root (ranging from 2.23 to 3.09 cm) (Table 7).

Table 7. Root induction (%), number of roots per explant and length of the longest root in *Sequoia sempervirens* elite adult trees cultured in ARN medium (Arnaud et al., 1993) supplemented with different auxin types [indole-3-butyric acid (IBA), 1-naphthaleneacetic acid (NAA)/IBA or NAA].

Variable	Auxin types		
	IBA	IBA/NAA	NAA
Root induction (%)	60.00±0.04 ^b	57.00±0.04 ^b	70.00±0.04 ^a
N° roots /explant	2.98±0.25 ^{n.s}	3.94±0.35 ^{n.s}	3.30±0.26 ^{n.s}
Length of longest root (cm)	2.77±0.21 ^{n.s}	2.23±0.17 ^{n.s}	3.09±0.25 ^{n.s}

Data are presented as mean values±S.E. Different letters within a row indicate significant differences ($p < 0.05$).n.s: Non-significant.

When evaluating the effect of the CK type over RI percentage significantly higher values were obtained from shoots induced in BA treatment when compared with those induced in m-T and KIN treatment (Table 8).

As aforementioned, no significant differences were found for the NR/E, independently of the SI treatment applied (ranging from 3.11 to 3.89) (Table 8).

The longest primary roots (3.32 cm) showed significantly higher values when shoots were induced in BA treatment (Table 8).

Table 8. Root induction (%), number roots per explant and length of longest root in different cytokinin types [6-benzyladenine (BA), kinetin (KIN) and meta-topolin (m-T)] from *Sequoia sempervirens* elite adult trees cultured in ARN medium (Arnaud et al., 1993).

Variable	Cytokinin type applied in the root induction stage		
	BA	m-T	KIN
Root induction (%)	66.00±0.04 ^a	58.00±0.04 ^b	62.00±0.04 ^b
N° roots /explant	3.11±0.24 ^{n.s}	3.89±0.33 ^{n.s}	3.16±0.29 ^{n.s}
Length of longest root (cm)	3.32±0.22 ^a	2.39±0.21 ^b	2.48±0.24 ^b

Data are presented as mean values ± S.E. Different letters within a row indicate significant differences ($p < 0.05$).n.s: Non-significant.

3.3. Acclimatization of rooted microplants

The analysis of deviance for survival (%) of rooted shoots propagated *in vitro* showed a significant effect of the auxin type applied in the RI stage and of the CK type applied in the SI phase (Table 9).

Table 9. Analysis of deviance for survival (%) of rooted shoots propagated *in vitro* coming from *Sequoia sempervirens* elite adult trees, after twelve weeks under *ex vitro* conditions.

Source	df	Survival (%)	
		X ² Test	p-Value
Auxin type (A)	2	244.40	5.679 ^{-5*}
Cytokinin type (C)	2	223.62	3.075 ^{-5*}
A x C	4	216.66	n.s

*Significant differences at $p < 0.05$, n.s: Non-significant at $p < 0.05$, df Degrees of freedom.

When the effect of auxin type on the survival percentage of rooted shoots was analyzed, the presence of IBA in the rooting medium provoked a significantly higher survival percentage (94%) than the mixture IBA/NAA and NAA treatments (Table 10).

Table 10. *Ex vitro* survival (%) of *in vitro* rooted shoots from *Sequoia sempervirens* elite adult trees, shoots were rooted in ARN medium (Arnaud et al., 1993) supplemented with different auxins [indole-3-butyric acid (IBA), 1-naphthaleneacetic acid (NAA)/IBA or NAA].

Variable	Auxin type		
	IBA	IBA/NAA	NAA
Survival %	94.00±0.03 ^a	79.00±0.06 ^b	68.00±0.05 ^b

Data are presented as mean values±S.E. Different letters indicate significant differences ($p < 0.05$).

Regarding the effect of the CK applied in the SI stage as shown in Table 11, the shoots induced in KIN treatment showed a significantly higher *ex vitro* survival percentage than those induced in BA or m-T treatments.

Table 11. *Ex vitro* survival (%) of *in vitro* rooted shoots from *Sequoia sempervirens* elite adult trees; shoots were propagated in ARN medium (Arnaud et al., 1993) supplemented with different cytokinins [6-benzyladenine (BA), kinetin (KIN) and meta-topolin (m-T)].

Variable	Cytokinin type		
	BA	m-T	KIN
Survival (%)	80.00±0.05 ^b	65.00±0.05 ^b	94.00±0.03 ^a

Data are presented as mean values±S.E. Different letters indicate significant differences by Tukey's post hoc test ($p < 0.05$).

At the end of the experiment, the micropropagated plants from the three elite adult trees evaluated showed a normal development, good growth and uniformity after 5 months under *ex vitro* conditions in the greenhouse (Figure 5).



Figure 5. Plants of *Sequoia sempervirens* cultured *in vitro* after 5 months growing under *ex vitro* conditions in the greenhouse, bar=4 cm.

4. DISCUSSION

The genotype is an endogenous factor that has an extremely important role for regenerative potential in the overall repeatability and reliability of tissue culture protocol (Coleman and Ernst, 1989). In this study, the SI, the NS/E, the RI and the NR/E were affected by the genotype of the explant. Our results are in concordance with the results reported by Meneguzzi et al. (2019), who observed different response among genotypes in shoot multiplication of *S. sempervirens*. In the same way, Sul and Korban (2005) found differences in shoot proliferation and shoot elongation of *S. sempervirens*.

The organogenesis processes are influenced by endogenous and exogenous factors (Sul and Korban, 2005; George, 2008; Aremu et al., 2012). There are different exogenous factors as the components in the tissue culture medium or the environmental conditions of culture. In this work we observed that the type of explant and chemical conditions in the culture medium (CK and auxin type) had a significant effect in SI and rooting response.

During the evaluation of the SI rate, the explants with the longest length (AG, AP and BG) showed the best induction rate and the highest NS/E were obtained. Similar results were found by Clapa et al. (2010) and Meneguzzi et al. (2020) who developed an *in vitro* protocol for *S. sempervirens* using shoot explants bigger than 1.5 cm of length. George (2008) reported that larger explants coming from larger parts of shoot apex or stem segments bearing one or more lateral buds could show advantages over smaller size explants.

Several studies about tissue culture protocols have reported positive effects on parameters such as shoot proliferation, shoot multiplication rate, alleviating physiological disorders, better rooting and acclimatization when using topolins for SI (Aremu et al., 2012; Moyo et al., 2018). In our study the explants induced in the presence of m-T exhibited the best results. In this sense, De Diego et al. (2010), obtained a high rate of organogenic response in adult buds of *Pinus sylvestris* using m-T and suggested that it could be used as an alternative CK to BA in micropropagation. Likewise, Bairu et al. (2007) obtained outstanding multiplication rates in *Aloe polyphylla* when used the same cytokinin. However, in the present study these explants from m-T treatments were more yellowish and less vigorous. In contrast, in *Prunus* rootstocks, *Pterocarpus marsupium* and *Corylus colurna* a positive effect of m-T on growth and quality of micropropagated shoots was found (Gentile et al., 2014, 2017; Ahmad and Anis, 2019).

According to Sul and Korban (2005), Meneguzzi et al. (2020), Valverde et al. (2004) and Montalbán et al. (2011) in forestry species, BA stimulated the axillary bud breakage and shoot elongation. In our study, a similar result was obtained in the BA treatment. It is important to mention that in the treatment with BA the quality of the shoots formed (robustness and color) was superior to

those shoots generated in the presence of m-T. In conclusion, in our study BA was the best CK in terms of SI and NS/E. This is in accordance with the results reported by Moncaleán et al. (2005), George (2008), Bairu et al. (2007) and Aremu et al. (2012) and confirming that BA is the most used CK in micropropagation due to its effectiveness and affordability. Regarding KIN treatment, in this work we obtained the worst results in agreement with those found in *Barleria greenii* and *Eucalyptus globulus* (Bennett et al., 1994; Amoo et al., 2011).

In this research the NAA treatment showed the best result in RI response. Our results agree with those reported in *S. sempervirens*, *Pinus radiata* *P. pinaster* and other coniferous species (De Diego et al., 2008, 2011; Halmagyi et al., 2010; Ragonezi et al., 2010; Montalbán et al., 2019). It is important to mention that the rooting percentages obtained in the present studies were higher than those recorded by Huang et al. (1992) who obtained about a 30% of rooting competence using adult stem sections of *S. sempervirens* in the presence of IAA/KIN. The NR/E obtained in the IBA/NAA treatment presented the highest values. In species such as *Eucalyptus sideroxylon*, *Rosa hybrida* and *Citrus aurantifolia* the best RI was obtained when using a mixture of IBA/NAA. Nonetheless, this treatment did not lead to the highest values in the NR/E (Kosh-Khui and Sink, 1982; Cheng et al., 1992; Al-Bahrany, 2002). The highest value in the LLR was also obtained in NAA treatment, being in accordance with the results obtained in *C. aurantifolia* (Al-Bahrany, 2002).

In our research the CK type used in induction stage had a significant effect over RI. Bairu et al. (2007), Werbrouck et al. (1996) and Aremu et al. (2012) found that m-T stimulated *in vitro* rooting activity, but in our work the m-T treatment presented the worst results. In this sense, Bairu et al. (2008) and Escalona et al. (2003) obtained negative carryover effects on rooting at too high m-T levels, so it is possible that the m-T concentration applied in our cultures might have had a detrimental effect in the subsequent stages of development.

Regarding the acclimatization stage, the highest survival percentage was observed in plantlets rooted in the presence of IBA and developed in the presence of KIN. This result agrees with those found in *E. globulus*, where low concentration of IBA showed the best survival rates; and in *Arachis paraguariensis* cultured in polyethylene terephthalate glycol vessel where IBA treatment was the best in survival (Bennett et al., 2003; Aina et al., 2015).

Aremu et al. (2012) explained that CKs generally have inhibitory effects on rooting, resulting in poor acclimatization rates afterwards. In this sense, the plants from m-T treatment showed the lowest survival. In contrast, several studies have shown that plantlets coming from m-T induction treatments have been successfully acclimatized (Bairu et al., 2007; Naaz et al., 2019).

In conclusion, the role of auxins and CKs in the micropropagation of different types of explants and their relationship with the survival and acclimatization of seedlings in *ex vitro* conditions was analyzed. The results of our study demonstrated that the apical explant ≥ 1.5 cm length, and BA showed the best results in the SI stage. Moreover, *in vitro* shoots rooted with IBA, led to a higher *ex vitro* survival. Finally, the results shown allow the development of forthcoming studies for large-scale propagation of this species in semi-solid systems and bioreactors.

5. REFERENCES

- Ahmad, A.; Anis, M. *Meta-topolin improves in vitro Morphogenesis, rhizogenesis and biochemical analysis in Pterocarpus marsupium* Roxb.: A potential drug-yielding tree. *J. Plant Growth Regul.* **2019**, *38*, 1007–1016. <https://doi.org/10.1007/s00344-018-09910-9>.
- Aina, O. O.; Quesenberry, K. H.; Gallo, M. Culture vessel and auxin treatments affect in vitro rooting and ex vitro survival of six *Arachis paraguariensis* genotypes. *Sci. Hortic.* **2015**, *183*, 167–171. <https://doi.org/10.1016/j.scienta.2014.12.006>.
- Al-Bahrany, A. M. Effect of phytohormones on in vitro shoot multiplication and rooting of lime *Citrus aurantifolia* (Christm.) Swing. *Sci. Hortic.* **2002**, *95*, 285–295. [https://doi.org/10.1016/S0304-4238\(01\)00349-1](https://doi.org/10.1016/S0304-4238(01)00349-1).
- Amoo, S. O.; Finnie, J. F.; van Staden, J. The role of meta-topolins in alleviating micropropagation problems. *Plant Growth Regul.* **2011**, *63*, 197–206. <https://doi.org/10.1007/s10725-010-9504-7>.
- Aremu, A. O.; Bairu, M. W.; Doležal, K.; Finnie, J. F.; Van Staden, J. Topolins: A panacea to plant tissue culture challenges? *Plant Cell Tissue Organ Cult.* **2012**, *108*, 1–16. <https://doi.org/10.1007/s11240-011-0007-7>.
- Arnaud, Y.; Franclet, A.; Tranvan, H.; Jacques, M. Micropropagation and rejuvenation of *Sequoia sempervirens* (Lamb) Endl: a review. *Ann. des Sci. For.* **1993**, *50*, 273–295. <https://hal.archives-ouvertes.fr/hal-00882846>.
- Bairu, M. W.; Stirk, W. A.; Dolezal, K.; Van Staden, J. Optimizing the micropropagation protocol for the endangered *Aloe polyphylla*: Can meta-topolin and its derivatives serve as replacement for benzyladenine and zeatin? *Plant Cell Tissue Organ Cult.* **2007**, *90*, 15–23 (2007). <https://doi.org/10.1007/s11240-007-9233-4>.
- Bairu, M. W.; Stirk, W. A.; Doležal, K.; Van Staden, J. The role of topolins in micropropagation and somaclonal variation of banana cultivars ‘Williams’ and ‘Grand Naine’ (*Musa* spp. AAA). *Plant Cell Tissue Organ Cult.* **2008**, *95*, 373–379. <https://doi.org/10.1007/s11240-008-9451-4>.
- Bennett, I.J.; McComb, J.A.; Tonkin, C.M.; McDavid, D.A.J. Alternating cytokinins in multiplication media stimulates in vitro shoot growth and rooting of *Eucalyptus globulus* Labill. *Ann. Bot.* **1994**, *74*, 53-58. <https://doi.org/10.1006/anbo.1994.1093>.
- Bennett, I.J.; McDavid, D.A.J.; McComb, J. The influence of ammonium nitrate, pH and indole butyric acid on root induction and survival in soil of micropropagated *Eucalyptus globulus*. *Biol. Plant.* **2003**, *47*, 355–360. <https://doi.org/10.1023/B:BIOP.0000023877.21262.a5>.
- Bonga, J. M. Conifer clonal propagation in tree improvement programs. In: *Vegetative Propagation of Forest Trees*; Park, Y.S., Bonga, J.M, Moon, H. K, Eds.; National Institute of Forest Science: Seoul, Korea, 2016; pp. 3-31.
- Cheng, B.; Peterson, C. M.; Mitchell, R. J. The role of sucrose, auxin and explant source on in vitro rooting of seedling explants of *Eucalyptus sideroxylon*. *Plant Sci.* **1992**, *87*, 207–214. [https://doi.org/10.1016/0168-9452\(92\)90152-C](https://doi.org/10.1016/0168-9452(92)90152-C).
- Clapa, D.; Fira, Al.; Hărșan, E.; Chiș L.; Roman, G. *Sequoia sempervirens* in Transylvania. *Acta Hortic.* **2010**, *885*, 85–90. <https://doi.org/10.17660/ActaHortic.2010.885.11>.

- Coleman, G.; Ernst, S. In vitro shoot regeneration of *Populus deltoides*: effect of cytokinin and genotype. *Plant Cell Rep.* **1989**, *8*, 459-462. <https://doi.org/10.1007/BF00269048>.
- De Diego, N.; Montalbán, I. A.; Fernandez De Larrinoa, E.; Moncaleán, P. In vitro regeneration of *Pinus pinaster* adult trees. *Can. J. For. Res.* **2008**, *38*, 2607–2615. <https://doi.org/10.1139/X08-102>.
- De Diego, N.; Montalbán, I. A.; Moncaleán, P. In vitro regeneration of adult *Pinus sylvestris* L. trees. *South African J. Bot.* **2010**, *76*, 158–162. <https://doi.org/10.1016/j.sajb.2009.09.007>.
- De Diego, N.; Montalbán, I. A.; Moncaleán, P. Improved micropropagation protocol for maritime pine using zygotic embryos. *Scand. J. For. Res.* **2011**, *26*, 202–211. <https://doi.org/10.1080/02827581.2011.559174>.
- De Souza, R. A.; Dantas, P. V.; Cavalcante, P. D. F.; Tenório, R. R.; Houllou, L. M. Basic procedure for the in vitro propagation of Brazilian trees for reforestation purposes. *J. Environ. Anal. Prog.* **2017**, *2*, 107–114. <http://dx.doi.org/10.24221/jeap.2.2.2017.1139.107-114>.
- Escalona, M.; Cejas, I.; Gonzáles-Olmedo, J.; Capote, I.; Roels, S.; Cañal, M.J.; Rodríguez, R.; Sandoval Fernandez, J.A.; Debergh, P. The effect of meta-topolin on plantain propagation using a temporary immersion bioreactor. *Infomusa* **2003**, *12*, 28-30. <https://www.musalit.org/seeMore.php?id=8153>.
- Gentile, A.; Jáquez Gutiérrez, M.; Martínez, J.; Frattarelli, A.; Nota, P.; Caboni, E. Effect of meta-Topolin on micropropagation and adventitious shoot regeneration in *Prunus* rootstocks. *Plant Cell Tissue Organ Cult.* **2014**, *118*, 373–381. <https://doi.org/10.1007/s11240-014-0489-1>.
- Gentile, A.; Frattarelli, A.; Nota, P.; Condello, E.; Caboni, E. The aromatic cytokinin meta-topolin promotes in vitro propagation, shoot quality and micrografting in *Corylus colurna* L. *Plant Cell Tissue Organ Cult.* **2017**, *128*, 693–703. <https://doi.org/10.1007/s11240-016-1150-y>.
- George, E.F. Plant Tissue Culture Procedure - Background. In *Plant Propagation by Tissue Culture*; George, E.F., Hall, M.A., De Klerk, G.J., Eds.; Springer: Dordrecht, The Netherlands, 2008; pp. 1–28. https://doi.org/doi.org/10.1007/978-1-4020-5005-3_1.
- Halmagyi, A.; Kevresan, S.; Kovacevic, B.; Orlovic, S.; Miladinovic, D.; Cirin-Novta, V.; Kuhajda, K. Effects of naphthenic acids on rooting of in vitro grown *Sequoia sempervirens*. *Acta Hortic.* **2010**, *885*, 139–144. <https://doi.org/10.17660/ActaHortic.2010.885.18>.
- Huang, L.; Lius, S.; Huang, B.; Murashige, T.; Mahdi, F.M.; Van Gundy, R. Rejuvenation of *Sequoia sempervirens* by repeated grafting of shoot tips onto juvenile rootstocks in vitro: Model for phase reversal of trees. *Plant Physiol.* **1992**, *98*, 166–173. <https://doi.org/10.1104/pp.98.1.166>.
- Kosh-Khui, M.; Sink, K.C. Rooting enhancement of *Rosa hybrida* for tissue culture propagation. *Sci. Hortic.* **1982**, *17*, 371–376. [https://doi.org/10.1016/0304-4238\(82\)90118-2](https://doi.org/10.1016/0304-4238(82)90118-2).
- Liu, C.; Xia, X.; Yin, W.; Huang, L.; Zhou, J. Shoot regeneration and somatic embryogenesis from needles of redwood (*Sequoia sempervirens* (D.Don.) Endl.). *Plant Cell Rep.* **2006**, *25*, 621–628. <https://doi.org/10.1007/s00299-006-0120-y>.

Meneguzzi, A.; Konzen, E. R.; Navroski, M. C.; Camargo, S. S.; Pereira, M. D. O.; Rufato, L.; Lovatel, Q. C. Shoot multiplication of two *Sequoia sempervirens* genotypes with addition of small concentrations of kinetin. *Pesqui. Florest. Bras.* **2019**, *39*, 1-8. <https://doi.org/10.4336/2019.pfb.39e201701550>.

Meneguzzi, A.; Camargo, S. S.; Navroski, M. C.; de Oliveira Pereira, M.; Souza, P. F.; Rufato, L. Agar and activated charcoal influence the in vitro development of shoot and root *Sequoia*. *Rev. Ciencias Agroveterinarias* **2020**, *18*, 20–24. <https://doi.org/10.5965/2238117118e2019020>.

Mihaljević, S.; Katavić, V.; Jelaska, S. Root formation in micropropagated shoots of *Sequoia sempervirens* using *Agrobacterium*. *Plant Sci.* **1999**, *141*, 73–80. <https://doi.org/10.1073/pnas.85.15.5536>.

Moncaleán, P.; Alonso, P.; Centeno, M. L.; Cortizo, M.; Rodríguez, A.; Fernández, B.; Ordás, R. J. Organogenic responses of *Pinus pinea* cotyledons to hormonal treatments: BA metabolism and cytokinin content. *Tree Physiol.* **2005**, *25*, 1–9. <https://doi.org/10.1093/treephys/25.1.1>.

Montalbán, I. A.; De Diego, N.; Moncaleán, P. Testing novel cytokinins for improved in vitro adventitious shoots formation and subsequent ex vitro performance in *Pinus radiata*. *Forestry* **2011**, *84*, 363–373. <https://doi.org/10.1093/forestry/cpr022>.

Montalbán, I. A.; Moncaleán, P. Rooting of *Pinus radiata* somatic embryos: factors involved in the success of the process. *J. For. Res.* **2019**, *30*, 65–71. <https://doi.org/10.1007/s11676-018-0618-5>.

Moyo, M.; Aremu, A.O.; Plačková, L.; Plíhalová, L.; Pěnčík, A.; Novák, O.; Holub, J.; Doležal, K.; Van Staden, J. Deciphering the growth pattern and phytohormonal content in Saskatoon berry (*Amelanchier alnifolia*) in response to in vitro cytokinin application. *N. Biotechnol.* **2018**, *42*, 85–94. <https://doi.org/10.1016/j.nbt.2018.02.001>.

Naaz, A.; Hussain, S. A.; Anis, M.; Alatar, A. A. Meta-topolin improved micropropagation in *Syzygium cumini* and acclimatization to ex vitro conditions. *Biol. Plant.* **2019**, *63*, 174–182. <https://doi.org/10.32615/bp.2019.020>.

Olson, D.F.; Douglass, F.R.; Walters, G.A. *Sequoia sempervirens* (D. Don) Endl. In: *Silvics of North America. 1. Conifers. Agriculture Handbook 654*; Burns, R. M, Honkala, B. H,Eds.; Forest Service, U.S. Department of Agriculture: Washington, USA, 1990; Volume 1, pp.541-551.

Otazua, F. Especies forestales. *Sequoia* de California (*Sequoia sempervirens* ENDL.). *Rev. la Asoc. For. Navarra.* **2016**, *39*, 06–09.

Ozudogru, E.A.; Kirdok, E.; Kaya, E.; Capuana, M.; De Carlo, A.; Engelmann, F. Medium-term conservation of redwood (*Sequoia sempervirens* (D. Don.) Endl.) in vitro shoot cultures and encapsulated buds. *Sci. Hortic.* **2011**, *127*, 431–435. <https://doi.org/10.1016/j.scienta.2010.10.013>.

Ragonezi, C.; Klimaszewska, K.; Castro, M. R.; Lima, M.; de Oliveira, P.; Zavattieri, M. A. Adventitious rooting of conifers: Influence of physical and chemical factors. *Trees - Struct. Funct.* **2010**, *24*, 975–992. <https://doi.org/10.1007/s00468-010-0488-8>.

Sul, I. W.; Korban, S. S. Effect of different cytokinins on axillary shoot proliferation and elongation of several genotypes of *Sequoia sempervirens*. *Vitr. Cell. Dev. Biol. – Plant.* **1994**, *30*, 131–135. <https://doi.org/10.1007/BF02632201>.

Sul, I. W.; Korban, S. S. Direct shoot organogenesis from needles of three genotypes of *Sequoia sempervirens*. *Plant Cell Tissue Organ Cult.* **2005**, *80*, 353–358. <https://doi.org/10.1007/s11240-004-1365-1>.

Taha, L. S.; Taie, H. A. A.; Metwally, S. A.; Fathy, H. M. Effect of laser radiation treatments on in vitro growth behavior, antioxidant activity and chemical constituents of *Sequoia sempervirens*. *Res. J. Pharm. Biol. Chem. Sci.* **2014**, *5*, 1024–1034.

Valverde-Cerdas, L.; Alvarado, L.; Hine, A. Micropropagation of clones from controlled crosses of *Gmelina arborea* in Costa Rica. *New For.* **2004**, *28*, 187–194. <https://doi.org/10.1023/B:NEFO.0000040945.75019.38>.

Vengadesan, G.; Ganapathi, A.; Amutha, S.; Selvaraj, N. In vitro propagation of *Acacia* species - A review. *Plant Sci.* **2002**, *163*, 663–671. [https://doi.org/10.1016/S0168-9452\(02\)00144-9](https://doi.org/10.1016/S0168-9452(02)00144-9).

Werbrouck, S.; Strnad, M.; Van Onckelen, H. Meta-topolin, an alternative to benzyladenine in tissue culture? *Physiol. Plant.* **1996**, *98*, 291–297. <https://doi.org/10.1034/j.1399-3054.1996.980210.x>

CHAPTER 7
GENERAL DISCUSSION

GENERAL DISCUSSION

In the last years the cultivated forest of *P. radiata* in the Basque Country has been affected by several diseases presumably due to climate change; in this scenario it is necessary to develop efficient propagation methods for alternative species to *P. radiata*. The best alternative to produce elite plants is the vegetative propagation. However, when plants show the desirable characteristics have reach maturity and propagation for traditional methods are impossible. For this reason, forestry countries combine traditional techniques with *in vitro* propagation methods to develop effective propagation systems.

In conifers, vegetative propagation is commonly used as a massive cloning method to obtain material from selected genotypes, to improve planting quality and to shorten the selection cycles for forest tree genetic improve programs (Niemi et al., 2004). Micropropagation is currently being used as effective method for mass cloning (Bonga, 2015), but in woody species it is widely recognized the difficulty of applying this technique; even more when explants from mature trees are used as initial explants (Giri et al., 2004).

In this work the possibility of using adult *P. radiata* trees with apparent tolerance to needle blight disease as a source of material to *in vitro* propagation was explored. As main findings it was observed that the collection date and genotype of the explants had a significant effect on the *in vitro* response, being the percentage of explants forming shoots genotype dependent. These results are in concordance with those carried out by and De Diego et al. (2008) in *P. pinaster* and Pereira et al. (2021) in *P. halepensis* in which they showed that the genotype affected the *in vitro* organogenesis response.

In this context, the objective of the present study was to develop *in vitro* propagation methods for *C. japonica*, *P. ponderosa* and *S. sempervirens*, species considered as an alternative to *P. radiata* for Basque Country forests. In this sense, this thesis contributes to the knowledge about how modifications applied in the physical and chemical environment during the different stages of micropropagation process can affect the success of the *in vitro* culture process and the morphological characteristics and metabolite profile in the plantlets obtained.

Among the modifications of the chemical environment is the addition of PGRs to the culture medium, in this work it has been verified that different types and concentrations of PGRs affect the micropropagation process. The function of CKs has extensively been studied in *in vitro* culture and their role in the suppression of apical dominance and promotion of the development of axillary buds has been recognized (Arab et al., 2014; Rodríguez et al., 2022). In this sense, although different CKs have been tested, BA was the most effective for shoot induction in elite trees of *P. radiata*, *C. japonica*, *P. ponderosa* and *S. sempervirens*. In this regard, BA is the most commonly CK used in plant tissue culture due to its effectiveness and affordability (Bairu et al., 2007; De Diego et al., 2008; Phillips and Garda, 2019). In accordance with our results, Montalbán et al. (2013) and Pereira et al. (2021) also reported a higher *in vitro* shoot organogenesis response when BA was utilized in organogenesis protocols from adult trees of different *Pinus* species. Contrary to our work, Castro-Garibay et al. (2023) in *S. sempervirens* and Cortizo et al. (2009) in *P. pinea* adult trees obtained a better organogenic response when they used KIN and TDZ, respectively. Furthermore, the use of a certain PGR during shoot induction can not only improve this stage but also influence subsequent ones including acclimatization. This thesis showed different results in relation to the CK applied along the induction period; we observed that a certain CK in the induction phase caused an improvement in the rooting phase but another of the evaluated CKs improved the acclimatization percentages as we have assessed in this work, where in *S. sempervirens* shoots induced with BA and rooted in NAA show significantly higher *in vitro* rooting; whereas plantlets induced in KIN and rooted in IBA show a significantly better acclimatization success.

All propagation methods have limitations; in micropropagation these difficulties may include low regeneration, necrosis, inability to continue elongation and low adventitious root regeneration among others (Von Aderkas and Bonga, 2000; Sarmast, 2018; Díaz-

Sala, 2019). Regarding the latter limitation mentioned, higher rooting rates were obtained when IBA and NAA were utilized in plantlets of *C. japonica* and *S. sempervirens*, respectively. According to our results, Halmagyi et al. (2010) in *S. sempervirens* also obtained higher rooting rates using NAA. Moreover, in *Cupressus sempervirens* Khamushi et al. (2019), reported that pulse treatment (30 s) in IBA and subsequent culturing in the woody plant medium supplemented with KIN and IBA was effective to root induction. These results demonstrated that the adventitious root regeneration also can be influenced by PGRs among other chemical factors (Zarei et al., 2020) and that the effect of them is species dependent.

During this work, in *P. ponderosa* and *P. radiata* the obtention of a good adventitious root system was very difficult. According with Von Aderkas and Bonga (2000), the *in vitro* conditions influence the maintenance of juvenility of the tissue. However, loss of organogenesis competence, progressive or sudden, reflects transition to the mature phase; in line with this, the low rooting response observed in these species may be due to tissue aging (Von Aderkas and Bonga, 2000). In hybrid hazelnuts (*Corylus avellana* × *Corylus americana*) the use of a temporary immersion system significantly improved the *in vitro* rooting efficiency compared with semi-solid medium (Nicholson et al., 2020). Likewise, San José et al. (2020) in *Alnus glutinosa* obtained rooted plantlets using a temporary immersion system. In this context, these recent strategies for rooting, together with testing other combinations and concentrations of auxins could be considered to improve *P. ponderosa* and *P. radiata* rooting in the future.

Light treatments during organogenesis also provoked different responses in all the species studied. Our results are in accordance with the results reported by Alallaq et al. (2020) in *Picea abies*. In this regard, *C. japonica* shoots produced a higher number of roots when they were grown under red LED lights. According to our results, Lotfi et al. (2019) reported a better rooting response *in vitro* in *Pyrus communis* plants when red LEDs were used. Likewise, *Protea cynaroides* rooting rates were the highest under red LEDs (Wu and Lin, 2012). However, this response seems to be also species dependent, for instance, in *Cucumis metuliferus* plantlets showed the best rooting response under white LEDs (Lai et al., 2022).

Significantly higher percentage of *P. ponderosa* shoots able to be rooted was detected in shoots growing under blue LEDs although the best rooting rate was observed in shoots growing under white LEDs. In this context, *Zingiber officinale* shoots exposed to blue LEDs showed the highest shoot regeneration (Gnasekaran et al., 2021). Likewise, in *Solanum tuberosum* when shoots were cultured under blue LEDs the greatest stem diameter and the highest health index were obtained (Chen et al., 2020). In contrast, Santana Costa Souza et al. (2022) studying *Eucalyptus grandis* × *E. urophylla* obtained the highest organogenic response when the *in vitro* culture was exposed to FL. Likewise, Guerra et al. (2022) using white LEDs in *Citronella mucronata* trees obtained the lowest rooting rates.

Higher root induction and number of roots per explant were observed in *P. radiata* shoots cultured under FL. Similar results were reported by Alvarenga et al. (2015) in *Achillea millefolium* where higher rooting rates were obtained in *in vitro* plants growing under FL. Contrary, in *P. cynaroides* and *Dendranthema × grandiflorum* the lowest rooting rates were obtained in plantlets growing under FL (Wu and Lin, 2012; Gil et al., 2020). According to our results Gupta and Jatothu (2013), also reported that at each stage of plant development the light requirements appeared to be different. In addition, even when applying the same light treatments different species respond differently which might be associated with the genotypic characteristics or the physiological state of explants (Da Silva et al., 2016; Lai et al., 2022). Many reports have suggested that a combination of different types of light could be interesting to improve the micropropagation process (Diaz-Rueda et al. 2021; Dewir et al. 2023), so future experiments should be made to evaluate that combination effect. Moreover, Allalaq et al. (2020) reported that different lights had a different effect in *Picea abies* vegetative propagation through modulation of hormone homeostasis, it would be interesting for future research to check the hormonal status of our tissues at different stages and light treatments.

Analysis of amino acids and carbohydrates during the rooting stage of *P. radiata* shoots were carried out to evaluate the effect of different physico-chemical factors along the organogenesis process. Fructose, glucose and sucrose levels were significantly different in the rooting stages analyzed. Carbohydrates in micropropagation process are related to the maintenance of osmotic potential and to serve as a carbon source for developmental processes such as root formation (Yaseen et al., 2013; Moradi et al., 2017). In this work, shoots with better *in vitro* rooting ability grown in culture medium supplemented with 3% sucrose concentration presented higher fructose, glucose and sucrose content. Also, Mingozi et al. (2011) in *Cydonia oblonga* reported a higher content of sucrose, glucose and fructose in shoots cultured with higher sucrose concentration.

Amino acid acts as block units for proteins, osmolytes, signaling molecules, phytohormone precursors, cofactors and amino acid levels readjust both nutritional and metabolic cell responses (Wu, 2009). In this study, higher threonine and tyrosine levels were obtained in shoots displaying the worst results in rooting percentage. Joshi et al. (2010) and Muthuramalingam et al. (2018) reported that threonine metabolites are involved in plant growth and development, cell division and responses to abiotic stresses. However, in our work, threonine level did not favor the plant growth and development. In this sense, Sarrobert et al. (2000) in an *Arabidopsis thaliana* mutant suggested that the excess of endogenous threonine is directly or indirectly toxic for rooting. Regarding to tyrosine, in *Alternanthera brasiliana* acts as precursor of secondary metabolites such as betacyanin (Silva et al., 2005).

On the other hand, it is widely recognized that the physico-chemical microenvironment of vessels used in micropropagation can induce anatomical, morphological and physiological abnormalities in plantlets (Shekhawat et al., 2017). This thesis reported the first microscopy assessment of *in vitro* plantlets from *P. ponderosa* exposed to LEDs treatments. Our analyses concluded that light treatments promoted differences in anatomical aspects of *in vitro* shoots; needles of micropropagated plants exposed to red and blue LEDs showed less stomata compared with needles from plantlets exposed to FL. In this way, Díaz-Rueda et al. (2021) reported minimal stomatal density when 70% red and 30% blue LEDs with $34 \mu\text{molm}^{-2}\text{s}^{-1}$ of light irradiance were used in *in vitro* culture of *Olea europaea*. In white strawberry (*Fragaria x ananassa*) (Pang et al., 2023), demonstrated that the stomata of the seedlings exposed to blue LED resulted damaged compared those from white, red, blue, and red+blue LEDs. As reported by Franco-Navarro et al. (2019) a lower stomatal frequency could reduce water loss through leaf transpiration and improved the *ex vitro* acclimatization. Further research is needed in our species to confirm this hypothesis.

Our work demonstrated that plant tissue culture could be used as a complementary cloning method of elite trees of *C. japonica*, *P. ponderosa*, *P. radiata* and *S. sempervirens* in future breeding program in order to improve the Basque Forest health in the current climate change scenario.

REFERENCES

- Alallaq, S.; Ranjan, A.; Brunoni, F.; Novák, O.; Lakehal, A.; Bellini, C. Red light controls adventitious root regeneration by modulating hormone homeostasis in *Picea abies* seedlings. *Front.Plant Sci.* **2020**, *11*. <https://doi.org/10.3389/fpls.2020.586140>.
- Alvarenga, I. C. A.; Pacheco, F. V.; Silva, S. T.; Bertolucci, S. K. V.; Pinto, J. E. B. P. In vitro culture of *Achillea millefolium* L.: quality and intensity of light on growth and production of volatiles. *Plant Cell Tissue Organ Cult.* **2015**, *122*, 299–308. <https://doi.org/10.1007/s11240-015-0766-7>.
- Arab, M. M.; Yadollahi, A.; Shojaeiyan, A.; Shokri, S.; Ghoghaj, S. M. Effects of nutrient media, different cytokinin types and their concentrations on in vitro multiplication of G × N15 (hybrid of almond × peach) vegetative rootstock. *J. Genet. Eng. Biotechnol.* **2014**, *12*, 81–87. <https://doi.org/10.1016/j.jgeb.2014.10.001>.

- Bairu, M. W.; Stirk, W. A.; Dolezal, K.; Van Staden, J. Optimizing the micropropagation protocol for the endangered *Aloe polyphylla*: Can meta-topolin and its derivatives serve as replacement for benzyladenine and zeatin?. *Plant Cell Tissue Organ Cult.* **2007**, *90*, 15–23. <https://doi.org/10.1007/s11240-007-9233-4>.
- Bonga, J. M. A comparative evaluation of the application of somatic embryogenesis, rooting of cuttings, and organogenesis of conifers. *Can. J. For. Res.* **2015**, *45*, 379–383. <https://doi.org/10.1139/cjfr-2014-0360>.
- Castro-Garibay, S. L.; Villegas-Monter, A.; Cruz-Larios, I. J. Concentración de Kinetina y tipo de explante en la multiplicación in vitro de *Sequoia sempervirens*. D. Don. *Endl. Polibotánica* **2023**, *55*, 81-94. <https://doi.org/10.18387/polibotanica.55.6>.
- Chen, L.; Zhang, K.; Gong, X.; Wang, H.; Gao, Y.; Wang, X.; Zeng, Z.; Hu, Y. Effects of different LEDs light spectrum on the growth, leaf anatomy, and chloroplast ultrastructure of potato plantlets in vitro and minituber production after transplanting in the greenhouse. *J. Integr. Agric.* **2020**, *19*, 108–119. [https://doi.org/10.1016/S2095-3119\(19\)62633-X](https://doi.org/10.1016/S2095-3119(19)62633-X).
- Cortizo, M.; De Diego, N.; Moncaleán, P.; Ordás, R. J. Micropropagation of adult Stone Pine (*Pinus pinea* L.). *Trees Struct. Funct.* **2009**, *23*, 835–842. <https://dx.doi.org/10.1007/s00468-009-0325-0>.
- Da Silva, J. A. T.; Winarto, B.; Dobránszki, J.; Cardoso, J. C.; Zeng, S. Tissue disinfection for preparation of *Dendrobium* in vitro culture. *Folia Hortic.* **2016**, *28*, 57–75. <https://doi.org/10.1515/fhort-2016-0008>.
- De Diego, N.; Montalbán, I. A.; Fernandez De Larrinoa, E.; Moncaleán, P. In vitro regeneration of *Pinus pinaster* adult trees. *Can. J. For. Res.* **2008**, *38*, 2607–2615. <https://doi.org/10.1139/X08-102>.
- Dewir, Y.H.; Al-Ali, A.M.; Rihan, H.Z.; Alshahrani, T.; Alwahibi, M.S.; Almutairi, K.F.; Naidoo, Y.; Fuller, M.P. Effects of artificial light spectra and sucrose on the leaf pigments, growth, and rooting of blackberry (*Rubus fruticosus*) microshoots. *Agronomy* **2023**, *13*, 89. <https://doi.org/10.3390/agronomy13010089>.
- Díaz-Rueda, P.; Cantos-Barragán, M.; Colmenero-Flores, J. M. Growth quality and development of olive plants cultured in vitro under different illumination regimes. *Plants* **2021**, *10*. <https://doi.org/10.3390/plants10102214>.
- Díaz-Sala, C. Molecular dissection of the regenerative capacity of forest tree species: Special focus on conifers. *Front. Plant Sci.* **2019**, *9*. <https://doi.org/10.3389/fpls.2018.01943>.
- Franco-Navarro, J.D.; Rosales, M.A.; Álvarez, R.; Cubero-Font, P.; Calvo, P.; Díaz-Espejo, A.; Colmenero-Flores, J.M. Chloride as macronutrient increases water use efficiency by anatomically-driven reduced stomatal conductance and increased mesophyll diffusion to CO₂. *Plant J.* **2019**, *99*, 815–831. <https://doi.org/10.1111/tpj.14423>.
- Gil, C. S.; Jung, H. Y.; Lee, C.; Eom, S. H. Blue light and NAA treatment significantly improve rooting on single leaf-bud cutting of *Chrysanthemum* via upregulated rooting-related genes. *Sci Hortic.* **2020**, *274*. <https://doi.org/10.1016/j.scienta.2020.109650>.
- Giri, C. C.; Shyamkumar, B.; Anjaneyulu, C. Progress in tissue culture, genetic transformation and applications of biotechnology to trees: An overview. *Trees* **2004**, *18*, 115–135. <https://doi.org/10.1007/s00468-003-0287-6>.

- Gnasekaran, P.; Rahman, Z. A.; Chew, B. L.; Appalasamy, S.; Mariappan, V.; Subramaniam, S. Development of micropropagation system of *Zingiber officinale* var. *rubrum* Theilade using different spectrum light-emitting diode (LED) irradiation. *Ind. Crops Prod.* **2021**, *170*. <https://doi.org/10.1016/j.indcrop.2021.113748>.
- Guerra, F.; Badilla, L.; Cautin, R.; Castro, M. Micropropagation of *Citronella mucronata* D. Don, a Vulnerable Chilean Endemic Tree Species. *Plants* **2022**, *11*. <https://doi.org/10.3390/plants11182425>.
- Gupta, S.; Jatothu, B. Fundamentals and applications of light-emitting diodes (LEDs) in in vitro plant growth and morphogenesis. *Plant Biotechnol. Rep.* **2013**, *7*, 211–220. <https://doi.org/10.1007/s11816-013-0277-0>.
- Halmagyi, A.; Kevresan, S.; Kovacevic, B.; Orlovic, S.; Miladinovic, D.; Cirin-Novta, V.; Kuhajda, K. Effects of naphthenic acids on rooting of in vitro grown *Sequoia sempervirens*. *Acta Hortic.* **2010**, *885*, 139–144. <https://doi.org/10.17660/actahortic.2010.885.18>.
- Joshi, V.; Joung, J. G.; Fei, Z.; Jander, G. Interdependence of threonine, methionine and isoleucine metabolism in plants: Accumulation and transcriptional regulation under abiotic stress. *Amino acids* **2010**, *39*, 933–947. <https://doi.org/10.1007/s00726-010-0505-7>.
- Khamushi, M.; Dehestani-Ardakani, M.; Zarei, A.; Aliabad, K.K. An efficient protocol for micropropagation of old cypress of Abarkuh (*Cupressus sempervirens* var. *horizontalis* [Mill.]) under in vitro condition. *Plant Cell Tissue Organ Cult.* **2019**, *138*, 597–601. <https://doi.org/10.1007/s11240-019-01645-z>.
- Lai, C. S.; Kho, Y. H.; Chew, B. L.; Raja, P. B.; Subramaniam, S. Organogenesis of *Cucumis metuliferus* plantlets under the effects of LEDs and silver nanoparticles. *S. Afr. J. Bot.* **2022**, *148*, 78–87. <https://doi.org/10.1016/j.sajb.2022.04.020>.
- Lotfi, M.; Mars, M.; Werbrouck, S. Optimizing pear micropropagation and rooting with light emitting diodes and trans-cinnamic acid. *Plant Growth Regul.* **2019**, *88*, 173–180. <https://doi.org/10.1007/s10725-019-00498-y>.
- Mingozi, M.; Morini, S.; Lucchesini, M.; Mensuali-Sodi, A. Effects of leaf soluble sugars content and net photosynthetic rate of quince donor shoots on subsequent morphogenesis in leaf explants. *Biol. Plant.* **2011**, *55*, 237–242. <https://doi.org/10.1007/s10535-011-0034-6>.
- Montalbán, I. A.; Novák, O.; Rolčík, J.; Strnad, M.; Moncaleán, P. Endogenous cytokinin and auxin profiles during in vitro organogenesis from vegetative buds of *Pinus radiata* adult trees. *Physiol. Plant.* **2013**, *148*, 214–231. <https://doi.org/10.1111/j.1399-3054.2012.01709.x>.
- Moradi, P.; Ford-Lloyd, B.; Pritchard, J. Metabolomic approach reveals the biochemical mechanisms underlying drought stress tolerance in thyme. *Anal. Biochem.* **2017**, *527*, 49–62. <https://doi.org/10.1016/j.ab.2017.02.006>.
- Muthuramalingam, P.; Krishnan, S. R.; Pandian, S.; Mareeswaran, N.; Aruni, W.; Pandian, S. K.; Ramesh, M. Global analysis of threonine metabolism genes unravel key players in rice to improve the abiotic stress tolerance. *Sci. Rep.* **2018**, *8*, 9270. <https://doi.org/10.1038/s41598-018-27703-8>.
- Nicholson, J.; Shukla, M. R.; Saxena, P. K. In vitro rooting of hybrid hazelnuts (*Corylus avellana* × *Corylus americana*) in a temporary immersion system. *Botany* **2020**, *98*, 343–352. <https://doi.org/10.1139/cjb-2019-0206>.

- Niemi, K.; Scagel, C.; Häggman, H. Application of ectomycorrhizal fungi in vegetative propagation of conifers. *Plant Cell Tissue Organ Cult.* **2004**, *78*, 83–91. <https://doi.org/10.1023/B:TICU.0000020379.52514.72>.
- Pang, W. Q.; Tan, S. T.; Mad'Atari, M. F.; Yoong, I. C. K.; Subramaniam, S. Establishment of an efficient micropropagation protocol for cameron highlands White Strawberry (*Fragaria x ananassa*) using light emitting diodes (LEDs) system. *S. Afri. J. Bot.* **2023**, *157*, 189–200. <https://doi.org/10.1016/j.sajb.2023.03.061>.
- Pereira, C.; Montalbán, I.A.; Pedrosa, A.; Tavares, J.; Pestryakov, A.; Bogdanchikova, N.; Canhoto, J.; Moncaleán, P. Regeneration of *Pinus halepensis* (Mill.) through organogenesis from apical shoot buds. *Forests* **2021**, *12*, 363. <https://doi.org/10.3390/f12030363>.
- Phillips, G.C.; Garda, M. Plant tissue culture media and practices: An overview. *In Vitro Cell. Dev. Biol. Plant.* **2019**, *55*, 242–257. <https://doi.org/10.1007/s11627-019-09983-5>.
- Rodríguez, S.M.; Ordás, R.J.; Alvarez, J.M. Conifer Biotechnology: An Overview. *Forests* **2022**, *13*, 1061. <https://doi.org/10.3390/f13071061>.
- San José, M. C.; Blázquez, N.; Cernadas, M. J.; Janeiro, L. V.; Cuenca, B.; Sánchez, C.; Vidal, N. Temporary immersion systems to improve alder micropropagation. *Plant Cell Tissue Organ Cult.* **2020**, *143*, 265–275. <https://doi.org/10.1007/s11240-020-01937-9>.
- Santana Costa Souza, D. M.; Fernandes, S. B.; Oliveira Silva, E.; Politi Duarte, V.; Santos Gonçalves, D.; de Carvalho, D.; Teixeira, G.; Ebling Brondani, G et al. Effect of light intensity on in vitro introduction and multiplication of *Eucalyptus grandis* × *Eucalyptus urophylla*. *In Vitro Cell. Dev. Biol.* **2022**, *58*, 225–239. <https://doi.org/10.1007/s11627-021-10237-6>.
- Sarmast, M. K. In vitro propagation of conifers using mature shoots. *J For Res.* **2018**, *29*, 565–574. <https://doi.org/10.1007/s11676-018-0608-7>.
- Sarrobot, C.; Thibaud, M.C.; Contard-David, P.; Gineste, S.; Bechtold, N.; Robaglia, C.; Nussaume, L. Identification of an *Arabidopsis thaliana* mutant accumulating threonine resulting from mutation in a new dihydrodipicolinate synthase gene. *Plant J.* **2000**, *24*, 357–367. <https://doi.org/10.1046/j.1365-313X.2000.00884.x>.
- Shekhawat, M. S.; Manokari, M.; Kannan, N. Micromorphological response towards altered environmental conditions in subsequent stages of in vitro propagation of *Morinda coreia*. *Environ. Exp. Biol.* **2017**, *15*, 37–46. <https://doi.org/10.22364/eeb.15.06>.
- Silva, N.C.B.; Macedo, A.F.; Lage, C.L.S.; Esquibel, M.A.; Sato, A. Developmental effects of additional ultraviolet a radiation, growth regulators and tyrosine in *Alternanthera brasiliana* (L.) Kuntze cultured in vitro. *Braz. Arch. Biol. Technol.* **2005**, *48*, 779–786. <https://doi.org/10.1590/S1516-89132005000600014>.
- Von Aderkas, P.; Bonga, J.M. Influencing micropropagation and somatic embryogenesis in mature trees by manipulation of phase change, stress and culture environment. *Tree Physiol.* **2000**, *20*, 921–928. <https://doi.org/10.1093/treephys/20.14.921>.
- Wu, G. Amino acids: Metabolism, functions, and nutrition. *Amino acids* **2009**, *37*, 1–17. <https://doi.org/10.1007/s00726-009-0269-0>.

Wu, H.C.; Lin, C.C. Red light-emitting diode light irradiation improves root and leaf formation in difficult-to-propagate *Protea cynaroides* L. plantlets in vitro. *HortScience* **2012**, *47*, 1490–1494. <https://doi.org/10.21273/HORTSCI.47.10.1490>.

Yaseen, M.; Ahmad, T.; Sablok, G.; Standardi, A.; Hafiz, I. A. Review: Role of carbon sources for in vitro plant growth and development. *Mol. Biol. Rep.* **2013**, *40*, 2837–2849. <https://doi.org/10.1007/s11033-012-2299-z>.

Zarei, M.; Salehi, H.; Jowkar, A. Controlling the barriers of cloning mature *Picea abies* (L.) H. Karst. via tissue culture and co-cultivation with *Agrobacterium rhizogenes*. *Trees* **2020**, *34*, 637–647. <https://doi.org/10.1007/s00468-019-01945-z>.

CONCLUSIONS

CONCLUSIONS

1. The collection date and the interaction between genotype and BA concentration have a significant effect on the percentage of explants forming shoots of adult *Pinus radiata* trees organogenesis. Shoots cultured in rooting medium supplemented with 3% sucrose and exposed to white FL show a significant higher root induction response.
2. The profile of carbohydrates and amino acids in *in vitro* shoots of *Pinus radiata* is influenced by the rooting medium. Fructose, glucose and sucrose contents increase in shoots during root induction stage. Significantly higher threonine and tyrosine levels are present in shoots after root expression stage.
3. *Cryptomeria japonica* basal explants from adult trees longer than 1.5 cm produce a significantly higher number of shoots than the rest of explants evaluated. QL medium supplemented with BA is the best chemical environment for the success of organogenic response.
4. *Cryptomeria japonica* explants induced with m-T and rooted with 1.5% sucrose display the best results in terms of root induction. Then, plantlets exposed to red LEDs show a significantly higher number of roots.
5. *Pinus ponderosa* whole zygotic embryos are the best initial explants to obtain developed shoots. *Pinus ponderosa* explants growing under blue LEDs produce a significantly higher number of shoots with rooting capacity, while plantlets grown under FL light during *in vitro* culture increase the acclimatization success.
6. LEDs show an effect in *Pinus ponderosa* needles decreasing the number of stomata in relation to FL.
7. *Sequoia sempervirens* needs different chemical environment in the *in vitro* stage depending on the objective of the process (improve the rooting or acclimatization stage).

SUPPLEMENTARY MATERIALS

CHAPTER 3

Influence of Physico-Chemical Factors on the Efficiency and

Metabolite Profile of Adult *Pinus radiata* D. Don Bud

Organogenesis

SUPPLEMENTARY MATERIALS CHAPTER 3

Table S1. Statistical analysis for explants forming shoots in different collection dates from *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)].

Source	df	Explants forming shoots (%)	
		χ^2 Test	p-Value
Collection dates	5	776.71	$\leq 0.05^*$

*Significant differences at $p < 0.05$, df Degrees of freedom.

Table S2. Statistical analysis for explants forming shoots from nine different genotypes of *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with 6-benzyladenine (BA) (22 and 44 μ M).

Source	df	Explants forming shoots (%)	
		χ^2 Test	p-Value
Genotype (G)	8	795.04	$\leq 0.05^*$
BA concentrations (B)	1	794.21	$> 0.05^{n.s}$
G x B	8	745.98	$\leq 0.05^*$

*Significant differences at $p < 0.05$, $^{n.s}$ Non-significant at $p < 0.05$, df Degrees of freedom.

Table S3. ANOVA for number of shoots formed per explant from *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with 6-benzyladenine (BA) (22 and 44 μ M).

Source	df	N° shoots/explant	
		F Test	p-Value
BA concentrations	1	0.06	$> 0.05^{n.s}$

$^{n.s}$ Non-significant at $p < 0.05$, df Degrees of freedom.

Table S4. Statistical analysis for root induction, number roots per explant and length of longest root from *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] according to 6-benzyladenine (BA) concentrations and light treatments applied in the root induction phase.

Source	df	Root induction		N° roots/explant		Length of the longest root	
		χ^2 Test	p-Value	F Test	p-Value	F Test	p-Value
BA concentrations (B)	1	251.24	$> 0.05^{n.s}$	0.83	$> 0.05^{n.s}$	0.31	$> 0.05^{n.s}$
Light treatments (L)	1	243.56	$\leq 0.05^*$	2.94	$> 0.05^{n.s}$	2.27	$> 0.05^{n.s}$
B x L	1	242.87	$> 0.05^{n.s}$	0.75	$> 0.05^{n.s}$	16.60	$\leq 0.05^*$

*Significant differences at $p < 0.05$, $^{n.s}$ Non-significant at $p < 0.05$, df Degrees of freedom.

Table S5. Statistical analysis for root induction (%), number roots per explant, and length of longest root from *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] according to 6-benzyladenine (BA) and sucrose concentrations applied in the root induction phase.

Source	df	Root induction		N° roots/explant		Length of the longest root	
		X ² Test	p-Value	F Test	p-Value	F Test	p-Value
BA concentrations (B)	1	251.24	>0.05 ^{n.s}	0.72	>0.05 ^{n.s}	0.25	>0.05 ^{n.s}
Sucrose concentrations (S)	1	242.33	≤0.05*	0.06	>0.05 ^{n.s}	0.95	>0.05 ^{n.s}
B x S	1	242.10	>0.05 ^{n.s}	0.32	>0.05 ^{n.s}	5.43	≤0.05*

*Significant differences at $p < 0.05$, ^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S6. Statistical analysis for root induction (%), number roots per explant, and length of longest root from *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] according to light treatments and sucrose concentrations applied in the root induction phase.

Source	df	Root induction		N° roots/explant		Length of the longest root	
		X ² Test	p-Value	F Test	p-Value	F Test	p-Value
Light treatment (L)	1	236.52	≤0.05*	4.82	≤0.05*	0.99	>0.05 ^{n.s}
Sucrose concentrations (S)	1	243.38	≤0.05*	0.11	>0.05 ^{n.s}	0.86	>0.05 ^{n.s}
L x S	1	232.75	>0.05 ^{n.s}	0.92	>0.05 ^{n.s}	3.28	>0.05 ^{n.s}

*Significant differences at $p < 0.05$, ^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S7. ANOVA for carbohydrates content of *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with a mixture of 5 μM 1-Naphthaleneacetic acid (NAA) and 10 μM indole-3-butyric acid (IBA) according to 6-benzyladenine (BA) concentrations and light treatments.

Carbohydrates	df	Source					
		BA concentrations (B)		Light treatments (L)		B x L	
		F Test	p-Value	F Test	p-Value	F Test	p-Value
Fructose	1	0.04	>0.05 ^{n.s}	0.90	>0.05 ^{n.s}	0.002	>0.05 ^{n.s}
Galactose	1	1.41	>0.05 ^{n.s}	0.83	>0.05 ^{n.s}	2.65	>0.05 ^{n.s}
Glucose	1	0.16	>0.05 ^{n.s}	0.94	>0.05 ^{n.s}	0.14	>0.05 ^{n.s}
Mannitol	1	0.66	>0.05 ^{n.s}	0.04	>0.05 ^{n.s}	0.71	>0.05 ^{n.s}
Sucrose	1	1.99	>0.05 ^{n.s}	1.19	>0.05 ^{n.s}	0.009	>0.05 ^{n.s}

^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S8. ANOVA for carbohydrates content of *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with a mixture of 5 μM 1-Naphthaleneacetic acid (NAA) and 10 μM indole-3-butyric acid (IBA) according to 6-benzyladenine (BA) and sucrose concentrations.

Carbohydrates	df	Source					
		BA concentrations (S)		Sucrose concentrations (B)		B x S	
		F Test	p-Value	F Test	p-Value	F Test	p-Value
Fructose	1	0.07	>0.05 ^{n.s}	14.04	≤0.05*	0.01	>0.05 ^{n.s}
Galactose	1	1.39	>0.05 ^{n.s}	1.90	>0.05 ^{n.s}	1.24	>0.05 ^{n.s}
Glucose	1	0.25	>0.05 ^{n.s}	13.73	≤0.05*	0.01	>0.05 ^{n.s}
Mannitol	1	0.76	>0.05 ^{n.s}	3.61	>0.05 ^{n.s}	0.20	>0.05 ^{n.s}

*Significant differences at $p < 0.05$, ^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S9. ANOVA for carbohydrates content of *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with a mixture of 5 μM 1-Naphthaleneacetic acid (NAA) and 10 μM indole-3-butyric acid (IBA) according to light treatments and sucrose concentrations.

Carbohydrates	df	Source					
		Light treatment (L)		Sucrose concentrations (S)		L x S	
		F Test	p-Value	F Test	p-Value	F Test	p-Value
Fructose	1	1.39	>0.05 ^{n.s}	13.71	$\leq 0.05^*$	0.59	>0.05 ^{n.s}
Galactose	1	0.54	>0.05 ^{n.s}	2.24	>0.05 ^{n.s}	0.97	>0.05 ^{n.s}
Glucose	1	1.31	>0.05 ^{n.s}	12.86	$\leq 0.05^*$	0.06	>0.05 ^{n.s}
Mannitol	1	0.10	>0.05 ^{n.s}	3.49	>0.05 ^{n.s}	0.08	>0.05 ^{n.s}
Sucrose	1	3.37	>0.05 ^{n.s}	9.83	$\leq 0.05^*$	2.50	>0.05 ^{n.s}

*Significant differences at $p < 0.05$, ^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S10. ANOVA for carbohydrates content of *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with 2 g L^{-1} activated charcoal, according to 6-benzyladenine (BA) concentrations and light treatments.

Carbohydrates	df	Source					
		BA concentrations (B)		Light treatments(L)		B x L	
		F Test	p-Value	F Test	p-Value	F Test	p-Value
Fructose	1	0.23	>0.05 ^{n.s}	0.51	>0.05 ^{n.s}	0.68	>0.05 ^{n.s}
Galactose	1	0.64	>0.05 ^{n.s}	1.09	>0.05 ^{n.s}	0.22	>0.05 ^{n.s}
Glucose	1	1.22	>0.05 ^{n.s}	0.95	>0.05 ^{n.s}	1.28	>0.05 ^{n.s}
Mannitol	1	0.43	>0.05 ^{n.s}	0.00	>0.05 ^{n.s}	2.96	>0.05 ^{n.s}
Sucrose	1	2.19	>0.05 ^{n.s}	0.82	>0.05 ^{n.s}	0.88	>0.05 ^{n.s}

^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S11. ANOVA for carbohydrates content of *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with 2 g L^{-1} activated charcoal, according to 6-benzyladenine (BA) and sucrose concentrations.

Carbohydrates	df	Source					
		BA concentrations (B)		Sucrose concentrations (S)		B x S	
		F Test	p-Value	F Test	p-Value	F Test	p-Value
Fructose	1	0.24	>0.05 ^{n.s}	1.71	>0.05 ^{n.s}	0.005	>0.05 ^{n.s}
Galactose	1	0.61	>0.05 ^{n.s}	0.16	>0.05 ^{n.s}	0.10	>0.05 ^{n.s}
Glucose	1	1.17	>0.05 ^{n.s}	0.56	>0.05 ^{n.s}	0.80	>0.05 ^{n.s}
Mannitol	1	0.39	>0.05 ^{n.s}	0.28	>0.05 ^{n.s}	0.64	>0.05 ^{n.s}
Sucrose	1	2.16	>0.05 ^{n.s}	0.83	>0.05 ^{n.s}	0.56	>0.05 ^{n.s}

^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S12. ANOVA for carbohydrates content of *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with 2 gL⁻¹ activated charcoal, according to light treatments and sucrose concentrations.

Carbohydrates	df	Source					
		Light treatments (L)		Sucrose concentrations (S)		L x C	
		F Test	p-Value	F Test	p-Value	F Test	p-Value
Fructose	1	0.65	>0.05 ^{n.s}	2.26	>0.05 ^{n.s}	0.006	>0.05 ^{n.s}
Galactose	1	0.79	>0.05 ^{n.s}	0.13	>0.05 ^{n.s}	0.01	>0.05 ^{n.s}
Glucose	1	1.23	>0.05 ^{n.s}	1.01	>0.05 ^{n.s}	0.41	>0.05 ^{n.s}
Mannitol	1	0.006	>0.05 ^{n.s}	0.20	>0.05 ^{n.s}	0.19	>0.05 ^{n.s}
Sucrose	1	1.27	>0.05 ^{n.s}	1.48	>0.05 ^{n.s}	1.36	>0.05 ^{n.s}

^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S13. ANOVA for amino acid content of *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with a mixture of 5 μ M 1-Naphthaleneacetic acid (NAA) and 10 μ M indole-3-butyric acid (IBA) according to 6-benzyladenine (BA) concentrations and light treatments.

Free amino acid	df	Source					
		BA concentrations (B)		Light treatments (L)		B x L	
		F Test	p-Value	F Test	p-Value	F Test	p-Value
Alanine	1	0.09	>0.05 ^{n.s}	0.19	>0.05 ^{n.s}	0.32	>0.05 ^{n.s}
Arginine	1	0.01	>0.05 ^{n.s}	0.81	>0.05 ^{n.s}	0.21	>0.05 ^{n.s}
Asparagine	1	0.81	>0.05 ^{n.s}	0.51	>0.05 ^{n.s}	1.50	>0.05 ^{n.s}
Aspartic acid	1	0.10	>0.05 ^{n.s}	2.36	>0.05 ^{n.s}	0.96	>0.05 ^{n.s}
Cystine	1	0.91	>0.05 ^{n.s}	2.34	>0.05 ^{n.s}	0.90	>0.05 ^{n.s}
Glutamic acid	1	0.28	>0.05 ^{n.s}	2.10	>0.05 ^{n.s}	0.15	>0.05 ^{n.s}
Glutamine	1	0.97	>0.05 ^{n.s}	0.11	>0.05 ^{n.s}	0.16	>0.05 ^{n.s}
Glycine	1	0.006	>0.05 ^{n.s}	0.86	>0.05 ^{n.s}	0.17	>0.05 ^{n.s}
Hydroxyproline	1	0.35	>0.05 ^{n.s}	0.27	>0.05 ^{n.s}	1.02	>0.05 ^{n.s}
Histidine	1	0.56	>0.05 ^{n.s}	0.24	>0.05 ^{n.s}	0.03	>0.05 ^{n.s}
Isoleucine	1	1.36	>0.05 ^{n.s}	0.02	>0.05 ^{n.s}	0.35	>0.05 ^{n.s}
Leucine	1	1.57	>0.05 ^{n.s}	0.06	>0.05 ^{n.s}	0.18	>0.05 ^{n.s}
Lysine	1	0.40	>0.05 ^{n.s}	0.001	>0.05 ^{n.s}	0.08	>0.05 ^{n.s}
Methionine	1	1.05	>0.05 ^{n.s}	0.10	>0.05 ^{n.s}	1.52	>0.05 ^{n.s}
Phenylalanine	1	2.99	>0.05 ^{n.s}	0.013	>0.05 ^{n.s}	1.35	>0.05 ^{n.s}
Proline	1	0.06	>0.05 ^{n.s}	0.82	>0.05 ^{n.s}	0.01	>0.05 ^{n.s}
Serine	1	1.28	>0.05 ^{n.s}	0.74	>0.05 ^{n.s}	1.04	>0.05 ^{n.s}
Threonine	1	1.64	>0.05 ^{n.s}	1.25	>0.05 ^{n.s}	0.34	>0.05 ^{n.s}
Tryptophan	1	1.14	>0.05 ^{n.s}	0.13	>0.05 ^{n.s}	0.22	>0.05 ^{n.s}
Tyrosine	1	3.53	>0.05 ^{n.s}	0.002	>0.05 ^{n.s}	0.72	>0.05 ^{n.s}
Valine	1	2.23	>0.05 ^{n.s}	0.67	>0.05 ^{n.s}	0.27	>0.05 ^{n.s}

^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S14. ANOVA for amino acid content of *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with a mixture of 5 μ M 1-Naphthaleneacetic acid (NAA) and 10 μ M indole-3-butyric acid (IBA) according to 6-benzyladenine (BA) and sucrose concentrations.

Free amino acid	df	Source					
		BA concentrations (B)		Sucrose concentrations (S)		B x S	
		F Test	p-Value	F Test	p-Value	F Test	p-Value
Alanine	1	0.09	>0.05 ^{n.s}	1.23	>0.05 ^{n.s}	0.77	>0.05 ^{n.s}
Arginine	1	0.02	>0.05 ^{n.s}	1.20	>0.05 ^{n.s}	2.03	>0.05 ^{n.s}
Asparagine	1	0.84	>0.05 ^{n.s}	2.93	>0.05 ^{n.s}	0.03	>0.05 ^{n.s}
Aspartic acid	1	0.08	>0.05 ^{n.s}	0.001	>0.05 ^{n.s}	0.00	>0.05 ^{n.s}
Cystine	1	0.82	>0.05 ^{n.s}	0.01	>0.05 ^{n.s}	0.69	>0.05 ^{n.s}
Glutamic acid	1	0.27	>0.05 ^{n.s}	0.95	>0.05 ^{n.s}	1.09	>0.05 ^{n.s}
Glutamine	1	0.98	>0.05 ^{n.s}	0.00	>0.05 ^{n.s}	0.52	>0.05 ^{n.s}
Glycine	1	0.006	>0.05 ^{n.s}	1.38	>0.05 ^{n.s}	0.45	>0.05 ^{n.s}
Hydroxyproline	1	0.35	>0.05 ^{n.s}	0.001	>0.05 ^{n.s}	1.31	>0.05 ^{n.s}
Histidine	1	0.57	>0.05 ^{n.s}	0.10	>0.05 ^{n.s}	0.63	>0.05 ^{n.s}
Isoleucine	1	1.42	>0.05 ^{n.s}	0.23	>0.05 ^{n.s}	1.12	>0.05 ^{n.s}
Leucine	1	1.67	>0.05 ^{n.s}	0.68	>0.05 ^{n.s}	1.01	>0.05 ^{n.s}
Lysine	1	0.46	>0.05 ^{n.s}	0.70	>0.05 ^{n.s}	2.85	>0.05 ^{n.s}
Methionine	1	0.94	>0.05 ^{n.s}	0.13	>0.05 ^{n.s}	0.01	>0.05 ^{n.s}
Phenylalanine	1	3.22	>0.05 ^{n.s}	0.16	>0.05 ^{n.s}	2.95	>0.05 ^{n.s}
Proline	1	0.06	>0.05 ^{n.s}	0.28	>0.05 ^{n.s}	0.58	>0.05 ^{n.s}
Serine	1	1.32	>0.05 ^{n.s}	0.97	>0.05 ^{n.s}	1.60	>0.05 ^{n.s}
Threonine	1	1.59	>0.05 ^{n.s}	0.40	>0.05 ^{n.s}	0.47	>0.05 ^{n.s}
Tryptophan	1	1.18	>0.05 ^{n.s}	0.04	>0.05 ^{n.s}	1.13	>0.05 ^{n.s}
Tyrosine	1	3.73	>0.05 ^{n.s}	0.00	>0.05 ^{n.s}	2.04	>0.05 ^{n.s}
Valine	1	2.26	>0.05 ^{n.s}	1.04	>0.05 ^{n.s}	0.28	>0.05 ^{n.s}

^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S15. ANOVA for amino acid content of *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with a mixture of 5 μ M 1-Naphthaleneacetic acid (NAA) and 10 μ M indole-3-butyric acid (IBA) according to light treatments and sucrose concentrations.

Free amino acid	df	Source					
		Light treatments (L)		Sucrose concentrations		L x S	
		F Test	p-Value	F Test	p-Value	F Test	p-Value
Alanine	1	0.18	>0.05 ^{n.s}	1.41	>0.05 ^{n.s}	0.50	>0.05 ^{n.s}
Arginine	1	0.85	>0.05 ^{n.s}	0.86	>0.05 ^{n.s}	0.00	>0.05 ^{n.s}
Asparagine	1	0.68	>0.05 ^{n.s}	2.62	>0.05 ^{n.s}	0.99	>0.05 ^{n.s}
Aspartic acid	1	2.14	>0.05 ^{n.s}	0.06	>0.05 ^{n.s}	0.01	>0.05 ^{n.s}
Cystine	1	1.86	>0.05 ^{n.s}	0.007	>0.05 ^{n.s}	0.00	>0.05 ^{n.s}
Glutamic acid	1	2.31	>0.05 ^{n.s}	0.60	>0.05 ^{n.s}	0.53	>0.05 ^{n.s}
Glutamine	1	0.05	>0.05 ^{n.s}	0.002	>0.05 ^{n.s}	0.38	>0.05 ^{n.s}
Glycine	1	0.89	>0.05 ^{n.s}	1.11	>0.05 ^{n.s}	0.42	>0.05 ^{n.s}
Hydroxyproline	1	0.34	>0.05 ^{n.s}	0.01	>0.05 ^{n.s}	1.91	>0.05 ^{n.s}
Histidine	1	0.17	>0.05 ^{n.s}	0.14	>0.05 ^{n.s}	0.000	>0.05 ^{n.s}
Isoleucine	1	0.06	>0.05 ^{n.s}	0.18	>0.05 ^{n.s}	0.77	>0.05 ^{n.s}
Leucine	1	0.01	>0.05 ^{n.s}	0.60	>0.05 ^{n.s}	0.54	>0.05 ^{n.s}
Lysine	1	0.009	>0.05 ^{n.s}	0.60	>0.05 ^{n.s}	0.000	>0.05 ^{n.s}
Methionine	1	0.15	>0.05 ^{n.s}	0.18	>0.05 ^{n.s}	0.15	>0.05 ^{n.s}
Phenylalanine	1	0.003	>0.05 ^{n.s}	0.14	>0.05 ^{n.s}	0.000	>0.05 ^{n.s}
Proline	1	0.82	>0.05 ^{n.s}	0.17	>0.05 ^{n.s}	1.30	>0.05 ^{n.s}
Serine	1	0.91	>0.05 ^{n.s}	0.67	>0.05 ^{n.s}	0.74	>0.05 ^{n.s}
Threonine	1	0.93	>0.05 ^{n.s}	0.23	>0.05 ^{n.s}	1.002	>0.05 ^{n.s}
Tryptophan	1	0.21	>0.05 ^{n.s}	0.08	>0.05 ^{n.s}	0.22	>0.05 ^{n.s}
Tyrosine	1	0.01	>0.05 ^{n.s}	0.00	>0.05 ^{n.s}	0.83	>0.05 ^{n.s}
Valine	1	0.43	>0.05 ^{n.s}	0.82	>0.05 ^{n.s}	1.13	>0.05 ^{n.s}

^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S16. ANOVA amino acid content of *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with 2 gL⁻¹ activated charcoal according to 6-benzyladenine (BA) concentrations and light treatments.

Free amino acid	df	Source					
		BA concentrations (B)		Light treatments (L)		B x L	
		F Test	p-Value	F Test	p-Value	F Test	p-Value
Alanine	1	1.45	>0.05 ^{n.s}	0.09	>0.05 ^{n.s}	1.36	>0.05 ^{n.s}
Arginine	1	0.01	>0.05 ^{n.s}	1.31	>0.05 ^{n.s}	1.64	>0.05 ^{n.s}
Asparagine	1	0.48	>0.05 ^{n.s}	0.82	>0.05 ^{n.s}	0.44	>0.05 ^{n.s}
Aspartic acid	1	0.21	>0.05 ^{n.s}	0.21	>0.05 ^{n.s}	0.004	>0.05 ^{n.s}
Cystine	1	0.11	>0.05 ^{n.s}	0.04	>0.05 ^{n.s}	0.54	>0.05 ^{n.s}
Glutamic acid	1	0.49	>0.05 ^{n.s}	0.50	>0.05 ^{n.s}	4.10	>0.05 ^{n.s}
Glutamine	1	1.04	>0.05 ^{n.s}	0.49	>0.05 ^{n.s}	0.30	>0.05 ^{n.s}
Glycine	1	0.29	>0.05 ^{n.s}	0.04	>0.05 ^{n.s}	0.01	>0.05 ^{n.s}
Hydroxyproline	1	2.66	>0.05 ^{n.s}	0.05	>0.05 ^{n.s}	0.03	>0.05 ^{n.s}
Histidine	1	0.03	>0.05 ^{n.s}	0.02	>0.05 ^{n.s}	1.12	>0.05 ^{n.s}
Isoleucine	1	0.01	>0.05 ^{n.s}	0.97	>0.05 ^{n.s}	0.006	>0.05 ^{n.s}
Leucine	1	0.10	>0.05 ^{n.s}	0.66	>0.05 ^{n.s}	0.02	>0.05 ^{n.s}
Lysine	1	1.55	>0.05 ^{n.s}	0.53	>0.05 ^{n.s}	1.38	>0.05 ^{n.s}
Methionine	1	0.77	>0.05 ^{n.s}	0.18	>0.05 ^{n.s}	0.006	>0.05 ^{n.s}
Phenylalanine	1	0.46	>0.05 ^{n.s}	1.43	>0.05 ^{n.s}	0.01	>0.05 ^{n.s}
Proline	1	0.001	>0.05 ^{n.s}	0.31	>0.05 ^{n.s}	0.006	>0.05 ^{n.s}
Serine	1	0.01	>0.05 ^{n.s}	1.03	>0.05 ^{n.s}	0.01	>0.05 ^{n.s}
Threonine	1	0.23	>0.05 ^{n.s}	0.02	>0.05 ^{n.s}	0.06	>0.05 ^{n.s}
Tryptophan	1	2.06	>0.05 ^{n.s}	0.47	>0.05 ^{n.s}	0.18	>0.05 ^{n.s}
Tyrosine	1	0.10	>0.05 ^{n.s}	0.24	>0.05 ^{n.s}	1.21	>0.05 ^{n.s}
Valine	1	0.46	>0.05 ^{n.s}	0.37	>0.05 ^{n.s}	0.15	>0.05 ^{n.s}

^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S17. ANOVA for amino acid content of *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with 2 gL⁻¹ activated charcoal according to 6-benzyladenine (BA) and sucrose concentrations.

Free amino acid	df	Source					
		BA concentrations (B)		Sucrose concentrations (S)		B x S	
		F Test	p-Value	F Test	p-Value	F Test	p-Value
Alanine	1	1.52	>0.05 ^{n.s}	1.37	>0.05 ^{n.s}	0.98	>0.05 ^{n.s}
Arginine	1	0.01	>0.05 ^{n.s}	0.08	>0.05 ^{n.s}	2.95	>0.05 ^{n.s}
Asparagine	1	0.48	>0.05 ^{n.s}	0.88	>0.05 ^{n.s}	0.43	>0.05 ^{n.s}
Aspartic acid	1	0.22	>0.05 ^{n.s}	0.99	>0.05 ^{n.s}	0.12	>0.05 ^{n.s}
Cystine	1	0.11	>0.05 ^{n.s}	1.53	>0.05 ^{n.s}	0.29	>0.05 ^{n.s}
Glutamic acid	1	0.44	>0.05 ^{n.s}	2.03	>0.05 ^{n.s}	0.47	>0.05 ^{n.s}
Glutamine	1	1.01	>0.05 ^{n.s}	0.11	>0.05 ^{n.s}	0.001	>0.05 ^{n.s}
Glycine	1	0.30	>0.05 ^{n.s}	0.76	>0.05 ^{n.s}	0.17	>0.05 ^{n.s}
Hydroxyproline	1	5.28	≤0.05*	2.38	>0.05 ^{n.s}	9.10	≤0.05*
Histidine	1	0.04	>0.05 ^{n.s}	2.22	>0.05 ^{n.s}	0.41	>0.05 ^{n.s}
Isoleucine	1	0.01	>0.05 ^{n.s}	2.50	>0.05 ^{n.s}	0.08	>0.05 ^{n.s}
Leucine	1	0.10	>0.05 ^{n.s}	0.82	>0.05 ^{n.s}	0.02	>0.05 ^{n.s}
Lysine	1	1.62	>0.05 ^{n.s}	0.05	>0.05 ^{n.s}	2.86	>0.05 ^{n.s}
Methionine	1	0.77	>0.05 ^{n.s}	0.06	>0.05 ^{n.s}	0.03	>0.05 ^{n.s}
Phenylalanine	1	0.44	>0.05 ^{n.s}	0.16	>0.05 ^{n.s}	0.74	>0.05 ^{n.s}
Proline	1	0.002	>0.05 ^{n.s}	3.03	>0.05 ^{n.s}	0.59	>0.05 ^{n.s}
Serine	1	0.01	>0.05 ^{n.s}	3.67	>0.05 ^{n.s}	0.11	>0.05 ^{n.s}
Threonine	1	0.31	*>0.05 ^{n.s}	6.09	≤0.05*	0.34	>0.05 ^{n.s}
Tryptophan	1	2.30	>0.05 ^{n.s}	0.16	>0.05 ^{n.s}	2.80	>0.05 ^{n.s}
Tyrosine	1	0.13	>0.05 ^{n.s}	7.86	≤0.05*	0.75	>0.05 ^{n.s}
Valine	1	0.48	>0.05 ^{n.s}	1.10	>0.05 ^{n.s}	0.19	>0.05 ^{n.s}

*Significant differences at $p < 0.05$, ^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S18. ANOVA amino acid content of *Pinus radiata* D. Don cultured in LP medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with 2 gL⁻¹ activated charcoal according to light treatments and sucrose concentrations.

Free amino acid	df	Source					
		Light treatments (L)		Sucrose concentrations (S)		L x S	
		F Test	p-Value	F Test	p-Value	F Test	p-Value
Alanine	1	0.41	>0.05 ^{n.s}	1.09	>0.05 ^{n.s}	0.53	>0.05 ^{n.s}
Arginine	1	1.31	>0.05 ^{n.s}	0.21	>0.05 ^{n.s}	1.21	>0.05 ^{n.s}
Asparagine	1	1.03	>0.05 ^{n.s}	0.57	>0.05 ^{n.s}	0.68	>0.05 ^{n.s}
Aspartic acid	1	0.42	>0.05 ^{n.s}	1.15	>0.05 ^{n.s}	9.30	≤0.05 *
Cystine	1	0.07	>0.05 ^{n.s}	1.38	>0.05 ^{n.s}	0.43	>0.05 ^{n.s}
Glutamic acid	1	0.57	>0.05 ^{n.s}	1.55	>0.05 ^{n.s}	0.07	>0.05 ^{n.s}
Glutamine	1	0.13	>0.05 ^{n.s}	1.24	>0.05 ^{n.s}	0.72	>0.05 ^{n.s}
Glycine	1	0.01	>0.05 ^{n.s}	0.87	>0.05 ^{n.s}	1.58	>0.05 ^{n.s}
Hydroxyproline	1	0.23	>0.05 ^{n.s}	2.23	>0.05 ^{n.s}	0.00	>0.05 ^{n.s}
Histidine	1	0.03	>0.05 ^{n.s}	2.63	>0.05 ^{n.s}	2.24	>0.05 ^{n.s}
Isoleucine	1	1.45	>0.05 ^{n.s}	2.81	>0.05 ^{n.s}	6.16	≤0.05 *
Leucine	1	0.71	>0.05 ^{n.s}	0.87	>0.05 ^{n.s}	4.34	>0.05
Lysine	1	0.77	>0.05 ^{n.s}	0.00	>0.05 ^{n.s}	1.28	>0.05 ^{n.s}
Methionine	1	0.30	>0.05 ^{n.s}	0.01	>0.05 ^{n.s}	0.74	>0.05 ^{n.s}
Phenylalanine	1	1.56	>0.05 ^{n.s}	0.14	>0.05 ^{n.s}	7.22	≤0.05 *
Proline	1	0.40	>0.05 ^{n.s}	3.05	>0.05 ^{n.s}	2.57	>0.05 ^{n.s}
Serine	1	1.38	>0.05 ^{n.s}	4.06	>0.05 ^{n.s}	4.29	≤0.05 ^{n.s}
Threonine	1	0.07	>0.05 ^{n.s}	6.68	≤0.05 *	4.18	≤0.05 *
Tryptophan	1	0.74	>0.05 ^{n.s}	0.01	>0.05 ^{n.s}	0.11	>0.05 ^{n.s}
Tyrosine	1	0.40	>0.05 ^{n.s}	11.88	≤0.05 *	11.31	≤0.05 *
Valine	1	0.29	>0.05 ^{n.s}	1.22	>0.05 ^{n.s}	2.59	>0.05 ^{n.s}

*Significant differences at $p < 0.05$, ^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

CHAPTER 4

Adult trees *Cryptomeria japonica* (Thunb. ex L.f.) D. Don

**Micropropagation: Factors Involved in the Success of the
Process**

SUPPLEMENTARY MATERIALS CHAPTER 4

Table S1. Variations of basal DCR (Gupta and Durzan, 1985), MS (Murashige and Skoog, 1962) and QL (Quoirin and Lepoivre, 1977) media tested at different stages of *Cryptomeria japonica* micropropagation process, Experiment 1.

Stage	Basal medium	PGRs (μM)	Sucrose (w/v)	AC (g L^{-1})	Agar (g L^{-1})
Induction	DCR	BA (8.8)	3%	-	Difco Agar [®] granulated (8)
Induction	MS	BA (8.8)	3%	-	Difco Agar [®] granulated (8)
Induction	QL	BA (8.8)	3%	-	Difco Agar [®] granulated (8)
Elongation	DCR	-	3%	(2)	Difco Agar [®] granulated (8.5)
Elongation	MS	-	3%	(2)	Difco Agar [®] granulated (8.5)
Elongation	QL	-	3%	(2)	Difco Agar [®] granulated (8.5)
Rooting	Half strength macronutrients QL	NAA (50)	3%	-	Difco Agar [®] granulated (8)
Rooting	Half strength macronutrients QL	NAA (50)	1.5%	-	Difco Agar [®] granulated (8)
Root expression	Half strength macronutrients QL	-	3%	(2)	Difco Agar [®] granulated (8.5)
Root expression	Half strength macronutrients QL	-	1.5%	(2)	Difco Agar [®] granulated (8.5)

PGRs, plant growth regulators; BA,6-benzyladenine; NAA, 1-naphthalenacetic acid; AC, activated charcoal.

Table S2. Variations of basal QL (Quoirin and Lepoivre, 1977) medium tested at different stages of *Cryptomeria japonica* micropropagation process, Experiment 2.

Stage	Basal medium	PGRs (μM)	Sucrose (w/v)	AC (g L^{-1})	Agar (g L^{-1})
Induction	QL	BA (8.8)	3%	-	Difco Agar [®] granulated (8)
Induction	QL	m-T (8.8)	3%	-	Difco Agar [®] granulated (8)
Induction	QL	TDZ (8.8)	3%	-	Difco Agar [®] granulated (8)
Elongation	QL	-	3%	(2)	Difco Agar [®] granulated (8.5)
Rooting	Half strength macronutrients QL	NAA (50)	1.5%	-	Difco Agar [®] granulated (8)
Root expression	Half strength macronutrients QL	-	1.5%	(2)	Difco Agar [®] granulated (8.5)

PGRs, plant growth regulators; BA,6-benzyladenine; m-T, meta-topolin; thidiazuron, TDZ; NAA, 1-naphthalenacetic acid; AC, activated charcoal.

Table S3. Statistical analysis for the survival (%) showed in *Cryptomeria japonica* per explant type (apical explants of >1.5 cm in length, apical explants of <1.0 cm length, basal explants of >1.5 cm length, and basal explants of <1.0 cm in length) and basal media (DCR [Gupta and Durzan, 1985], MS [Murashige and Skoog, 1962] and QL [Quoirin and Lepoivre, 1977]).

Source	df	Survival (%)	
		χ^2 Test	p-Value
Explant type (E)	3	492.11	<0.05*
Basal media (B)	2	490.60	≥ 0.05 ^{n.s}
B x C	6	490.60	≥ 0.05 ^{n.s}

* Significant differences at $p < 0.05$, ^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S4. Statistical analysis for shoot induction (%) and number of shoots per explant showed in *Cryptomeria japonica* per explant type (apical explants of >1.5 cm in length, apical explants of <1.0 cm length, basal explants of >1.5 cm length, and basal explants of <1.0 cm in length) and basal media (DCR [Gupta and Durzan, 1985], MS [Murashige and Skoog, 1962] and QL [Quoirin and Lepoivre, 1977]).

Source	df	Shoot induction (%)		N° shoots / explant	
		χ^2 Test	p-Value	F Test	p-Value
Explant type (E)	3	266.95	<0.05*	4.31	<0.05*
Basal media (B)	2	266.00	≥ 0.05 ^{n.s}	9.09	<0.05*
E x B	6	256.35	≥ 0.05 ^{n.s}	2.59	<0.05*

* Significant differences at $p < 0.05$, ^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S5. Statistical analysis for root induction (%), number roots per explant and length of longest of *Cryptomeria japonica* shoots cultured in QL medium (Quoirin and Lepoivre, 1977), supplemented with 3% (w/v) sucrose or 1.5% (w/v) sucrose, according to light treatment.

Source	df	Root induction (%)		N° roots / explant		Length of longest root	
		χ^2 Test	p-Value	F Test	p-Value	F Test	p-Value
Sucrose concentration (S)	1	152.78	≥ 0.05 ^{n.s}	1.38	<0.05 ^{n.s}	4.13	<0.05*
Light treatment (L)	1	149.44	≥ 0.05 ^{n.s}	22.70	<0.05*	4.8	<0.05*
S x L	1	149.44	≥ 0.05 ^{n.s}	0.01	≥ 0.05 ^{n.s}	0.30	≥ 0.05 ^{n.s}

* Significant differences at $p < 0.05$, ^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S6. Statistical analysis for the survival (%) of rooted shoots propagated *in vitro* coming from *Cryptomeria japonica* adult trees, after four weeks under *ex vitro* conditions.

Source	df	Survival (%)	
		χ^2 Test	p-Value
Sucrose concentration (S)	1	67.94	≥ 0.05 ^{n.s}
Light treatment (L)	1	67.07	≥ 0.05 ^{n.s}
S x L	1	60.17	<0.05*

* Significant differences at $p < 0.05$, ^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S7. Statistical analysis for the survival (%) showed in *Cryptomeria japonica* explants (basal explants of >1.5 cm length) cultured in QL medium (Quoirin and Lepoivre, 1977), supplemented with 6-benzyladenine (BA), meta-Topolin (m-T) and thidiazuron (TDZ) (8.8 μ M).

Source	df	Survival (%)	
		X ² Test	p-Value
Cytokinin type	2	703.95	≥ 0.05 ^{n.s}

^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S8. Statistical analysis for shoot induction (%) and number of shoots per explant showed in *Cryptomeria japonica* explants (basal explants of >1.5 cm length) cultured in QL medium (Quoirin and Lepoivre, 1977), supplemented with 6-benzyladenine (BA), meta-Topolin (m-T) and thidiazuron (TDZ) (8.8 μ M).

Source	df	Shoot induction (%)		N ^o shoots / explant	
		X ² Test	p-Value	F Test	p-Value
Cytokinin type (E)	1	405.51	<0.05*	0.19	≥ 0.05 ^{n.s}

*Significant differences at $p < 0.05$, ^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S9. Statistical analysis for root induction (%), number root per explant and length of longest root showed in *Cryptomeria japonica* explants (basal explants of >1.5 cm length) cultured in QL medium (Quoirin and Lepoivre, 1977), supplemented with 50 μ M 1-naphthaleneacetic acid (NAA), according to cytokinin type [6-benzyladenine (BA), meta-Topolin (m-T) and Thidiazuron (TDZ)] (8.8 μ M).

Source	df	Root induction		N ^o root / explant		Length of longest root	
		X ² Test	p-Value	F Test	p-Value	F Test	p-Value
Cytokinin type	1	118.52	<0.05*	0.70	≥ 0.05 ^{n.s}	0.49	≥ 0.05 ^{n.s}

*Significant differences at $p < 0.05$, ^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S10. Statistical analysis for the survival (%) showed in *Cryptomeria japonica* explants (basal explants of >1.5 cm length) cultured in QL medium (Quoirin and Lepoivre, 1977), supplemented with 50 μ M 1-naphthaleneacetic acid (NAA), according to cytokinin type [6-benzyladenine (BA) and meta-Topolin (m-T) (8.8 μ M)].

Source	df	Survival (%)	
		X ² Test	p-Value
Cytokinin type	1	14.53	≥ 0.05 ^{n.s}

^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

CHAPTER 5

**Testing Explant Sources, Culture Media, and Light Conditions
for the Improvement of Organogenesis in *Pinus ponderosa* P.**

Lawson and C. Lawson

SUPPLEMENTARY MATERIALS CHAPTER 5

Table S1. Statistical analysis for the survival (%) showed in explants of *Pinus ponderosa* per 6-benzyladenine (BA) (4.4, 22 and 44 μ M) and cultured media (LP or half LP macronutrients) [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)].

Source	df	Survival (%)	
		χ^2 Test	p-Value
BA concentration (B)	2	600.83	$\leq 0.05^*$
Culture medium (C)	1	608.99.	$> 0.05^{n.s}$
B x C	1	597.87	$> 0.05^{n.s}$

* Significant differences at $p < 0.05$, ^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S2. Statistical analysis for the survival (%) showed in explants of *Pinus ponderosa* per explant type (cotyledons or whole zygotic embryos) and sterilization protocol [5% commercial bleach and 10% H₂O₂ (v/v)].

Source	df	Survival (%)	
		χ^2 Test	p-Value
Explant type (E)	1	596.21	$\leq 0.05^*$
Sterilization protocol (S)	1	593.11	$> 0.05^{n.s}$
E x S	1	592.96	$> 0.05^{n.s}$

* Significant differences at $p < 0.05$, ^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S3. Statistical analysis for the explants forming shoots (%) showed in explants of *Pinus ponderosa* per 6-benzyladenine (BA) (4.4, 22 and 44 μ M), culture media (LP or half LP macronutrients) [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)], explant type (cotyledons or whole zygotic embryos) and sterilization protocol [5% commercial bleach and 10% H₂O₂ (v/v)].

Source	df	Explants forming shoots (%)	
		χ^2 Test	p-Value
BA concentration (B)	2	496.36	$> 0.05^{n.s}$
Cultured media (C)	1	498.04	$\leq 0.05^*$
Explant type (E)	1	534.93	$\leq 0.05^*$
Sterilization protocol (S)	1	534.93	$> 0.05^{n.s}$
E x S	1	496.36	$> 0.05^{n.s}$
E x C	1	454.59	$\leq 0.05^*$
C x S	1	453.84	$> 0.05^{n.s}$
B x E	2	453.54	$> 0.05^{n.s}$
B x S	2	452.56	$> 0.05^{n.s}$
B x C	2	452.18	$> 0.05^{n.s}$
E x C x S	1	448.81	$> 0.05^{n.s}$
B x E x S	2	448.19	$> 0.05^{n.s}$
B x C x E	2	443.74	$> 0.05^{n.s}$
B x C x S	2	441.34	$> 0.05^{n.s}$
B x C x E x S	2	436.19	$> 0.05^{n.s}$

* Significant differences at $p < 0.05$, ^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S4. Statistical analysis for the survival (%) showed in explants of *Pinus ponderosa* per cytokinin type and light treatment cultured in half LP macronutrients medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)].

Source	df	Survival (%)	
		χ^2 Test	p-Value
Cytokinin type (C)	1	29.206	$\leq 0.05^*$
Light treatment (C)	3	21.311	$\leq 0.05^*$
C x L	3	21.311	$> 0.05^{n.s.}$

*Significant differences at $p < 0.05$, ^{n.s.} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S5. Statistical analysis for the explants forming shoots (%) showed in explants of *Pinus ponderosa* cultured in half LP macronutrients medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with 6-benzyladenine (BA) and meta-Topolin (m-T) according to light treatment.

Source	df	Explants forming shoots (%)	
		χ^2 Test	p-Value
Cytokinin type (C)	1	393.51	$> 0.05^{n.s.}$
Light treatment (C)	3	392.40	$> 0.05^{n.s.}$
C x L	3	391.23	$> 0.05^{n.s.}$

^{n.s.} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S6. ANOVA for total number of shoots produced per initial explant showed in explants of *Pinus ponderosa* cultured in half LP macronutrients medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with 6-benzyladenine (BA) and meta-Topolin (m-T) (13.1 μ M) according to light treatment.

Source	df	N° shoots / explant	
		F Test	p-Value
Cytokinin type (C)	1	0.31	$> 0.05^{n.s.}$
Light treatment (C)	3	1.00	$> 0.05^{n.s.}$
C x L	3	1.07	$> 0.05^{n.s.}$

^{n.s.} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S7. ANOVA for shoots elongated sufficiently to be rooted (%) showed in explants of *Pinus ponderosa* cultured in half LP macronutrients medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] supplemented with 6-benzyladenine (BA) and meta-Topolin (m-T) (13.1 μ M) according to light treatment.

Source	df	Shoots elongated sufficiently to be rooted (%)	
		F Test	p-Value
Cytokinin type (C)	1	0.02	$> 0.05^{n.s.}$
Light treatment (C)	3	3.94	$\leq 0.05^*$
C x L	3	0.21	$> 0.05^{n.s.}$

*Significant differences at $p < 0.05$, ^{n.s.} Non-significant at $p < 0.05$, df Degrees of freedom.

Table S8. Statistical analysis for root induction (%), number root per explant and length of longest root showed in explants of *Pinus ponderosa* cultured in half LP macronutrients medium [Quoirin and Lepoivre (1977) modified by Aitken-Christie et al. (1988)] according to light treatment.

Source	df	Root induction		N° root /explant		Length of longest root	
		X ² Test	p-Value	F Test	p-Value	F Test	p-Value
Light treatment (L)	3	132.79	>0.05 ^{n.s}	0.66	>0.05 ^{n.s}	0.42	>0.05 ^{n.s}

^{n.s} Non-significant at $p < 0.05$, df Degrees of freedom.

Development of biotechnological tools for the *in vitro* propagation of elite trees for the Basque Country forest

Alejandra Rojas Vargas