



IODP-INVEST
WHITE PAPER

**NICOYA, COSTA RICA:
AN IODP TARGET WHERE THE SEISMOGENIC ZONE OF LARGE EARTHQUAKES
CAN BE DRILLED AND INSTRUMENTED**

Marino Protti¹, Victor González¹, Yoshiyuki Kaneda² and Kevin Brown³

(1) Costa Rica Volcanological and Seismological Observatory

Universidad Nacional

Apartado 1718-3000, Heredia, Costa Rica

jprotti@una.ac.cr

(2) IFREE-JAMSTEC

(3) SCRIPPS-UCSD

ABSTRACT

The Nicoya segment of Middle America Trench has generated large earthquakes in 1883, 1900 and 1950. Since the Nicoya peninsula sits right over the seismogenic zone, it has proved to be a unique site for near field monitoring of interplate interactions. Recent crustal deformation data indicates that locking on this plate boundary starts as close as 30 km from the trench where the plate interface is around 6 km deep. These results make this subduction segment an attractive target for drilling the plate interface and seismogenic zone at several different depths in a margin where collected data suggests is close to rupture. This temporal proximity to failure gives also a chance to monitor several parameters (such as fluid flow, geochemistry, strain, temperature, tilt, among others) in holes at several distances from the trench prior to rupture, possibly during rupture and most of the changes induced by postseismic slip.

This area is part of the target region of both, the Seismogenic Zone Experiment (SEIZE) and the Subduction Factory (SUBFAC) initiatives of MARGINS and it has been studied for many years with different techniques (high resolution marine seismics, refraction profiles, broad band land and ocean seismic stations, scan mapping, an ALVIN cruise, ODP and IODP legs, CGPS, electronic tiltmeters), so there is already sufficient information to start an IODP proposal. The on-land recorded deformation rates are very high precluding success in recording deformation offshore.

Submitted to the following working groups of the upcoming INVEST meeting (IODP New Ventures in Exploring Scientific Targets) in Bremen, September 23-25, 2009:

WG2.5: Subduction zones and volcanic arcs.

WG5.1: Geohazards: earthquakes.

WG6.1: Observatories.

WG6.6: Balancing long-term projects and single expeditions.

NICOYA, COSTA RICA: AN IODP TARGET WHERE THE SEISMOGENIC ZONE OF LARGE EARTHQUAKES CAN BE DRILLED AND INSTRUMENTED

Nicoya is a segment of the subduction zone at Middle American Trench. Here the Cocos plate subducts at almost 90 mm/yr under the Caribbean plate (Fig. 1). With the exception of the Nicoya segment, the entire subduction zone along the Middle American Trench has ruptured with large earthquakes in the last 60 years. Nicoya had large earthquakes ($M_w > 7$) in 1853, 1900 and in 1950. Therefore the Nicoya segment constitutes a mature seismic gap (Protti et al., 2001).




Figure 1. Tectonic setting of the Nicoya, Costa Rica, subduction segment. Solid circles show the location of large ($M_w > 7$) earthquakes in Costa Rica. Solid triangles represent active volcanoes. Lines parallel to the trench are isodepth contours of the top of the Wadati-Benioff zone in 20 km intervals. QSC is the Quesada Sharp Contortion, a tear fault in the subducted Cocos slab.

The Nicoya subduction segment has two important advantages over most subduction zones in the world. First it has a peninsula sitting right over the seismogenic zone of large ($M_w > 7$) earthquakes that allows recording crustal deformation in the very near field with geodetic techniques such as continuous GPS (CGPS) and tiltmeters, as well as with seismic instrumentation. The second advantage is that the upper limit of the seismogenic zone is reachable by drilling with the current drilling vessels capabilities.

Geodetic and seismic instrumentation have recorded two slow slip events on the Nicoya segment, one, in September 2003, believed to have occurred within the seismogenic zone (Protti et al., 2004) and another in May 2007 that went from the upper limit to the downdip limit of the

seismogenic zone (Psencik et al., 2008). According to Brown et al., 2005 and LaBonte et al., 2009, slow slip events have also been recorded, in this segment, near the trench through transient fluid pulsing and seismic tremor, demonstrating the importance of continuous borehole monitoring close to the trench.

Recent crustal deformation data from the Nicoya peninsula indicates that locking on this plate boundary starts as close as 30 km from the trench (Norabuena et al 2004) where the plate interface is around 6 km deep. These results make this subduction segment an attractive target for drilling the plate interface and seismogenic zone at several different depths in a margin where collected data suggests is close to rupture.




Figure 2. The Nicoya segment of the Middle American Subduction zone (DeShon, et. al., 2005). Color pattern shows the quality of locking along the plate interface (Norabuena et. al., 2004) with the strongest patch where the plate interface is from 5 to 15 km deep (the plate interface is represented by a red line labeled 5 km and solid black lines labeled 10, 20 30, 40 and 50). Orange circles represent interplate seismicity. Temperature isotherms from *Spinelli and Saffer* (2004) updip and *Harris and Wang* (2002) downdip; the grey box indicates the potential variance in

the modeled 300°C isotherm. Yellow solid lines mark the 500 m and 2500 m depth contours, which is the operational water depth interval for Chikyu. Within this depth range Chikyu could target the seismogenic zone at 6 to 7 km.

WHY NICOYA?

IT GENERATES LARGE EARTHQUAKES
SHORT RECURRENCE TIMES
CLOSE TO FAILURE
FAST DEFORMATION RATES
POLITICALLY EASY
LOTS OF DATA AVAILABLE
SMALL AREA (10s of km rather than 100s of km)
LAND CLOSE BY FOR LOGISTICS AND DATA LINKS
THE SEISMOGENIC ZONE IS REACHABLE BY DRILLING

TWO WAYS AND TIMES YOU CAN DO IT:

NOW AND CHEAP
NON-RISER HOLES WITH IN-SITU RECORDING SYSTEMS
LATER AND WITH MORE RESOURCES
CHIKIU DRILLING AND OCEAN BOTTOM CABLE NETWORKS

References

- Brown, K., et al. (2005), Correlated transient fluid pulsing and seismic tremor in the Costa Rica subduction zone, *Earth and Planetary Science Letters*, 238, 189-203.
- DeShon, H. R.; S. Schwartz, A. Newman, V. González, M. Protti, L. M. Dorman, T. H. Dixon, E. O. Norabuena, E. R. Flueh; 2006; Seismogenic zone structure beneath the Nicoya Peninsula, Costa Rica, from 3D local earthquake P- and S-wave tomography; *Geophysical Journal International*; Vol 164, issue 1, January.
- Harris, R.N. and K. Wang (2002), Thermal models of the Middle America trench at the Nicoya Peninsula, Costa Rica, *Geophys. Res. Letts.*, 29, 2010, doi:10.1029/2002GL015406.
- LaBonte, A.L. Brown, K.M. Fialko Y., Hydrologic detection and finite-element modeling of a slow-slip event in the Costa Rica prism toe, *J Geophys. RESEARCH*, VOL. 114, B00A02, doi:10.1029/2008JB005806, 2009.
- Newman, A.; S. Schwartz, V. Gonzalez , H. DeShon, M. Protti, and L. Dorman; Along-strike variability in the seismogenic zone below Nicoya Peninsula, Costa Rica; *Geophysical Research Letters*; Vol. 29, No. 20, 2002.
- Norabuena, E., T.H. Dixon, S.Y. Schwartz, H.R. DeShon, M. Protti, L. Dorman, E.R. Flueh, V. González, P. Lundgren, A. Newman, F. Pollitz, and D. Sampson 2004 Geodetic and seismic constraints on some seismogenic zone processes in Costa Rica, *J. Geophys. Res.*, 109, B11403, doi: 10.1029/2003JB002931.
- Protti, M., F. Güendel, and E. Malavassi (2001), *Evaluación del Potencial Sísmico de la Península de Nicoya*, Editorial Fundacion UNA, Heredia, Costa Rica, 144 p.
- Protti, M; V. González; T. Kato; T. Iinuma; S. Miyasaki; K. Obana; Y. Kaneda; P. LaFemina; T. Dixon; S. Schwartz; A Creep Event on the Shallow Interface of the Nicoya Peninsula,

Presented before the INVEST meeting (IODP New Ventures in Exploring Scientific Targets) in Bremen, Germany, September 23-25, 2009.

Costa Rica Seismogenic Zone; *Eos Trans. AGU*, 85(47), Fall Meet. Suppl., Abstract S41D-07; 2004.

Psencik, K, Dixon, T H, Schwartz, S, Walter, J, Protti, M, Gonzalez, V, 2008, Updated Episodic Tremor and Slip on the Cocos-Caribbean Subduction zone as measured by a GPS and Seismic Network on the Nicoya Peninsula, Costa Rica, *Eos Trans. AGU*, 89(53), Fall Meet. Suppl., Abstract U33A-0044.

Spinelli, G.A. and D. Saffer (2004), Along-strike variations in underthrust sediment dewatering on the Nicoya margin, Costa Rica related to the updip limit of seismicity, *Geophys. Res. Letts.*, 31, 4613, doi:10.1029/2003GL018863.